

The effect of gyre-scale property changes on the North Atlantic overturning over the last 50 years




Vassil Roussenov¹, Richard Williams¹, M. Susan Lozier² and Doug Smith³

1. Department of Earth and Ocean Sciences, University of Liverpool, UK 2. Duke University, Durham, NC 27708, USA 3. Met Office, Exeter, UK



1. Historical data analysis

Two independent hydrographic data sets (HYDROBASE and Met Office) have been used to assess the changes in the temperature, salinity and density over the last 50 years. HYDROBASE is based on hydrographic station data binned into 1°x1° grid for the North Atlantic T/S and averaged over two 20 year periods: 1950-1970 and 1980-2000 (Lozier, et al., 2010). The Met Office data set is compiled by an advanced objective analysis of ocean temperature and salinity using covariances from a global climate model (Smith and Murphy, 2007). It covers the period 1950 to 2009 with monthly temporal resolution.




The long term property changes show:

- freshening and cooling in the subtropical gyre; warmer and saltier waters in the subtropical gyre
- density compensation of T and S changes reflected by reduced density changes
- T and S are not completely compensated leading to a decrease of density in 1980-2000
- T/S changes in the Met Office data set show similar gyre confined and opposing signals with slightly reduced amplitudes and smoother patterns
- SSH changes comparable with reconstructions of sea level rise between 1950 and 2000 of typically 3 mm yr⁻¹ along 40° N (Church et al., 2004)


2. Model set-up and test experiments

The overturning is diagnosed using a dynamical adjustment in the MITGCM (global 1°x1° set-up and 23 vertical levels). The model is initialised from gridded T/S data and integrated for few years to allow the density field to dynamically adjust and to spin-up the wind-driven circulation. The model is forced by monthly NCEP winds averaged for each 20 year period. Internal 3D T/S relaxation towards initial fields has been used with 3 years restoring timescale.

- For the test case we use 20 year mean T/S from GECCO (GECCO, Koehl et al., 2006) and compare the model MOC with the transports directly calculated from GECCO monthly velocities.
- Then we replace the North Atlantic GECCO T/S by HYDROBASE data for the two 20 year periods, repeat the runs and use the model transports to diagnose the MOC change.
- Same procedure has been repeated with the Met Office global data on annual basis, using interannually and monthly varying NCEP winds.




3. MOC changes based on HYDROBASE



Uncertainties in the MOC changes have been assessed by Bayesian-type perturbation experiments carried out by randomly perturbing initial fields using the density standard errors for the historical data. The ensemble mean of 80 random members is close to the estimate from the default run with confidence limits based on the standard deviation of the ensemble. The transport change in the subtropical gyre is typically - 2 Sv (± 1 Sv) and + 1 Sv (± 0.5 Sv) for the subtropical gyre. In the case of removed data points along the topography (interior test, green line), the model communicates the interior T/S along the boundaries, recovering west-east density contrast and leading to a transport change very close to the default case.

4. MOC variability estimated from the Met Office data



The long term MOC changes estimated from the Met Office data are in accord with the previous results with opposing signals in subtropical and subpolar gyres. The changes in the later period (2000 to 2009) show the opposite pattern: increase in the subtropical and decrease in the subpolar gyres. MOC changes, estimated on annual basis, show more complicated structures with annual amplitudes of 1 - 1.5 Sv superimposed on the long term decadal variations.

5. Conclusions

- data analysis shows different responses in the long term property changes in the subtropical/subpolar gyres: freshening and cooling in the subpolar gyre; warmer and saltier waters in the subtropical gyre
- density compensation of T and S changes reflected by reduced density changes
- T and S not completely compensated leading to a decrease of the density in 1980 - 2000
- model estimates of the long term MOC changes show weakening over the subtropical gyre by - 2 ± 1 Sv and slight increase over the subpolar gyre by + 1 ± 0.5 Sv
- MOC variations estimated on annual basis from Met Office data set have similar amplitudes, but show more complicated patterns and opposing changes in the latest period

References:

Lozier, MS, V Roussenov, MSC Reed and RG Williams 2010. On the spatial pattern of meridional overturning changes in the North Atlantic. *Nature Geoscience*, DOI: 10.1038/NGE094.

Smith, DM and JM Murphy, 2007. An objective ocean temperature and salinity analysis using covariances from a global climate model. *JGR*, 112, C02022.

