

A. Richter, A. Hilboll, A. Zien, and J. P. Burrows

Email: [Andreas.Richter@iup.physik.uni-bremen.de](mailto:Andreas.Richter@iup.physik.uni-bremen.de)

Institute of Environmental Physics/Remote Sensing, University of Bremen, FB 1, P.O. Box 330440, D-28334 Bremen, Germany


## Introduction

- satellite observations of trace gases such as O<sub>3</sub>, NO<sub>2</sub>, or SO<sub>2</sub> provide important information on tropospheric composition
- clouds interfere with the retrievals and need to be taken into account
- study of variation in NO<sub>2</sub> signal for different cloud situations can give an idea on the uncertainties introduced and also on the vertical NO<sub>2</sub> distribution

## Cloud Effects

- clouds **shield** the atmosphere below them from view
- clouds enhance visibility of trace gases above them (**albedo effect**)
- clouds enhance visibility of trace gases in their upper part (**multiple scattering**)
- the effect of clouds on the light path depends on cloud fraction, cloud height, NO<sub>2</sub> profile, surface reflectivity and aerosol loading
- the NO<sub>2</sub> vertical profile in the presence of clouds will differ from that under clear sky because of changes in photolysis, convection, uplifting in frontal systems
- the satellite observed signal is larger over clouds
- the interference from surface effects is smaller in cloudy scenes
- clouds can potentially also interfere with the spectral retrieval of NO<sub>2</sub>

## Instrument and Retrieval



### GOME-2 Instrument:

- launched on MetOp-A in October 2006
- data since January 2007
- 4 channel nadir viewing UV/visible spectrometer
- first in a series of three identical instruments
- 80 x 40 km<sup>2</sup> pixel size
- global coverage in 1.5 days
- 09:30 LT equator crossing

### DOAS Analysis:



- 425 - 497 nm fitting window
- spectral spike correction
- liquid water cross-section to remove interference from water absorption

### Stratospheric Correction:

- reference sector over the Pacific (180° - 220° E), no cloud screening

### Airmass Factors:


- no airmass factor was applied



**Figure top:** Comparison of GOME-2 NO<sub>2</sub> slant columns for September 2010 using different cloud selection criteria of 0.0-0.1 (left), 0.1-0.2 (middle) and 0.5-1.0 (right). As can be seen, there are only moderate differences between the first two, but much reduced tropospheric signals for the last case. Note the occurrence of sporadic hotspots over the oceans in cloudy measurements (transport and lightning). A reduction of values is observed for large cloud fractions also in unpolluted regions from shielding of the tropospheric background.


## NO<sub>2</sub> dependence on cloud fraction

### Approach:




- All data over an area was collected for 2009 and 2010 (unless noted otherwise) and grouped into cloud fraction bins
- very large scatter is observed even strongly polluted regions under clear conditions => atmospheric variability
- seasonal differences are observed over cities => less vertical mixing in winter leads to more shielding

### Pollution hotspots:




- rapid increase of tropospheric NO<sub>2</sub> slant columns for lower cloud fraction => most NO<sub>2</sub> close to the surface
- effect is most pronounced over hotspots, less over large polluted areas (e.g. Hongkong vs. East Central China)
- over biomass burning regions, smaller dependence on cloud fraction => elevated NO<sub>2</sub>

### Outflow regions:




- reduced cloud fraction dependence of NO<sub>2</sub> in outflow regions => mixture of transport above and below clouds
- inverted behaviour over outflow region west of Africa => NO<sub>2</sub> above low clouds
- as shown for region 5, individual events of very large NO<sub>2</sub> are found at all cloud fractions

## Clouds and transport events



- many export events are associated to clouds
- NO<sub>2</sub> must be above clouds or within their top part to be visible from space
- cloudy and clear observations sometimes represent very different situations

## Effect of simple cloud screening



**Figure:** Normalised cloud fraction dependence. Dashed line is extrapolation by eye to 0 cloud fraction, large open symbol is value for cloud screening with cf < 0.2

- the current IUP NO<sub>2</sub> product applies only cloud screening (cf < 20% FRESCO), no additional cloud correction
- from extrapolation of cloud fraction dependence, this leads to underestimations of 23% in summer and 16% in winter over an intense hotspot such as Hongkong
- errors will be smaller in most other regions
- for individual measurements, errors can be much larger
- uncertainties in cloud fraction also affect these estimates

## Conclusions

- clouds have large impact on tropospheric NO<sub>2</sub> columns over polluted regions
- retrieved slant columns are much larger for cloud free situations
- effect is largest over hotspots and in winter
- over export regions, results vary depending on altitude of clouds and NO<sub>2</sub> layer
- export events are often associated to clouds, leading to biases in results derived by combining both types of data
- the simplified approach of selecting data with cloud fraction < 20% instead of detailed cloud treatment leads on average to underestimations of about 20% for an intense hotspot such as Hongkong and less elsewhere

## Selected References

- Beirle et al., Impact of clouds on tropospheric trace gas retrievals, poster at the NO<sub>2</sub> workshop Utrecht, 2007
- Boersma, K. F., H. J. Eskes and E. J. Brinksma, Error Analysis for Tropospheric NO<sub>2</sub> Retrieval from Space, *J. Geophys. Res.* 109 D04311, doi:10.1029/2003JD003962, 2004
- Boersma, K. F., H. J. Eskes, E. W. Meijer, and H. M. Kelder, Estimates of lightning NO<sub>x</sub> production from GOME satellite observations, *Atmos. Chem. Phys.*, 5, 2311–2331, 2005
- Koelemeijer, R. B. A., P. Stammes, J. W. Hovenier, and J. F. de Haan, A fast method for retrieval of cloud parameters using oxygen A band measurements from the Global Ozone Monitoring Experiment, *J. Geophys. Res.* 106, 3475–3490, 2001
- Martin, R. V., K. Chance, D. J. Jacob, T. P. Kurosu, R. J. D. Spurr, E. Bucsela, J. F. Gleason, P. I. Palmer, I. Bey, A. M. Fiore, Q. Li, R. M. Yantosca and R. B. Koelemeijer, An improved retrieval of tropospheric nitrogen dioxide from GOME, *J. Geophys. Res.* 107, 4437–4456, 2002.
- Richter, A. and J. P. Burrows, Retrieval of Tropospheric NO<sub>2</sub> from GOME measurements, *Adv. Space Res.*, 29, 16673–1683, 2002.