Mediterranean agriculture: More efficient irrigation needed to compensate increases in future irrigation water requirements

Marianela Fader, Sinan Shi, Werner von Bloh, Alberte Bondeau, Wolfgang Cramer









Outline


- 1. Introduction & research questions
- 2. Main tool: LPJmL
- 3. Current irrigation water use and efficiency
- 4. Irrigation water needs under climate and demographic change
- 5. Take home messages







Introduction





EGU 2016

Marianela Fader






Research questions

- Irrigation water needs today & water-intensity of crops
- 2. Countries' potentials for saving water
- 3. Impact of climate change on irrigation requirements
- 4. Influence of demographic change and water scarcity in future water requirements






Modelling tool: LPJmL





Irrigated and rainfed agriculture, grasslands, natural vegetation

C, H₂O exchange



Photosynthesis ⇔ Water availability

- River discharge (Gerten et al., 2004; Biemans et al., 2009)
- Irrigation water requirements (Rost et al., 2008)
- Water consumption of crops (Fader et al., 2010)

(Sitch et al. 2003, Bondeau et al. 2007, Rost et al. 2008, Behringer et al., 2008, Fader et al., 2010, Schapfhoff et al. 2013, Fader et al. 2015a)







10 additional crops included in LPJmL

- Fruit trees
- Nut trees
- Citrus trees
- Olive trees
- Potatoes
- Vegetables
- Fodder grasses
- Date Palms
- Cotton
- Grapes





- 1. Land use input
- 2. Parametrisation
- 3. Dev. phenology approach
- 4. Dev. management module
 - 5. Calibration
 - 6. Validation








Current irrigation water needs and efficiencies

Irrigation water requirements today



Average: 2000-2009

Total irrigation water requirements by plants = 128 km^3 = net irrigation water requirements (NIR) Total water "lost" from source to plant = 95 km^3

Current water withdrawal from source = 223 km³ = gross irrigation water requirements (GIR)

Withdrawal using drip irrigation and optimising distribution systems = 143 km^3

That means saving potentials of ~35%




(Fader et al., 2016, HESS)

(i)


CC



National gross irrigation water requirements and saving potentials



Agricultural trees have higher water requirements



Crops water needs = green water consumption + NIR. Average: 2000-2009. Error bars show the maximum and minimum values for crop water needs published by FAO (1989).










Mediterranean irrigation water requirements under climate and demographic change

Change in net irrigation requirements (2080-2090)





0





>45

>15



EGU 2016

NIR Change %

<-45 <-15

0

Marianela Fader

(Fader et al., 2016, HESS)


Ð

BY

(cc)



Southern Mediterranean strongest affected by population growth





EGU 2016

(Fader et al., 2016, HESS)

)ED

Ð


BY

(cc)

Will they have enough water?

- Scenarios of water availability, irrigation technology, efficiency of conveyance systems, management of aquatic ecosystems, climate and population growth.
- Algeria, Libya, Israel, Jordan, Lebanon, Syria, Serbia, Morocco, Tunisia and Spain have problems to meet irrigation water requirements in some scenarios (2080-2090).
- That is 10 out of 22 Mediterranean countries!





(Fader et al., 2016, HESS)

(i)

CC



Take home messages

Agricultural trees essential for simulation of irrig. requirements & carbon

Climate change will increase irrigation requirements (4-18%) Mediterranean region could save 35% of water

4

More efficient irrigation may partially compensate those increases

5

Population growth may increase irrig. req. even stronger (additionally 56%)

Water scarcity may pose problems for irrigated agriculture in 10 out of 22 countries

Ð

CC



EGU 2016

Marianela Fader



More details here ③

Geoscientific Model Development

Geosci. Model Dev., 8, 3545–3561, 2015 www.geosci-model-dev.net/8/3545/2015/ doi:10.5194/gmd-8-3545-2015 © Author(s) 2015. CC Attribution 3.0 License.

Modelling Mediterranean agro-ecosystems by including agricultural trees in the LPJmL model

M. Fader¹, W. von Bloh², S. Shi³, A. Bondeau¹, and W. Cramer¹

¹Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale, Aix-Marseille Université, CNRS, IRD,
Avignon Université, Technopôle Arbois-Méditerranée, Bâtiment Villemin, BP 80, 13545 Aix-en-Provence CEDEX 04, France
²Potsdam Institute for Climate Impact Research, Telegraphenberg, 14473 Potsdam, Germany
³Research Software Development Group, Research IT Services, University College London, Podium Building (1st Floor),
Gower Street, London WC1E 6BT, UK

Correspondence to: M. Fader (marianela.fader@imbe.fr)

Received: 1 June 2015 – Published in Geosci. Model Dev. Discuss.: 30 June 2015 Revised: 13 October 2015 – Accepted: 14 October 2015 – Published: 5 November 2015

Abstract. In the Mediterranean region, climate and land use change are expected to impact on natural and agricultural ecosystems by warming, reduced rainfall, direct degradation of ecosystems and biodiversity loss. Human population growth and socioeconomic changes, notably on the eastHydrol. Earth Syst. Sci., 20, 953–973, 2016 www.hydrol-earth-syst-sci.net/20/953/2016/ doi:10.5194/hess-20-953-2016 © Author(s) 2016. CC Attribution 3.0 License.

ample assessments of the influence of change impacts. Mediterranean irrigation under climate change: more efficient irrigation needed to compensate for increases in irrigation water requirements

M. Fader^{1,2}, S. Shi^{2,4}, W. von Bloh³, A. Bondeau¹, and W. Cramer¹

¹Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale, Aix-Marseille Université, CNRS, IRD, Avignon Université, Technopôle Arbois-Méditerranée, Bâtiment Villemin, BP 80, 13545 Aix-en-Provence CEDEX 4, France ²Laboratory of Excellence OT-Med. Europôle Méditerranéen de l'Arbois, Bâtiment Gérard Mégie, Avenue Louis Philibert, 13857 Aix-en-Provence CEDEX 3, France

Hydrology and Earth System

Sciences

 ³Potsdam Institute for Climate Impact Research, Telegraphenberg, Building A31, 14473 Potsdam, Germany
⁴Research Software Development Group, Research IT Services, University College London, Podium Building, 1st Floor, 1 Eversholt Street, London, NW1 2DN, UK

Correspondence to: M. Fader (marianela.fader@imbe.fr)

More details here 🙂

Geosci. Model Dev., 8, 3545-3561, 2015 www.geosci-model-dev.net/8/3545/2015/ doi:10.5194/gmd-8-3545-2015 © Author(s) 2015. CC Attribution 3.0 License.

Geoscientific Model Development EĞU

Modelling Mediterranean agro-ecosystems by including agricultural trees in the LP.ImL model

M. Fader¹, W. von Bloh², S. Shi³, A. Bondeau¹, and W. Cramer¹

¹Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale, Aix-Marseille Université, CNRS, IRD, Avignon Université, Technopôle Arbois-Méditerranée, Bâtiment Villemin, BP 80, 13545 Aix-en-Provence CEDEX 04, France ²Potsdam Institute for Climate Impact Research, Telegraphenberg, 14473 Potsdam, Germany ³Research Software Development Group, Research IT Services, University College London, Podium Building (1st Floor), Gower Street, London WC1E 6BT, UK

Correspondence to: M. Fader (marianela.fader@imbe.fr)

Received: 1 June 2015 - Published in Geosci. Model Dev. Discuss.: 30 June 2015 Revised: 13 October 2015 - Accepted: 14 October 2015 - Published: 5 November 2015

Abstract. In the Mediterranean region, climate and land use change are expected to impact on natural and agricultural ecosystems by warming, reduced rainfall, direct degradation of ecosystems and biodiversity loss. Human population growth and socioeconomic changes, notably on the east-







nstitut méditerranéen de biodiversité et d'écologie Aix*Marseille

Hydrol. Earth Syst. Sci., 20, 953-973, 2016 www.hydrol-earth-syst-sci.net/20/953/2016/ doi:10.5194/hess-20-953-2016 © Author(s) 2016. CC Attribution 3.0 License.

Thank you for your attention!



¹Mediterranean irrigation under climate change: ample assessments of sitions, the influence of more efficient irrigation needed to compensate for change impacts. increases in irrigation water requirements

M. Fader^{1,2}, S. Shi^{2,4}, W. von Bloh³, A. Bondeau¹, and W. Cramer¹

¹Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale, Aix-Marseille Université, CNRS, IRD, Avignon Université, Technopôle Arbois-Méditerranée, Bâtiment Villemin, BP 80, 13545 Aix-en-Provence CEDEX 4, France ²Laboratory of Excellence OT-Med. Europôle Méditerranéen de l'Arbois, Bâtiment Gérard Mégie, Avenue Louis Philibert, 13857 Aix-en-Provence CEDEX 3, France

³Potsdam Institute for Climate Impact Research, Telegraphenberg, Building A31, 14473 Potsdam, Germany ⁴Research Software Development Group, Research IT Services, University College London, Podium Building, 1st Floor, 1 Eversholt Street, London, NW1 2DN, UK

Correspondence to: M. Fader (marianela.fader@imbe.fr)