Four decades of hindsight into a complex hydropower system

[EGU2018-16245]

José P. Matos[1], Pedro Manso[1], Bettina Schaefli[2], Benno Schwegler[3], Andres Fankhauser[3], Martin Seiler[3] and Anton J. Schleiss[1]

FlexSTOR

0 Goal

Explore ways to identify and quantitatively describe the modes of operation that take place in a complex hydropower system across several decades (1980 – 2014).

1 The KWO system, in Switzerland



Fig 1. The KWO system and estimated glacier cover in 1993 [1, 2].

- **The *Kraftwerke Oberhasli AG* (KWO) hydropower system (Fig. 1):**
  - a) 10 power plants,
  - b) 29 turbines,
  - c) 1368 MW.
  - d) 4 main reservoirs.

2 Data

A wealth of data was used:

- Daily series of all **water fluxes** in the system (since 1980).
- Daily series of **reservoir levels** (since 1980).
- Hourly series of **SPOT market prices** (since 2008).



Fig 2. Outflows from the system (a) and SPOT energy prices (b).

3 Visualization

Sankey plots provide an appealing way of visualizing and interpreting system operation states.

- The width of each arrow is proportional to the magnitude of the flux (in this case discharge).
- These plots can be produced automatically, enabling a practical view of the system.



Fig 3. Mean behavior of the KWO system (1980 to 2014).

4 Clustering techniques

How to **reduce the dimensionality** of the dataset so that it can be interpreted ?

- **It depends** on the question being addressed.



Fig 4. Illustration of how two clustering algorithms operate.

5 Seasonal operations

Seasonal clusters of daily operations can be found using the K-means algorithm.

Significant changes since the 1980's (pumped storage).



Fig 5. Seasonal clustering (K-means).



Fig 6. Evolution of seasonal operations (5-year averages).

6 Outlier operation modes

Using DBSCAN clustering one can quickly identify rare operation modes.

- All production through the Gadmental
- Run-of-the-river production
- Filling of the Grimsel reservoir
- Absorbing energy from the system through pumping



Fig 7. Outlier operations (DBSCAN).

7 Conclusions

- Sankey plots are effective tools to visualize the operation of complex hydropower systems.
- Clustering can be used to reduce the dimensionality and facilitate the interpretation of the operations of a complex hydropower system.
- KWO has undergone significant changes in its operation that can mostly be explained by energy market conditions (stable hydrology on average terms).

8 References and acknowledgements

[1] M. P. Bieri, Operation of Complex Hydropower Schemes and its Impact on the Flow Regime in the Downstream River System under Changing Scenarios. Thèse EPFL, n° 5433, 2012.
[2] S. Terrier et al., Impact du retrait glaciaire et adaptation du potentiel hydroélectrique dans les Alpes suisses, La Houille Blanche, 2015 (1).
[3] A. K. Jain, Data clustering: 50 years beyond K-means, Pattern Recognition Letters, 31 (1), 2010.
[4] J. Sander et al., Density-based clustering in spatial databases: the algorithm GDBSCN and its applications, Data Mining and Knowledge Discovery, 2 (2), 1998.
[5] J. P. Matos et al. Operation changes of a complex hydropower system over decades. SCCER-SoE Annual Conference 2017.

We are thankful for the support provided by KWO and *Innosuisse*, which provided the bulk of the funding through project 17902.3 PFIW-IW, within the framework of the Swiss Competence Center for Energy Research (SCCER) on Supply of Electricity (SoE).

ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE
SCCER SoE SWISS COMPETENCE CENTER for ENERGY RESEARCH SUPPLY of ELECTRICITY
UNIL | Université de Lausanne
KWO GRIMSELSTROM

(1) Laboratory of Hydraulic Constructions, École Polytechnique Fédérale de Lausanne, Switzerland. pedro.manso@epfl.ch
(2) Faculty of Geosciences and Environment, University of Lausanne, Switzerland.
(3) Kraftwerke Oberhasli AG, Innertkirchen, Switzerland.

EGU European Geosciences Union General Assembly 2018
Vienna | Austria | 8–13 April 2018