









Internal conductance of Scots pine varies with atmospheric moisture

Lenka Foltýnová^{1,2}*, Yann Salmon^{2,3}, Linda M. J. Kooijmans^{4,5}, Ivan Mammarella², Kukka-Maaria Erkkilä², Huilin Chen⁴, Kadmiel Maseyk⁶, Wu Sun⁷, Ulli Seibt⁷, Timo Vesala^{2,3,8} & Teemu Hölttä³

Introduction

Carbonyl sulfide (COS) measurements can be used to compute stomatal and internal conductance to COS of plants as COS uptake in leaves is a one-way flux and does not require light, in contrast to photosynthesis. The possibility to separate the stomatal conductance $(g_{s,cos})$ from the full conductance pathway leaves the ability to study the internal conductance to COS ($g_{i,cos}$). $g_{i,cos}$ comprises the combined mesophyll (g_m) , chloroplast membrane (g_{ch}) and biochemical $(g_{bch,CA})$ conductances of COS (Fig 1).

The main goal of this study was to identify environmental drivers of the internal conductance to COS which was never done before, especially using carbonyl sulfide flux data.




Fig. 1 Scheme of COS and CO₂ diffusion pathways


Conductances denoted in the scheme are: g_s – stomatal (stomata and intercellular spaces, grey area), g_m – mesophyll (cell wall, plasma membrane and cytosol), gch - chloroplast membrane, $g_{st} - chloroplast$ stroma, $g_{bch,CA} - biochemical$ capacity of carbonic anhydrase (CA) located at inner side of chloroplast membrane, where majority of COS is decomposed (Stimler et al. 2010). Red area represents the internal conductance of COS $(g_{i,COS})$. Part of CA is also located in stroma, and therefore gst also contributes in part to $g_{i,COS}$. CO₂ equivalent of $g_{i,COS}$ represents partial internal conductance to CO_2 (g_{ip,CO_2}). Yellow area represents internal conductance of CO_2 (g_{i,CO_2}), which, in addition to being a part of the stroma resistance, is equal to $g_{i,COS}$. The green area representing the biochemical activity of RuBisCO ($g_{bch,RuB}$) is not involved in $g_{i,CO2}$.

Methods

Station for Measuring Forest Ecosystem – Atmosphere Relations (SMEAR II) in Hyytiälä, Finland (61°51′ N, 24°17′ E, 181 m a.s.l.), which is dominated by Scots pine (Pinus sylvestris L.).

Dendrometers were used to measure xylem diameter changes, which has been shown to be proportional to changes in water potential (WPP).

Air temperature (Ta), relative humidity (RH), vapor pressure radiation measured.







deficit (VPD), water vapor fluxes Fig 2: Two automated leaf gas-exchange chambers were (F_{H2O}) and photosynthetically active installed at the top of the canopy in one Scots pine tree. were also The chambers were of a different type with different spacing of the needles.

Ball-Berry model (Ball et al. 1987) was used to obtain g_s to water vapor, which was then converted to g_s to COS ($g_{s,COS}$). The parameters of empirical slope and intercept were determined from g_s , which was computed using F_{H2O} and VPD during lower RH conditions, RH < 70% (Kooijmans et al. 2018).

With assumptions that boundary layer conductance is negligible (ventilated chambers) and concentration of COS inside chloroplasts is near zero (COS is fully destroyed), we computed the internal conductance of COS from COS flux (F_{COS}) , ambient COS concentration $(C_{a,COS})$ and $g_{s,COS}$ as follows (Seibt et al. 2010, Wehr et al.

$$g_{i,COS} = \frac{F_{COS} \cdot g_{s,COS}}{-C_{a,COS} \cdot g_{s,COS} - F_{CO}}$$

Results and discussion

Univariate dependencies of g_{i.Cos} on Ta, RH, PAR, VPD, WP and F_{H2O} were analysed to to up to 3^{rd} degree of combinations of all examine whether $g_{ip,CO2}$ is reflecting micrometeorological conditions, tree water status and transpiration rate.

We observed the strongest dependency on VPD and RH, weaker dependency is apparent also on Ta and WPP (in averages from both chambers). The $g_{ip,CO2}$ did not seem to reflect the F_{H2O} at all in chamber 2. There was a weak dependence observed in chamber 1, which is an artefact caused by filtering out the high RH data, done only in chamber 1 (Fig 3).




Fig 3: Univariate dependencies in chamber 1. R² - coefficient of determination of linear regression

Multivariate dependencies of all six variables were tested for significance. We considered nonlinear relationships with g_{ip.CO2}, interactions variables and used non-parametric regression method (MARS) to be sure that intercorrelated drivers would not increase explanatory strength artificially.




Fig 4: Relative contribution of each variable to explained variability of gip.CO2

The full set of variables explained 62% and 51% of variability in $g_{ip,CO2}$ in chambers 1 and 2 respectively. Ta, RH and $g_{s,CO2}$ solely can explain the most of variability in chamber 1 and 2 (47% and 42% respectively, Fig 4).

PAR, WPP and F_{H2O} are not important drivers of g_{ip,CO2}. VPD can be explained by Ta and **RH** by definition.

Less explained variability in chamber 2 **Tab 1:** Results of path analysis suggests that the differences in leaf arrangement had an impact on the response of leaves to environmental conditions.




Fig 5: Diagram of the path analysis

We employed the path analysis to determine the causality of the relationships obtained from single- and multivariate analysis - the dependence of g_{ip.CO2} on VPD.

It is not evident, whether the $g_{ip,CO2} \times VPD$ dependency is caused by the direct VPD impact or indirect through other variables, i.e. WPP or $g_{s,CO2}$ (Fig 5).

The direct effect of VPD is the strongest one (Tab 1), however, it is hard to explain the mechanism. It could be caused by leaf osmolality (as a function of water and sugar content) or water potential (our proxy might not be good enough).

Direct Indirect Indirect Indirect	aı
effect effect (g _{s,cos}) effect (WPP) effe	ect (R ²)
Chamber 1 0.33 0.10 0.10 0.53	3
Chamber 2 0.32 0.14 0.03 0.49	9

References

Ball, J. T., Woodrow, I. E. & Berry, J. A. A model predicting stomatal conductance and its contribution to the control of photosynthesis under different environmental conditions. In: Biggens, J. (Ed.), Progress in Photosynthesis Research, vol. IV. 221-224 (1987)

Seibt, U., Kesselmeier, J., Sandoval-Soto, L., Kuhn, U. & Berry, J. A. A kinetic analysis of leaf uptake of COS and its relation to transpiration, photosynthesis and carbon isotope fractionation. Biogeosciences 7, 333-341, doi:10.5194/ bg-7-333-2010 (2010)

Wehr, R. et al. Dynamics of canopy stomatal conductance, transpiration, and evaporation in a temperate deciduous forest, validated by carbonyl sulfide uptake. Biogeosciences 14, 389-401, doi:10.5194/bg-14-389-2017 (2017)

Kooijmans, L. M. J. et al. Influences of light and humidity on carbonyl sulfidebased estimates of photosynthesis. PNAS, online (2019)

Abbreviations

CA ... carbonic anhydrase C_{a.COS} ... ambient concentration of COS

F_{H2O} ... water vapor flux

F_{COS} ... carbonyl sulphide flux g_{bch.CA} ... biochemical capacity of CA

g_{bch.RuB} ... biochemical capacity of RuBisCO g_{ch} ... cloroplast membrane conductance

g_{i,COS} ... internal conductance to COS

g_{i,CO2} ... internal conductance to CO₂ g_{ip,CO2} ... partial internal conductance to CO₂

g_m ... mesophyll conductance ... stomatal conductance

g_{s.COS} ... stomatal conductance to COS

g_{st} ... chloroplast stroma conductance PAR ... photosynthetically active radiation

RH ... relative air humidity Ta ... air temperature VPD ... vapor pressure deficit

WPP ... leaf water potential proxy

Acknowledgements

Affiliations

¹ Global Change Research Institute, Czech Academy of Sciences, Brno, Czech Republic

⁴ Centre for Isotope Research, University of Groningen, Groningen, The Netherlands

- ² Institute for Atmospheric and Earth System Research/Physics, University of Helsinki, Finland
- ³ Institute for Atmospheric and Earth System Research/Forest Sciences, University of Helsinki, Finland
- ⁵ Meteorology and Air Quality, Wageningen University and Research Center, Wageningen, The Netherlands ⁶ School of Environment, Earth and Ecosystem Sciences, Open University, Milton Keynes, UK
 - Department of Atmospheric and Oceanic Sciences, University of California Los Angeles, Los Angeles, CA, USA
 - ⁸ Viikki Plant Science Centre, University of Helsinki, Helsinki, Finland

This work was supported by the Ministry of Education, Youth and Sports of the Czech Republic within the National Sustainability Programme I (NPU I), grant number LO1415. The financial support by the Academy of Finland Centre of Excellence (118780), Academy Professor projects (312571 and 282842), ICOS-Finland (281255), the ERC-advanced funding scheme (AdG 2016 Project number: 742798, Project Acronym: COS-OCS), and the European Union's Horizon 2020 research and innovation programme (654182) are acknowledged. We thank Juho Aalto, Janne Levula, and Helmi Keskinen for technical support during the field campaign. KME thanks the Vilho, Yrjö and Kalle Väisälä Foundation for its support.