Impacts of acidity on multiphase chemistry in aqueous particles and clouds

Thomas Schaefer¹, Andreas Tilgner¹, Havala O. T. Pye², V. Faye McNeill³ and Hartmut Herrmann¹

¹Leibniz Institute for Tropospheric Research (TROPOS), Atmospheric Chemistry Department (ACD), Leipzig, 04318, Germany

²Office of Research and Development, US Environmental Protection Agency, Research Triangle Park, NC, 27711, USA

³Department of Chemical Engineering, Columbia University, New York, NY, 10027, USA

TROPOS Leibniz Institute for

Tropospheric Research



Introduction

- Acidity of aqueous atmospheric solutions is a key parameter
- Drives the partitioning of semi-volatile acidic or basic trace gases and their linked aqueous-phase chemistry
- Aqueous-phase chemistry affectes the acidity of atmospheric aqueous phases, e.g. deliquesced aerosol particles, cloud and fog droplets



- Figure 1 illustrates the most important processes where acidity plays a role
- Feedbacks in acidity and chemistry have crucial implications: (i) tropospheric lifetime of air pollutants, hence air quality and atmospheric aerosol composition, (ii) deposition input into other terrestrial and oceanic ecosystems, (iii) the visibility, (iv) climate and (v) human health
- The present study outline impacts of acidity (i) on the hydration of organic carbonyl compounds and (ii) multiphase chemistry of dissociating organic compounds in aqueous particles and clouds
- A comprehensive literature survey presents the current state of knowledge of acidity dependencies

Influence of acidity on hydration constants

- Hydration processes are typically acid- or base-catalyzed
- Equilibrium constant K_{Hvd} describes the ratio of the diol and carbonyl fraction of the aldehyde or ketone
- Hydration of simple aldehydes and ketones and dicarbonyls is unaffected by pH
- Multifunctional carbonyl compounds that contain pH sensitive moieties, such as a-oxocarboxylic acids, are highly influenced by the pH



OH, NO₃ radical studies on aqueous-phase chemistry

- OH or NO₃ radicals can react either by H-atom abstraction, electron transfer or by addition to a double bond
- OH radicals usually react by H-atom abstraction with saturated compounds in the range of $k = 10^7 - 10^9 M^{-1} s^{-1}$
- Unsaturated and aromatic compounds react by OH radical addition in the range of $k = 10^9 - 10^{10} M^{-1} s^{-1}$



O₂ studies on aqueous-phase chemistry

- Ozone is an electrophilic and very selective oxidant for organic compounds
- Rather high acidity dependencies exist for phenolic compounds and hydroxylated benzoic acids



- Acidity effect on reactivity of OH with organics is generally small
- Exceptions are small organic acids such as fomic, acetic, oxalic and malonic acids, which are more dependent on the acidity of the surrounding environment
- NO₃ reactions of protonated saturated aliphatic carboxylic acids are in the range of $k = 10^4 - 10^6 M^{-1} s^{-1}$

Fig. 4: Reactivity ratios of different dissociating organics. The ratio of the dianion and protonated dicarboxylic acid is indicated by (di).



Fig. 3: Ratio of ozone reaction rate constants of deprotonated and protonated organics in aqueous solution.

References

Lopalco, A. et al., 2016 J. Pharm. Sci. 105, 664-672. Pocker, Y. et al., 1969 J. Phys. Chem. 73, 2879-2882. Rapf, R.J. et al., 2017a J. Phys. Chem. A 121, 8368-8379. Rapf, R.J. et al., 2017b J. Phys. Chem. A 121, 4272-4282. Reed Harris, A.E. et al., 2014 J. Phys. Chem. A 118, 8505-8516. Tilgner, A. et al., 2020 in prep. Tur'yan, Y. I. Croat. Chem. Acta 1999, 72, 13-24.

Summary

- Hydration processes of multifunctional carbonyl compounds, such as a-oxocarboxylic acids, are highly influenced by the pH-value
- OH radical reaction are less affected by the pH, with exception of small carboxylic acids such as formic acid and oxalic acid
- NO₃ and O₃ reactions with organic acids, are highly affected by the acidity of the surrounding aqueous phase