# Mobile or not mobile: exploring the linkage between deep mantle composition and early Earth surface mobility


- **Keely A. O'Farrell[1], Sean J. Trim[2] and Samuel L. Butler[2]**
  - **[1]University of Kentucky, [2]University of Saskatchewan**

Initial Conditions

- Isothermal, free-slip top/bottom with reflecting sidewalls
- Run forward for approximately 2 Gyr (t=0.01)



# Vary parameters in basal magma ocean

- $Ra_T=2x10^3$ and $\Delta\eta_T=10^6$ (with $\tau_{yield}=\infty$ and $\eta^*=10\text{-}6$)
- Primordial layer may contain a high concentration of radiogenic element, patches of low viscosity melt or metals
- Primordial layer has one of the following:
  - Increased internal heating (H=20)
  - Decreased viscosity ($\Delta\eta_C=10$)
  - Increased conductivity (k=2)
- The Earth's mantle was hotter in the past
- Low viscosity gives rise to a high mean $Ra_T$
- Model mean $Ra_T$ ranges between $10^8$ - $10^9$

# Convective Vigor

- Model mean $Ra_T$ ranges between $10^8$ - $10^9$
- Our models have high Rayleigh numbers (convective vigor)
- Early Earth should be hotte and thus more convective (ie higher Ra)



# Potential for surface yielding



# Non-Newtonian rheology



# Potential for surface yielding – non-newtonian

- Episodic-lid convection
- Compositional features increase peak lithosphere mobility
- Onset time of lithosphere mobility affected by primordial composition