



Aerospace Information Research Institute(AIR)
Chinese Academy of Sciences(CAS)

Evaluating crop water requirements and actual crop water use with center pivot irrigation system in Inner Mongolia of China

Peejush Pani^{1,2},

Li Jia¹, Massimo Menenti^{1,3}, Guangcheng Hu¹, Chaolei Zheng¹, Qiting Chen¹, Yelong Zeng^{1,2}

¹State Key Laboratory of Remote Sensing Science, Aerospace Information Research Institute, Chinese Academy of Sciences, Beijing 100101, China;

²University of Chinese Academy of Sciences, Beijing 100049, China;

³Faculty of Civil Engineering and Earth Sciences, Delft University of Technology, 2628, Delft, Netherlands



7 May, 2020

Pani, Jia, Menenti et al., EGU Assembly 2020



Background

■ Objective:

Assess **Irrigation Performance Indicator (IPI)** using high resolution satellite remote sensing


■ Definition:

- **Crop water requirement (CWR)**: water required for crops to grow under optimal situation with **un-restricted water supply**.
- **Actual crop water use (CWU)**: water used by crops growing under **actual scenario** of water supply including application losses.

■ New approach:

- To estimate (gross = crop water requirements + loss) and actual crop water.
- Integrate farmer's irrigation practice & **application losses** at farm-scale for **center pivot irrigation systems (CPIS)**
- **Sentinel-2 MSI** and **Landsat-8 OLI** with meteorological forcing data and soil moisture retrievals.

Components of CWU in CPIS




... Martin et al., 2013



www.aircas.ac.cn

Methodology: Framework




$GIWR$ = Gross irrigation water requirement (mm)

$GIWUIS$ = Gross irrigation water use by sprinkler irrigation (I_S)

Study Area

CPIS irrigated area in Inner Mongolia, China




Located in north-central **Inner Mongolia** autonomous region of China, extending between **41°59' N** to **42°01' N** latitude and **115°55' E** to **115°57' E** longitude.



Results: Crop Water Requirement (CWR)



- Daily CWR using k_c times reference ET shows **distinct variation** of CWR for **potato** and **wheat**.
- Clear and distinct stages of each crop is reflected in the k_c graphs based on **daily reconstructed NDVI** from **MSI** and **OLI** data following k_c -NDVI method.
- Daily **fluctuation** in the ET_c is because of variation in **meteorological condition**





Results: Daily Estimates of GIWU components


- GIWU using **fine resolution MSI & OLI multi-spectral remote sensing** for each CPIS field.
- E_T ranges between **6-7mm/day**; E_S between **1.5-2.5 mm/day**; whereas, I_S **0.5-2.5 mm/irrigation**; E_A **simulation model** between **0.2 - 4.0 mm/irrigation**.
- **Lines** = **daily** estimates whereas, discrete **Bars** = **Irrigation events**.
- ET_C (black dash line) > ET_a (red line) [$E_S + E_T$] estimates.




Results: Model Validation

- Estimated I_S shows a good correlation with measured values R^2 0.91 & 0.97 for wheat and potato respectively, with lesser RMSE over wheat (0.65) compared to potato (1.54)
- E_A estimated on (cross = 13th July; diamond=14th July) shows distinct trend because of varying daily meteorological condition and agrees highly with measured values (R^2 0.97).
- The ETMonitor estimates shows a good correlation with the Eddy covariance instrument measured ET.


Estimated vs Measured I_S




Estimated vs Measured E_A



Estimated $E_T + E_S$ vs Eddy Cov.
Potato





Estimated $E_T + E_S$ vs Eddy Cov.
Wheat



Results: Irrigation Performance Indicator

Growing season (May-Sept, 2019)



- Figure 1 (Potato) & 2 (wheat) shows V and percentage of estimated losses by CPIS.
- IPI1 shows that V is **insufficient** (1.61) to fulfill the requirement of CPIS.
- IPI2 shows CPIS has **performed better** with respect to the **actual use** over wheat (1.14) compared to potato (1.36) in fulfilling the requirement.
- Out of total water allocated, **23.7** & **25.4** % is **lost** over potato and wheat field, respectively.



Conclusion & Discussion

- The **current approach** of evaluating separately the gross and net irrigation water requirement and actual water use considering the water losses, is **very essential** in evaluating the performance of any irrigation system.
- **ETMonitor** along with **modified RS-Gash** and **modified Yasar's algorithm** can play a vital role in estimating the **actual water use** of irrigated agriculture.
- Combined use of **Sentinel-2 MSI level-2** and **Landsat-8 OLI level-2** multi-spectral images provide data at higher temporal frequency to construct a continuous crop's growth stage for daily estimates at field-scale.
- Considering the **method of irrigation application** and **farmer's practice** is highly recommended for estimating CWR & CWU through remote sensing as well as evaluating the performance of an irrigation system.



Reference

- Allen, R.G.; Pereira, L.S.; Raes, D.; Smith, M. (1998) Crop Evapotranspiration—Guidelines for Computing Crop Water Requirements; FAO Irrigation and Drainage Paper 56; FAO: Rome, Italy, 1998, p. 300. 17.
- Cui Y.K., L. Jia, 2014, A Modified Gash Model for Estimating Rainfall Interception Loss of Forest Using Remote Sensing Observations at Regional Scale, *Water*, 2014, 6(4), 993-1012; doi:10.3390/w6040993.
- Cui Y.K., L. Jia, G.C. Hu, and J. Zhou, 2015, Mapping of Interception Loss of Vegetation in the Heihe River Basin of China Using Remote Sensing Observations, *IEEE Geoscience and Remote Sensing Letters (IEEE GRSL)*, 12(1), 23 - 27; doi:10.1109/LGRS.2014.2324635.
- Hu, G., & Jia, L. (2015). Monitoring of evapotranspiration in a semi-arid inland river basin by combining microwave and optical remote sensing observations. *Remote Sensing* (Vol. 7). <https://doi.org/10.3390/rs70303056>
- Martin D. L., Kincaid D. C., & Lyle W. M., (2013). Design and Operation of Sprinkler Systems - Chapter 16. Design and Operation of Farm Irrigation Systems, 2nd Edition, American Society of Agricultural and Biological Engineers, Michigan, 557-631. <https://doi.org/10.13031/2013.23699>
- Menenti, M., Visser, T. N. M., Morabito J. A. and Drovandi, A, 1989. Appraisal of irrigation performance with satellite data and georeferenced information. J.R. Rydzewski and C.F. Ward (eds.). *Irrigation Theory and Practice*. Inst. of Irrigation Studies, Southampton Univ., UK: 785-801
- Yazar, A., 1984. Evaporation and drift losses from sprinkler irrigation systems under various operating conditions. *Agric. Water Manag.* 8, 439-449. [https://doi.org/10.1016/0378-3774\(84\)90070-2](https://doi.org/10.1016/0378-3774(84)90070-2)

Thank you!

**Aerospace Information Research Institute(AIR)
Chinese Academy of Sciences(CAS)**

www.aircas.ac.cn

