

The role of history and strength of the oceanic forcing in sea-level projections from Antarctica with the Parallel Ice Sheet Model

Ronja Reese¹, Anders Levermann^{1,2,3}, Torsten Albrecht¹, H el ene Seroussi⁴, Ricarda Winkelmann^{1,2}

<https://doi.org/10.21941/2019-330>
Preprint. Discussion started: 21 January 2020
  Author(s) 2020. CC BY 4.0 license.



The role of history and strength of the oceanic forcing in sea-level projections from Antarctica with the Parallel Ice Sheet Model

Ronja Reese¹, Anders Levermann^{2,3}, Torsten Albrecht¹, H el ene Seroussi⁴, and Ricarda Winkelmann^{1,2}

¹Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, P.O. Box 60 12 03, 14412 Potsdam, Germany

²University of Potsdam, Institute of Physics and Astronomy, Karl-Liebknecht-Str. 24-25, 14476 Potsdam, Germany

³LDEO, Columbia University, New York, USA

⁴Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

Correspondence: Ronja Reese (ronja.reese@pik-potsdam.de)

Abstract. Mass loss from the Antarctic Ice Sheet constitutes the largest uncertainty in projections of future sea-level rise. Ocean-driven melting underneath the floating ice shelves and subsequent acceleration of the inland ice streams is the major reason for currently observed mass loss from Antarctica and is expected to become more important in the future. Here we show that for projections of future mass loss from the Antarctic Ice Sheet, it is essential (1) to better constrain the sensitivity of sub-shelf melt rates to ocean warming, and (2) to include the historic trajectory of the ice sheet. In particular, we find that while the ice-sheet response in simulations using the Parallel Ice Sheet Model is comparable to the median response of models in three Antarctic ice-sheet intercomparison projects – ISMIP1, LARMIP2 and ISMIP6 – conducted with a range of ice-sheet

What?

We analyse projection of Antarctica's future sea-level rise – ISMIP6 and LARMIP-2 - with all simulations and initial configurations.

Main finding 1

Sea-level projections for the highest emission scenario

Why?



Main difference is how the basal-melt rates change in the ISMIP6 projections is used.

Main finding 2

Starting the experiments after a preceding history of 5 and 50% in comparison to a 'cold-start'.

Implications

To better constrain Antarctic sea-level projection rates and (2) make 'hindcasting' experiments.



Go to TCD manuscript

Very short summary

Read about the role of the history

Read about the oceanic forcing

Go to the outlook



- 1 Potsdam Institute for Climate Impact Research, Potsdam, Germany
- 2 University of Potsdam, Institute of Physics and Astronomy, Potsdam, Germany
- 3 LDEO, Columbia University, New York, USA
- 4 Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA



Tipping Points in Antarctic Climate Components

References & Acknowledgements

1

Very short summary

What?

We analyse projection of Antarctica's future sea-level contribution following two recent Model Intercomparison Projects – ISMIP6 and LARMIP-2 - with all simulations done with the Parallel Ice Sheet Model and started from the same initial configurations (using only ocean forcing; using PICO in ISMIP6, model response comparable to median in both MIPs).

Main finding 1

Sea-level projections for the highest emission scenario RCP8.5 vary by an order of magnitude.

Why?



Main difference is how the basal-melt rates change: in the LARMIP-2 projections a sensitivity three times as large as in our ISMIP6 experiments is used.

Main finding 2

Starting the experiments after a preceding historic simulation from 1850 to 2015, the sea-level contribution increases between 5 and 50% in comparison to starting from equilibrium.


So what?

To better constrain Antarctic sea-level projections, it is important to (1) improve the sensitivity of sub-shelf melt rates and (2) make 'hindcasting' experiments.



2

The role of history



The historic simulation increases mass loss in future projections by up to 50%.



Back to start

➤ All simulations were done with the Parallel Ice Sheet Model



Learn more about PISM (website)

- Sea-level contribution between 2015 and 2100 for different simulations driven by CMIP5 ocean forcing as provided by ISMIP6 (Seroussi et al., under review). The simulations start from two different initial states, one with and one without a preceding historic simulation from 1850 to 2014.

Why could this be the case? Some ideas:


- After the historic simulations, the configuration is losing mass and shows thinning in many regions
- Small changes in geometry over the historic simulation might have made the state more vulnerable to ocean changes
- Non-linearities, e.g., in temperature field or buttressing
- Historic state is closer to an instability

3

The role of the oceanic forcing




Projections for RCP 8.5



Why is the sea-level contribution in our ISMIP6 experiments an order of magnitude smaller?

Central reason: Higher melt rate sensitivity to ocean warming in LARMIP experiments...



Melt rates increase differently for similar ocean warming: changes in LARMIP (grey lines) are based on ocean modelling studies, those in our ISMIP6 experiments calculated with PICO.

LARMIP: red curve shows median, red shaded region likely and very likely ranges.
ISMIP6: three different CMIP5 forcing experiments.



Future sea-level projections: a comparison of two Model Intercomparison Projects (ISMIP6 and LARMIP)

- In line with LARMIP-2, only ocean-forcing from ISMIP6 is applied, basal melting in ISMIP6 experiments is calculated with PICO
- Model response shown here is comparable to the median found in both MIPs (also true for initMIP)
- Sea-level contributions in our ISMIP6 experiments are an order of magnitude smaller than estimates from LARMIP based on the same initial setup, both modelled with the Parallel Ice Sheet Model
- This can be explained by the different sensitivity of melt rates to ocean changes: the changes in LARMIP-2 range from 9 to 16 m/a/K, based on oceanographic estimates (Jenkins 1991, et al. Payne 2007)
- We find that this sensitivity is consistent with a coupled ice-ocean modelling study of Thwaites glacier (Seroussi et al., 2017)
- Changes in ISMIP6 were simulated with PICO which – for the current choice of parameters that were tuned to fit present-day melt rates – yields smaller sensitivities of 2.2 m/a/K over the entire ice shelves and 5.3 m/a/K close to the grounding lines
- Similar findings on role of the basal melt rate parameterization's sensitivity were reported in Jourdain et al. (under review)

Future sea-level projections: role of history

- Mass loss estimates increase by up to 50% when started after a historic simulation in contrast to a cold start simulation based on the same initial setup
- Increases are smaller for the LARMIP-2 estimates (5-7%) where the ocean forcing is stronger and induces larger overall mass loss

Outlook

- Improve sub-shelf melt parameterizations such as PICO by re-tuning with melt sensitivities derived from observations and modelling
- Use hindcasting of observed changes over the past decades to better constrain ice sheet model projections



References & further reading

- Reese et al.: The role of history and strength of the oceanic forcing in sea-level projections from Antarctica with the Parallel Ice Sheet Model, *The Cryosphere Discuss.*, <https://doi.org/10.5194/tc-2019-330>, in review, 2020.
- Seroussi et al. : ISMIP6 Antarctica: a multi-model ensemble of the Antarctic ice sheet evolution over the 21st century, *The Cryosphere Discuss.*, <https://doi.org/10.5194/tc-2019-324>, in review, 2020.
- Levermann et al. : Projecting Antarctica's contribution to future sea level rise from basal ice shelf melt using linear response functions of 16 ice sheet models (LARMIP-2), *Earth Syst. Dynam.*, 11, 35–76 <https://doi.org/10.5194/esd-11-35-2020>, 2020.
- Jourdain et al. : A protocol for calculating basal melt rates in the ISMIP6 Antarctic ice sheet projections, *The Cryosphere Discuss.*, <https://doi.org/10.5194/tc-2019-277>, in review, 2019.
- Seroussi, et al. (2017), Continued retreat of Thwaites Glacier, West Antarctica, controlled by bed topography and ocean circulation, *Geophys. Res. Lett.*, 44, 6191– 6199, doi:[10.1002/2017GL072910](https://doi.org/10.1002/2017GL072910).
- Jenkins: Ice shelf basal melting: Implications of a simple mathematical model, *Filchner-Ronne Ice Shelf Programme Report*, 5, 32–36, 1991
- Payne et al.: Numerical modeling of ocean-ice interactions under Pine Island Bay's ice shelf, *Journal of Geophysical Research: Oceans*, 112, 2007, doi: <https://doi.org/10.1029/2006JC003733>

Acknowledgements

We thank the Climate and Cryosphere (CliC) effort, which provided support for ISMIP6 through sponsoring of workshops, hosting the ISMIP6 website and wiki, and promoted ISMIP6. We acknowledge the World Climate Research Programme, which, through its Working Group on Coupled Modelling, coordinated and promoted CMIP5. We thank the climate modeling groups for producing and making available their model output, the Earth System Grid Federation (ESGF) for archiving the CMIP data and providing access, the University at Buffalo for ISMIP6 data distribution and upload, and the multiple funding agencies who support CMIP5 and ESGF. We thank the ISMIP6 steering committee, the ISMIP6 model selection group and ISMIP6 dataset preparation group for their continuous engagement in defining ISMIP6. Development of PISM is supported by NASA grant NNX17AG65G and NSF grants PLR-1603799 and PLR-1644277. The authors gratefully acknowledge the European Regional Development Fund (ERDF), the German Federal Ministry of Education and Research and the Land Brandenburg for supporting this project by providing resources on the high performance computer system at the Potsdam Institute for Climate Impact Research. Computer resources for this project have been also provided by the Gauss Centre for Supercomputing/Leibniz Supercomputing Centre (www.lrz.de) under Project-ID pr94ga and pn69ru. R.R. is supported by the Deutsche Forschungsgemeinschaft (DFG) by grant WI4556/3-1 and through the TiPACCs project that receives funding from the European Union's Horizon 2020 research and innovation programme under grant agreement no. 820575. T.A. is supported by the Deutsche Forschungsgemeinschaft (DFG) in the framework of the priority program 'Antarctic Research with comparative investigations in Arctic ice areas' by grant WI4556/2-1 and WI4556/4-1 and in the framework of the PalMod project (FKZ: 01LP1925D), supported by the German Federal Ministry of Education and Research (BMBF) as a Research for Sustainability initiative (FONA). H.S. was supported by grants from NASA Cryospheric Science and Modeling, Analysis, Predictions Programs.



Tipping Points in Antarctic
Climate Components



CR 1.1
D2556