Probabilistic flood loss estimation for residential buildings in Europe

Max Steinhausen^{1,2}, Kai Schröter¹, Stefan Lüdtke¹ and Heidi Kreibich¹

¹ Helmholtz Centre Potsdam, GFZ German Research Centre for Geosciences, Section Hydrology ² Humboldt University Berlin, Geography Department



EGU 2020 | max.steinhausen@gfz-potsdam.de



Introduction and Research Questions

European wide flood loss assessments are essential for governments to:

- support climate change adaptation policies (Van Renssen, 2013)
- manage the European Union solidarity fund (Hochrainer et al., 2010)
- comply with the European Flood Directive (EU, 2007)

Furthermore the (re-)insurance sector relies on flood loss estimations to calculate premiums.

Flood loss is often estimated by simple stagedamage-curves and lacks adequate uncertainty information.



With our research we aim to:

- provide Europe-wide probabilistic flood loss estimations with inherent uncertainty quantification
 - identify the drivers that contribute to flood loss changes in the present and under future climate conditions
- make innovative modelling approaches more accessible



EGU 2020 | max.steinhausen@gfz-potsdam.de

Features of BN-FLEMOps

- Bayesian Network Flood Loss
- Estimation Model for the private sector
- Direct loss to private buildings
- Multi-variable model for better damage process representation
- Probabilistic results provide uncertainty information for e.g. decision-making
- Transferable in location, scale and time
- Predictions are possible with incomplete input data





EGU 2020 | max.steinhausen@gfz-potsdam.de

Probabilistic flood loss estimation



$$P(X_i, \dots, X_n) = \prod_{i=1}^n P(X_i | parents(X_i))$$

Equation for the joint probability distribution in BNs

- BN-FLEMOps uses discretized variables
- Joint probability distributions are represented as node probability tables (example on the left)
- These tables contain the conditional probabilities for each node and their associated child node(s)
- ←The example on the left shows a simplified node probability table for an example of the BN where only water depth information is available to estimate relative loss



EGU 2020 | max.steinhausen@gfz-potsdam.de

Consistent European input data



GFZ Helmholtz Centre Potsdam

EGU 2020 | max.steinhausen@gfz-potsdam.de

Validation in case studies

Validation of the BN-FLEMOps was performed by estimating flood loss for historic events in 3 case studies in Italy, Austria and Germany (Lüdtke et al. 2019).

In a comparison with 20 flood loss models we were able to show that BN-FLEMOps outperforms a model ensemble. It requires less data and setup work than an ensemble and provides probabilistic results with inherent uncertainty information (Steinhausen et al. *in review*).

	Caldogno	Lech	Mulde
Reported loss [mil.€]	7.5	1.9	240
Estimated median loss [mil.€]	9.5	7.8	136
Interquartile range [mil.€]	4.0 – 18.5	3.6 - 3.4	87.1- 246.0
No. of data points	295	22	74





EGU 2020 | max.steinhausen@gfz-potsdam.de

European application

100-year European flood loss map (undefended scenario, reference climate)



The map on the left shows the 20% quantile (Q20), the median (Q50) and the 80% quantile (Q80) of a flood loss estimation performed with BN-FLEMOps for a continent-wide 100-year flood scenario. Results are aggregated on the NUTS-3 level.

For this flood scenario, the highest flood losses are estimated in the flood plains of major European rivers, such as Rhine and Meuse, Danube, Seine, Loire, and Po. The flood scenario does not account for any flood protection infrastructure in place.

The total accumulated loss for residential buildings in Europe is estimated to 79.0 billion \in (Q20 = 32.3; Q80 = 213.8) (Lüdtke et al. 2019).



EGU 2020 | max.steinhausen@gfz-potsdam.de

OASIS-LMF implementation



The OASIS Loss Modelling Framework (LMF) is an open source platform for developing, deploying and executing catastrophe models to enable the "plug and play" of hazard and vulnerability modules (<u>https://oasislmf.org/</u>).

To make BN-FLEMOps easier to use and transfer new approaches from research into practice the model was implemented in the OASIS LMF.

Data for the use of BN-FLEMOps are available on the OASIS HUB (<u>https://oasishub.co/dataset/european-exposure-data-for-bn-flemo-models-gfz</u>).

← The figure on the left shows the structure of compliant tables in the OASIS LMF

GFZ Helmholtz Centre POTSDAM

EGU 2020 | max.steinhausen@gfz-potsdam.de

Summary and Outlook

- The Bayesian Network methodologies of BN-FLEMOps produces probabilistic loss estimates
- BN-FLEMOps is validated in case studies and comparisons show good performance
- Consistent European input data enables continent-wide loss estimation
- First results show that urban areas along major rivers would suffer most of the 79.0 billion € estimated loss for residential buildings in Europe
- The OASIS LMF implementation makes BN-FLEMOps more accessible



This project has received funding from the European Commission's, Horizon 2020 research and innovation programme under Grant Agreement number: 730381 — H2020_Insurance

- An improved asset methodology will be used for future loss estimations (figure below)
- Exposure scenarios for future climate conditions will be incorporated
- · Adaptation by private precaution will be studied



9 HELMHOLTZ

EGU 2020 | max.steinhausen@gfz-potsdam.de