


# Identifying sources of changed precipitation in paleoclimate studies through moisture tracking



## A case study for orbital extremes over the Mediterranean Sea

Ruud van der Ent<sup>1,2</sup>, Joyce Bosmans<sup>2,3</sup>, Rein Haarsma<sup>4</sup>, Sybren Drijfhout<sup>2,4,5</sup>, and Frits Hilgen<sup>2</sup>

<sup>1</sup>Delft University of Technology, Netherlands (r.j.vanderent@tudelft.nl)

<sup>2</sup>Utrecht University, Utrecht, Netherlands

<sup>3</sup>Radboud University, Nijmegen, Netherlands (joyce.bosmans@ru.nl)

<sup>4</sup>Royal Netherlands Meteorological Institute (KNMI), Netherlands

<sup>5</sup>University of Southampton, United Kingdom

# Moisture tracking

What are the sources of precipitation?

What is the fate of evaporation?


Questions can be answered with ‘offline’ moisture tracking models, e.g. WAM-2layers ([Van der Ent et al., 2014](#))

Required input is daily or sub-daily gridded fields of:


- Precipitation
- Evaporation
- Wind speed (at several pressure or model levels)
- Humidity (at several pressure or model levels)

# Applications of moisture tracking


Sources of precipitation ([Guo et al., 2019](#))




Land-use change impacts ([Wang-Erlandsson et al., 2018](#))



Fate of land evaporation from specific river basins ([Link et al., 2020](#))



Future water cycle changes ([Findell et al., 2019](#))



# First application in a paleoclimate study: Orbital extremes over the Mediterranean Sea

## Paleoceanography and Paleoclimatology



**AGU100** ADVANCING  
EARTH AND  
SPACE SCIENCE

### RESEARCH ARTICLE

10.1029/2019PA003655

#### Key Points:

- Moisture sources for precession-induced enhanced winter precipitation are local in fall and from the Atlantic in late winter
- For obliquity, precipitation changes are smaller; local and Atlantic sources play an equal role
- The Atlantic sources are not related to storm tracks but to low-latitude surface pressure changes

## Precession- and Obliquity-Induced Changes in Moisture Sources for Enhanced Precipitation Over the Mediterranean Sea

**J.H.C. Bosmans<sup>1,2</sup>, R.J. van der Ent<sup>1,3</sup>, R.J. Haarsma<sup>4</sup>, S.S. Drijfhout<sup>4,5,6</sup>, and F.J. Hilgen<sup>1</sup>**

<sup>1</sup>Faculty of Geosciences, Utrecht University, Utrecht, Netherlands, <sup>2</sup>Faculty of Science, Radboud University, Nijmegen, Netherlands, <sup>3</sup>Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, Netherlands, <sup>4</sup>Royal Netherlands Meteorological Institute (KNMI), De Bilt, Netherlands, <sup>5</sup>Ocean and Earth Science, National Oceanography Centre, University of Southampton, Southampton, UK, <sup>6</sup>Faculty of Science, Utrecht University, Utrecht, Netherlands

# Precession-induced enhanced precipitation over the Mediterranean

What is the source of freshwater input, thought to cause sapropels?

Winter precipitation could play an important role

([Bosmans et al., 2015](#)).

Is **enhanced winter precipitation** related to **local processes** or **Atlantic storm tracks**?



Sapropels in Sicily from the Miocene [Tuenter \(2004\)](#)

# Methods


Orbitally extreme experiments performed with the state-of-the-art climate model EC-Earth.

Moisture tracking with WAM-2layers to compute:

- $\rho$  = precipitation recycling ratio (fraction of precipitation from Mediterranean evaporation)
- $\varepsilon$  = evaporation recycling ratio (fraction of evaporation from ending up as Mediterranean precipitation)

# Results for precession

Precipitation differences between precession minimum (Pmin) and precession maximum (Pmax) related to changes in the importance of local vs. remote moisture sources





# Precession September+October

20% more precipitation during precession minimum.




Moisture sources during Pmin:



Differences between Pmin and Pmax

Blue = stronger sources during Pmin

Red = weaker sources during Pmin



>2.5 mm/2months = 79% of  $P^M$







# Precession January + February

50% more precipitation during Pmin.




Stronger sources locally and from the Atlantic:

Pmin - Pmax January + February



Blue = more storm track activity during Pmin



Standard deviation of 500 hPa geopotential height (m)




Enhanced precipitation driven by storm track activity over Atlantic

# Conclusions

Enhanced precipitation during precession minimum due to different mechanisms.

Stronger **local moisture recycling** in September and October.

Increased **Atlantic storm track activity** during January and February.



# More information and contact

Bosmans, J. H. C., van der Ent, R. J., Haarsma, R. J., Drijfhout, S. S. and Hilgen, F. J.: Precession- and Obliquity-Induced Changes in Moisture Sources for Enhanced Precipitation Over the Mediterranean Sea, *Paleoceanogr. Paleoclimatology*, 35(1), 1–14, [doi:10.1029/2019PA003655](https://doi.org/10.1029/2019PA003655), 2020.

Information on orbital extremes in the Mediterranean:

Joyce Bosmans ([joyce.bosmans@ru.nl](mailto:joyce.bosmans@ru.nl))

Collaborations on moisture tracking for other paleoclimate studies:

Ruud van der Ent ([r.j.vanderent@tudelft.nl](mailto:r.j.vanderent@tudelft.nl))

