

# Coastal geomorphic response to volcano-tectonic activity in the Campi Flegrei Caldera: new insight from the geoarchaeological study of Portus Julius (Pozzuoli Gulf, Italy).

Claudia Caporizzo <sup>1</sup>, Pietro P.C. Aucelli <sup>1</sup>, Gaia Mattei <sup>1</sup>, Aldo Cinque <sup>2</sup>, Salvatore Troisi <sup>1</sup>, Francesco Peluso <sup>1</sup>, Michele Stefanile <sup>3</sup>, Gerardo Pappone <sup>1</sup>.

<sup>1</sup>Dipartimento di Scienze e Tecnologie, Università degli Studi di Napoli Parthenope, Centro Direzionale Is. C4, 80121 Napoli, Italy.

<sup>2</sup>Dipartimento di Scienze della Terra, dell'Ambiente e delle Risorse, Università degli Studi di Napoli Federico II, Largo San Marcellino, 10, 80138 Napoli, Italy.

<sup>3</sup>Dipartimento Asia Africa e Mediterraneo, Università degli Studi di Napoli L'Orientale, Piazza S. Domenico Maggiore, 12, 80134 Napoli, Italy.




Figure 1. Geological map of the Campi Flegrei with the main volcanic events and the location of Portus Julius.




Figure 2. Archaeological map of Portus Julius with the archaeological sea level markers detected.

## The Study Area

The ancient Portus Julius is one of the largest underwater archaeological sites in Italy and it represents a great point of interest for marine archaeology studies [1]. This site, located within the Campi Flegrei Caldera (Figure 1) and precisely inside the Underwater Archeological Park of Baia, it's a perfect example of submerged archaeological heritage that records fundamental traces of the ancient coastal morphologies, highly vulnerable to present coastal processes because of recent submersion.

During the Roman period, Portus Julius (Figure 2) experienced at least two different construction phases: a brief military phase (37 BC – 12 BC) during which the entry channel was built, with a length of 300 m and a protection of pier structures with huge *pilae*, and a long commercial phase (after 12 BC) during which the area was restored with the construction of numerous warehouses (*horrea*) and fish tanks. The fish tank here studied represents one of the few examples of *piscinae* related to commercial activities and not to private maritime villas and its recovery within the entry channel can be related to the presence of a fish market or areas intended for the production of the *garum*. According to the historical sources the harbor was abandoned during the IV century AD due to a bradyseismic crisis.

This study presents the results of high-precision surveys on the most reliable archaeological markers of the ancient sea level, aimed at the evaluation of the ground movements affecting the area of Portus Julius during the Roman period.




Figure 3. a: measuring point in the fish tank (MHW: Mean High Water; MLW: Mean Low Water); b: intact sluice gate of the fish tank of Portus Julius; c: measuring point in the pilae; d: concrete change in one of the pilae of Portus Julius; e: TLP in the villa close to Portus Julius; f: one of the floors of the maritime villa of Republican age.

## Methods

Direct surveys were carried out on three different types of archaeological sea level markers, by using a metric roll and a depth gauge, to measure the size and submersion of the archaeological investigated structures related to different constructive phases of the port. The submersion measurements were taken from the tide gauge in the Port of Naples

Among those sea-level markers, the fish tank's sluice gate (*cataracta*) (Figure 3a, 3b) and the concrete change of the *pilae* (sensu Mattei et al., [2]) (Figure 3c, 3d) are considered high-precision sea level index points (SLIPs) [3, 4].

The former RSLs were calculated considering the Indicative Range (IR, sensu Shennan et al. [3]) and the functional clearance of each marker using the following equation:

$$SLIP_{\text{marker}} \text{ (m)} = E - fc_{\text{marker}} - IR/2$$

The third marker (Figure 3e, 3f) is a floor belonged to a maritime villa built on the ancient sandy coast, therefore considered a terrestrial limiting point (TLP, [3, 4]).

Moreover, a morphometric reconstruction of the fish tank - key element of our geoarchaeological interpretations - has been carried out to obtain a detailed morphometric analysis of the marker and to preserve a 3D documentation of this important and unique historical testimony (Figure 4).




Figure 4. a: reconstruction of the photogrammetric system used during the survey; b: zoom of the 3D reconstruction of the well preserved gate; c: position of the photogrammetric cameras during the survey; d: 3D sketch of the surveyed fish tank.

## Results and Conclusions

From the submersion values, a RSL of -4.7/-5.0 m MSL related to the period before the 37 BC from the living floors, a RSL of -3.1 m MSL related to the 37 BC from the *pilae* and a RSL of 3.1 m MSL related to the 12 BC from the fish tank were determined.

The results achieved in terms of the RSL variations allowed us to reconstruct the evolution of the coastal sector over the last 2100 years (Figure 5). It is possible to affirm that, at the end of the Republican Age, the study area was shaped as a low coastal sector located next to a sandy spit, above which the former *Via Herculanea* was constructed (Figure 5).

The maritime villa, analyzed in this paper, was located near to the above-mentioned spit in a flat area at a depth of about -3.5 m MSL. Therefore, considering its functional clearance, an RSL at -4.70/-5.2

m MSL can be presumed for that period. The RSL at -3.2 m MSL measured from the *pilae* demonstrates that some subsidence had already occurred when the military port was built in 37 BC. As well, we can suppose that the subsidence, which produced an increase in the water depth, encouraged the construction of the harbor facility.

Finally, the standing of the RSL from the fish tank at about -3.2 m MSL, even after 12 BC, demonstrates that the area experienced a period of volcano-tectonic stability.

Moreover, by comparing our measurements with the Glacio-Hydro-Isostatic Adjustment (GIA) models proposed by several authors [5, 6] (Figure 6), it is possible to determine that the only contribution of the vertical ground movements (VGMs) occurred between the early first century BC and the early first century AD. In this period, the GIA produced an SL variation in our study area of a centimetric amount (from -1.05 to -1.09 m MSL). Consequently, at the same time and in terms of VGMs, the study area suffered a subsidence of about 4.00 meters, and experienced a rapid increase (bradyseismic crisis) of another 2.00 m before 37 BC.

## References:

- Gianfrotta, P.A. Ricerche nell'area sommersa del "Portus Iulius" (1988-'90 e successive): Un riepilogo. Atlante Temat. Topogr. Antica 2012, 22, 1-20.
- Mattei, G.; Troisi, S.; Aucelli, P.P.C.; Pappone, G.; Peluso, F.; Stefanile, M. Sensing the Submerged Landscape of Nisida Roman Harbour in the Gulf of Naples from Integrated Measurements on a USV. *Water* 2018, 10, 1-31.
- Shennan, I.; Long, A.J.; Horton, B.P. Handbook of Sea-Level Research; John Wiley & Sons: Oxford, UK; New York, USA, 2015.
- Rovere, A.; Raymo, M.E.; Vacchi, M.; Lorscheid, T.; Stocchi, P.; Gómez-Pujol, L.; Harris, D.L.; Casella, E.; O'Leary, M.J.; Hearty, P.J. The analysis of the Last Interglacial (MIS 5e) relative sea-level indicators: Reconstructing sea-level in a warmer world. *Earth Sci. Rev.* 2016, 159, 404-427.
- Lambeck, K.; Antonioli, F.; Anzidei, M.; Ferranti, L.; Leoni, G.; Scicchitano, G.; Silenzi, S. Sea level change along the Italian coasts during Holocene and prediction for the future. *Quat. Int.* 2011, 232, 250-257.
- Vacchi, M.; Ghilardi, M.; Melis, R.T.; Spada, G.; Giaime, M.; Marriner, N.; Lorscheid, T.; Morhange, C.; Burjachs, F.; Rovere, A. New relative sea-level insights into the isostatic history of the Western Mediterranean. *Quat. Sci. Rev.* 2018, 201, 396-408.