

Climate warming does not adequately translate to increased radial stem growth of coniferous species along the Alpine treeline ecotone

W Oberhuber¹, U Bendler¹, V Gamper¹, J Geier¹, A Hölzl¹, H Krismer¹, W Kofler¹, B Waldböth¹, G Wieser²

¹Department of Botany, University of Innsbruck, Austria; ²Department of Alpine Timberline Ecophysiology, BFW, Innsbruck, Austria



Background

Climate observations reveal a pronounced rise of air temperature in the Central European Alps (CEA) in recent decades (Gobiet et al. 2014) and it is also well established that tree growth at high altitude is mainly limited by low temperature during the growing season (Körner & Paulsen 2004; Wieser et al. 2019).

Aim and Method

We developed tree ring series of coniferous species at subalpine sites (n=194 trees) and evaluated whether trends in basal area increment (BAI) are in line with two phases of climate warming. Differences in temperature and BAI during 5-yr long periods at the start and end of these warming phases were compared.




Fig. 1 Two phases of climate warming (W_{ph} I and II) in the Alpine region in the period 1915–2015 (data from Auer et al. 2007).




Fig. 2 Trend in basal area increment (BAI) from 1915–2015 in dominant conifers in CEA (n>50 trees per species).

Species	W_{ph} I	W_{ph} II	W_{ph} III
<i>Pinus cembra</i>	5.35±0.84 ^a	1.89±1.78 ^b	4.84±1.33 ^a
<i>Larix decidua</i>	4.49±1.91 ^a	4.65±0.54 ^a	6.66±1.05 ^b
<i>Picea abies</i>	4.24±0.82 ^a	4.28±1.23 ^a	10.57±1.22 ^c

Table 1 Increase in BAI (cm²) during phases of climate warming (W_{ph} I and II) and the whole study period (W_{ph} III). Different letters indicate significant differences at $P < 0.05$.

Results

Although enhanced tree growth occurred at high altitude in response to climate warming, results revealed that growth increase has declined in *Pinus cembra* and remained constant in *Picea abies* and *Larix decidua* despite the more intensified warming taking place since late 1970s (**Fig. 1, Table 1**).

Pinus cembra dominates at high altitude in the CEA and therefore was expected to mainly benefit from climate warming. However, increase in air temperature during the growing season primarily favored growth of *Larix decidua* and *Picea abies* (**Fig. 2, Table 1**).

Conclusions

We explain missing adequate growth response to recent climate warming by

- (i) intensified competition for resources (nutrients, light, water) in increasingly denser stands leading to changes in carbon allocation among tree organs, and
- (ii) species-specific sensitivity to climate variables beyond the growing season.

References

Auer I et al. (2007) Int J Climatol; Gobiet et al. (2014) Sci Total Environ; Körner & Paulsen (2004) J Biogeogr; Wieser et al. (2019) Forests.



For more details please download paper (open access):

<https://www.mdpi.com/1999-4907/11/2/132>



forests


(2020) 11,132

doi:10.3390/f11020132



Article

Growth Trends of Coniferous Species along Elevational Transects in the Central European Alps Indicate Decreasing Sensitivity to Climate Warming

Walter Oberhuber ¹ , Ursula Bendler ¹, Vanessa Gamper ¹, Jacob Geier ¹, Anna Hölzl ¹,
Werner Kofler ¹, Hanna Krismer ², Barbara Waldböth ¹ and Gerhard Wieser ^{3,*} 