

Radiative Properties of Aerosols and Clouds from Observations and Models over the Southeast Atlantic

Ian Chang and the

NASA ORACLES Science Team

School of Meteorology
University of Oklahoma, Norman, OK, USA

EGU 2020 Virtual Presentation

ian.chang@ou.edu



Science Contributors

U of Oklahoma

I. Chang

C. Flynn

L. Gao

J. Redemann

Aircraft teams

S. Burton

H. Chen

R. Ferrare

S. Leblanc

M. Kacenenbogen

K. Pistone

M. Segal-Rozenhaimer

S. Schmidt

Y. Shinozuka

Modeling groups

P. Castellanos

Y. Feng

G. Ferrada

K. Longo

M. Mallet

P. Saide

Leadership contributors

S. Christopher

S. Doherty

M. Diamond

K. Meyer

R. Wood

P. Zuidema

lan.chang@ou.edu

Introduction


Quantifying above-cloud aerosol optical depth (ACAOD) and underlying cloud optical depth (COD) from observations and chemical transport models remains challenging, complicating the estimate of the aerosol direct radiative effects.

Objectives



- Evaluate the performance of above-cloud aerosol optical depth (ACAOD) retrievals from satellite and the impact of temporal collocation on the retrieval results.
- Compare cloud optical depth (COD) retrievals between satellite and NASA ORACLES aircraft.
- Evaluate chemical transport model diagnostics of AOD and COD over the southeast Atlantic.

Study domain: September 2016 in South Atlantic

Aqua true color on 19 September 2016



**Aerosol
above
cloud**



Above-cloud aerosol optical thickness

Yellow to brown shading: above-cloud AOD

Black contours: Cloud fraction

Land shading: Fire frequency

Arrows: wind vectors

Part 1 results: Satellite and aircraft intercomparisons

Data Sets

Aircraft

1) 4STAR (Sun-Tracking Atmospheric Research)

- Hyperspectral AOD (355-1650nm) from sun photometer

2) HSRL-2 (High Spectral Resolution Lidar)

- Aerosol extinction at 532 nm and aerosol backscatter and depolarization at 532 and 1064 nm


3) SSFR (Solar Spectral Flux Radiometer)

- Upwelling / downwelling solar spectral irradiance (350 nm - 2150 nm) to retrieve COD

Satellite retrievals







- **Type 1:** Optimal estimation method from six MODIS solar channels (Meyer et al., 2015)
- **Type 2:** MODIS 0.47/0.87 μ m reflectance ratio (Jethva et al., 2013)
- **Type 3:** SEVIRI 0.64/0.81 μ m reflectance ratio (Chang and Christopher, 2016)

Results: 4STAR-HSRL ACAOD comparisons








- RMSE increases with coarser temporal collocation due to increased differences in the ACAOD measurements.
- The increased ACAOD differences suggests the presence of aerosol heterogeneity in this region.

Results: 4STAR-satellite ACAOD comparisons



Finer temporal collocation improves the retrieval performance both for R and $RMSE$

Results: Cloud optical depth comparisons








- Finer temporal collocation reduces the COD differences between satellite and aircraft retrievals
- SEVIRI still underestimates COD due to its coarser spatial resolution

Part 2 results: Model and aircraft intercomparisons


Model Descriptions

- 1) **WRF-Chem** model that couples with the Community Atmosphere Model version 5 (CAM5) physics package (36-km horizontal grid resolution)
- 2) **ALADIN**: Regional climate model based on bi-spectral semi-implicit semi-lagrangian model (12 km horizontal resolution)
- 3) **GEOS-5**: The Goddard Earth Observing System Version 5 (GEOS-5) is a global modeling system developed at NASA Global Modeling and Assimilation Office (GMAO) (25 × 31.25 km horizontal resolution)
- 4) **GEOS-CHEM**: A global 3-D model of atmospheric composition driven by assimilated meteorological data GEOS-FP data from the GMAO (25 × 31.25 km horizontal resolution)
- 5) **MERRA-2**: Modern-Era Retrospective-analysis for Research and Applications- Version 2 uses GEOS-5 that is radiatively coupled to GOCART (50 × 61.25 km horizontal resolution)

Results: ORACLES 2016 aircraft-model AOD








±30-minute collocation




Example of one model grid

Results: ORACLES 2017 aircraft-model AOD



Results: $10^{\circ} \times 10^{\circ}$ COD Frequency Distributions (Sep 2016)

Domain: 5°E to 15°E , 20°S to 10°S



Summary and future work

- ACAODs over the southeast Atlantic shows heterogeneity beyond 15 minutes, implying that polar-orbiting satellite retrievals need to impose stringent temporal collocation.
- Correction to CODs due to overlying aerosol attenuation has shown to be comparable to aircraft COD retrievals.
- Models and 4STAR AODs have a higher agreement during 2016 than during 2017. Major sources of model uncertainties need to be diagnosed.
- Models tend to produce optically thicker low-level liquid clouds than satellite-retrieved CODs over the southeast Atlantic.
- Future work needs to revisit the direct aerosol radiative effects in southeast Atlantic. The impact of mid- and high-level clouds on satellite retrievals of underlying aerosols is under investigation.