

NaFoLiCA: Synoptic-scale controls of fog and low cloud variability in the Namib Desert

Hendrik Andersen, Jan Cermak, Julia Fuchs, Peter Knippertz, Marco Gaetani, Julian Quinting, Sebastian Sippel, and Roland Vogt

Karlsruhe Institute of Technology (KIT)




Motivation: Fog relevant to local ecosystems but poorly understood



L: *Stipagrostis sabulicola*: Roth-Nebelsick et al. (2013),
R: *Onymacris unguicularis*: Anderson (2008) [modified]

Previous work I: Satellite detection of Fog and Low Clouds (FLC)


- SEVIRI-based FLC detection at native resolution (15 min, 3x3 km)
- Validation with station measurements shows good skill



Andersen and Cermak 2018, AMT

→ Continuous observations since 2004, collaborations welcome!

Previous work II: Mapping of spatiotemporal FLC patterns



Andersen et al. 2019, ACP

- Vertical position of low-cloud layer varies by season.
- Timing of FLC occurrence tightly related to the distance to the coast.
- But: Knowledge gaps concerning drivers of FLC variability


Scientific goal and approach

- Improve scientific understanding of the drivers of day-to-day variability of fog and low clouds in the Namib Desert.
- Analysis is done for two distinct FLC seasons: April, May, June (low-lying FLCs), and September, October, November (higher-level FLCs)
- Data basis: Satellite observations of FLCs from Andersen and Cermak (2018) and ERA5 reanalysis.





Image credits: Julia Fuchs

The seasonal large-scale setting



Top: 14-year average mean sea level surface pressure and 10m winds,
bottom: SST and LTS. Andersen et al. 2020, ACP


AMJ: comparison of meteo. FLC vs. clear days



Left: MSLP and 10m wind differences, right: Z500 and 500hPa wind differences;
Andersen et al. 2020, ACP)

- Significant differences of atmospheric dynamics on FLC vs. clear days at synoptic scales
- Combination of synoptic-scale forcing (Z500) and heat low (MSLP) on FLC days


AMJ: comparison of meteo. FLC vs. clear days



Left: T2m differences, right: TCWV differences; Andersen et al. 2020, ACP

- Higher continental temperatures on FLC days
- T2m anomalies related to increased TCWV from large-scale moisture transport on FLC days (correlation of anomalies over land: 0.75)


Comparison: meteo. FLC vs. clear days



Winds, Q at 700hPa during AMJ (top) and SON (bottom) during FLC days (left), clear days (center) and differences (right); Andersen et al. 2020, ACP)

- Synoptic-scale disturbance → moisture transport during AMJ
- Dry anomaly in free-troposphere where FLCs form in both seasons


Comparison: meteo. on FLC and clear days



Differences on fog vs. clear days: winds, specific humidity (Q) at different levels;
Andersen et al. 2020, ACP


- Dry anomaly in free-troposphere overlies moist anomaly in marine boundary layer on FLC days
- Enhanced radiative cooling at cloud-top helps maintain FLCs
- Higher SSTs upwind suggest increased MBL moisture fluxes

Comparison of backtrajectories




Top: FLC days, bottom: clear days; Andersen et al. 2020, ACP

Concept: Seasonal synoptic-scale controls



- Free-tropospheric advection of dry air
- Longwave cooling
- Dry slot or dry air intrusion
- Heat low



- Free-tropospheric advection of moist air
- Longwave heating
- Onshore advection
- Boundary-layer warm air advection

Andersen et al. 2020, ACP

What have we learned... and what is next?

Conclusions

- FLCs are associated with advection of marine boundary-layer air.
- Synoptic-scale disturbances drive moisture transport, long wave radiative cooling(ocean)/warming(land) and control FLC variability.

Outlook

- Predict FLC occurrence and future changes with statistical models.
- Detailed-case studies based on field campaign.
- Link synoptic to local scales and models.

NaFoLiCA resources

- Satellite FLC-detection paper: 10.5194/amt-11-5461-2018
- Spatiotemporal patterns paper: 10.5194/acp-19-4383-2019
- Synoptic-scale controls paper: 10.5194/acp-20-3415-2020
- IOP paper: 10.1175/bams-d-18-0142.1