



ARISTOTLE  
UNIVERSITY  
OF THESSALONIKI



*Comparison of one year of  $XCH_4$  and  $XCO$  measurements  
using a EM27/SUN low resolution FTIR spectrometer  
to S5P/TROPOMI methane and carbon monoxide columns  
at Thessaloniki, Greece (**revised results**)*

Topaloglou Chrysanthi<sup>1</sup>, Mermigkas Marios<sup>1</sup>, Koukouli Maria-Elissavet<sup>1</sup>, Balis Dimitrios<sup>1</sup>, Hase Frank<sup>2</sup>, Jochen Landgraf,<sup>3</sup> Ilse Aben<sup>3</sup>

contact e-mail address : chtopal@auth.gr

<sup>1</sup>Laboratory of Atmospheric Physics, Department of Physics, Aristotle University of Thessaloniki, Greece

<sup>2</sup>Institute of Meteorology and Climate Research (KIT-IMK), Karlsruhe Institute of Technology, Karlsruhe, Germany

<sup>3</sup>SRON Netherlands Institute for Space Research, Utrecht, The Netherlands







The Total Carbon Column Observing Network (TCCON) (Wunch et al., 2011) measures total columns of CO<sub>2</sub> and CH<sub>4</sub>, using Bruker IFS 125HR spectrometers. These high resolution instruments are rather expensive and need large infrastructure to be set up and expert maintenance, which has to be performed on site. (Frey et al, 2019)

The **Bruker EM27/SUN** portable FTIR spectrometer (Gisi et al., 2011; Frey et al., 2015; Hedelius et al., 2016) developed by KIT in collaboration with Bruker Optics™, is a promising instrument to overcome the above-mentioned shortcomings as it is a mobile, reliable, easy-to-deploy and low-cost supplement to the Bruker IFS 125HR spectrometer used in the TCCON network (Frey et al, 2019)


The **COllaborative Carbon Column Observing Network (COCCON)**, constituted by 60 EM27/SUN spectrometers, with stations around the globe, produces greenhouse gas observations based on common instrumental standards and data analysis procedures, working as an important supplement of TCCON for the quantification of local sinks and sources, to increase the global density of column-averaged greenhouse gas observations



# TCCON- COCCON NETWORK



## COCCON – Next Step





## The Bruker EM27/SUN FTIR spectrometer



It consists of a spectrometer body with dimensions of 35x40x27cm and a solar tracker directly mounted on the spectrometer. It weighs approximately 25 kg and its design achieves high stability and is resistant to thermal influences and mechanical disturbances.

# The Bruker EM27/SUN FTIR spectrometer

	Spectral window (cm-1)	Spectral resolution (cm-1)	Semi-FOV (mrad)	Detector	Retrieval software	Spectroscopy	A priori profiles & averaging kernels
O <sub>2</sub>	7765-8005	0.5	2.36	InGaAs Spectral range:	PROFFAST (Sha et al. 2019)	HITRAN O <sub>2</sub> : 2007 (Geoff Toon's ATM line list)	TCCON map files Column sensitivities from retrieval software
CH <sub>4</sub>	5897-6145			main channel standard (5000 cm-1 low wavenumber cut off)		H <sub>2</sub> O : 2009 +empirical corrections	
CO	4208-4318			CO channel extended (4000 cm-1 low wavenumber cut off).		CH <sub>4</sub> : 2008	
				CO: 2012			



## EM27/SUN PRODUCTS

In this study, column-averaged dry-air mole fractions  $X_{CH_4}$  and  $X_{CO}$  are presented for one year of measurements in Thessaloniki, Greece, an urban site (40.5N°, 22.9E, 60 m a.s.l). For a gas G:

$$X_{Gas} = \frac{Column_{Gas}}{Column_{dryair}}$$

Since  $O_2$  's mixing ratio in the atmosphere can be considered stable

$$Column_{dryair} = \frac{Column_{O_2}}{0.2095}$$

the column-averaged dry-air mole fractions (DMF), denoted  $X_G$  for gas G, are computed using the retrieved  $O_2$  columns as a measure of the dry air column

$$X_{Gas} = \frac{Column_{Gas}}{Column_{O_2}} \cdot 0.2095$$

Dividing by  $O_2$  improves the precision of the measurement by significantly reducing the effects of instrumental or measurement errors that are common to both the gases (tracker problems leading to mis-pointing and zero-level offsets) or by errors the use of in the surface pressure



## EM27/SUN PRODUCTS

EM27/SUN measurements have undergone correction with air mass dependent (ADCF) and air mass independent (AICF) factors in order to bring COCCON data in agreement with TCCON (<https://www.imk-asf.kit.edu/>)

Species	MW	AICF $a_{\text{gas}}$	ADCF $b_{\text{gas}}$
XH <sub>2</sub> O	(8353.4, 8463.1)	0.8300	0.000
XAIR	(7765.0, 8005.0)	0.9737	−0.007
XCO <sub>2</sub>	(6173.0, 6390.0)	0.9862	0.005
XCH <sub>4</sub>	(5897.0, 6145.0)	0.9905	−0.014
XCO	(4208.7, 4318.8)	0.9250	0.103
XCH <sub>4</sub> <sup>(b)</sup>	(4208.7, 4318.8)	0.9727	−0.017

(b) XCH<sub>4</sub> S5P (Sentinel-5 Precursor)



## EM27/SUN PRODUCTS

X<sub>air</sub> is defined as

$$X_{\text{air}} = \frac{0.2095}{V_{\text{CO}_2} \cdot \bar{\mu}} \cdot \left( \frac{P_s}{g} - V_{\text{CH}_2\text{O}} \cdot \mu_{\text{H}_2\text{O}} \right)$$


where  $\mu$  and  $\mu_{\text{H}_2\text{O}}$  are molecular masses of dry air and water vapor  
 $g$  is column averaged gravitational acceleration,  $P_s$  is the surface pressure  
and  $V_{\text{CO}_2}$ ,  $V_{\text{CH}_2\text{O}}$  are the total columns of oxygen and water vapor

For an ideal measurement and retrieval  
with accurate O<sub>2</sub> and H<sub>2</sub>O spectroscopy and surface pressure,  
X<sub>air</sub> would be 1.

Due to a 2.0% bias in the O<sub>2</sub> spectroscopy,

- TCCON measurements X<sub>air</sub> is typically ~ 0.98 (Wunch et al., 2015).
- EM27/SUN measurements show a factor of ~ 0.97 (Frey et al., 2015; Hase et al., 2015; Klappenbach et al., 2015).

Large deviations (~ 1 %) from these values indicate severe problems,  
e.g., errors with the surface pressure, pointing errors, timing errors or  
changes in the optical alignment of the instrument. (Frey et al, 2019)



X<sub>air</sub> at Thessaloniki passes TCCON standard  
quality check (between 0.96 and 1.04) with a  
mean value of 0.996 and a STD of ~ 0.003



## EM27/SUN PRODUCTS

Two methane products of EM27/SUN measurements:


- in the 1.6 $\mu$ m band (standard product)
- in the S5P band (2.6 $\mu$ m) (not official)

are in good agreement, with a mean ratio of 1.0026  
( between 0.993 and 1.012 and a std <0.003)

Differences can be attributed to

- inconsistent spectroscopic linelists between bands
- different vertical sensitivity of the retrievals
- spectral interference of water vapor on the S5P band being stronger

A new linelist to be implemented into PROFFAST will be available soon, for an improved XCH<sub>4</sub> S5P band product (Frank Hase, personal





## EM27/SUN comparison to TROPOMI

Methane	Carbon monoxide
<p><u>TROPOMI</u>: XCH<sub>4</sub> OFFL product, no bias-corrected, qa=100</p> <p><u>EM27/SUN</u>: XCH<sub>4</sub> standard (1.6μm) XCH<sub>4</sub>S5P using S5P band (2.3μm)</p> <p>Mission data requirements for XCH<sub>4</sub>: 1.5% ±1%</p>	<p><u>TROPOMI</u>: XCO OFFL product, qa=100</p> <p><u>EM27/SUN</u>: XCO</p> <p>Mission data requirements for XCO: 15% ±10%</p>
100km & 1h co-location criteria <sup>1</sup>	50km & 1h co-location criteria <sup>1</sup>
<sup>1</sup> as used in TROPOMI Validation Reports	




## EM27/SUN vs TROPOMI full time series

Examining the full time series with no spatial/time criteria, TROPOMI seems to show

- higher CO values compared to ground based measurements
- lower XCH<sub>4</sub> values than the standard ground based XCH<sub>4</sub> and almost similar to the S5P band XCH<sub>4</sub>



Seasonal variations captured by EM27 are reproduced by TROPOMI





## EM27/SUN vs TROPOMI (co-located measurements)

### Methane ( $\text{XCH}_4$ EM27/SUN standard product)





$\% \text{ bias} = -0.0105\% \pm 0.502\%$ , well within mission requirements ( $1.5\% \pm 1.0\%$ )

Results in agreement with those of the Validation Reports of the Copernicus Sentinel-5 Precursor Operational Data Products that show TROPOMI 's underestimation of  $\text{XCH}_4$  vs NDACC and TCCON data with an overall agreement of  $-0.68\%$  (negative bias) for not bias-corrected TROPOMI  $\text{XCH}_4$  and a standard deviation of the relative bias is on an average  $0.6\%$



## EM27/SUN vs TROPOMI (co-located measurements)

**Methane (XCH<sub>4</sub> EM27/SUN standard product)**





% bias = -0.165% ± 0.546%, well within mission requirements (1.5% ± 1.0%)

Results in agreement with those of the Validation Reports of the Copernicus Sentinel-5 Precursor Operational Data Products that show TROPOMI 's underestimation of XCH<sub>4</sub> vs NDACC and TCCON data with an overall agreement of -0.68% (negative bias) for not bias-corrected TROPOMI XCH<sub>4</sub> and a standard deviation of the relative bias is on an average 0.6%



# EM27/SUN vs TROPOMI (co-located measurements)

## Carbon monoxide (XCO)



$\text{XCO } \% \text{ bias} = +2.41\% \pm 6.26\%$ , well within mission requirements ( $15\% \pm 10\%$ )

Results in agreement with those of the Validation Reports of the Copernicus Sentinel-5 Precursor Operational Data Products that show TROPOMI 's overestimation of XCO vs NDACC and TCCON data with an overall agreement of 7-10% (positive bias) and a standard deviation of the relative bias is on an average 5%



## EM27/SUN vs TROPOMI comparison for other distances

Distance (km)	XCH <sub>4</sub> standard (1h, qa=1) %bias	XCO (1h, qa=1) % bias
30	-0.0946	+4.416
50	+0.2088	+2.411
100	-0.0105	+0.736

- As the distance from Thessaloniki overpass increases, the bias decreases demonstrating a lower value of TROPOMI average value for this area, probably caused by fewer CO sources in the surrounding area of Thessaloniki, which is a high polluted urban site.
- For methane, it is not clear whether conclusions can be drawn mainly due to a very low number of common measurement days between EM27/SUN and TROPOMI. A detailed recording of CO and CH<sub>4</sub> sources in the area of Thessaloniki and within 100km, could provide more insight



## Summary

- The very first year of measurements of  $\text{XCH}_4$  and  $\text{XCO}$  using a Bruker EM27/SUN low resolution spectrometer provided by Karlsruhe Institute of Technology, are presented for Thessaloniki, an urban site in Northern Greece. Our measurements show similar levels of  $\text{XCH}_4$  and  $\text{XCO}$  for similar sites

- Comparison to TROPOMI measurements show that seasonal variations are captured. Mission requirements are met for both the bias level and the standard deviation:
  - $\Rightarrow \text{XCH}_4$  (1 hour, 100km) : % bias =  $-0.0105\% \pm 0.502\%$  (within  $1.5\% \pm 1.0\%$ )
  - $\Rightarrow \text{XCO}$  (1 hour, 50km) : % bias =  $+2.41\% \pm 6.26\%$  (within  $15\% \pm 10\%$ )

**(these are revised results compared to the positive TROPOMI bias for methane stated in the abstract)**

- Data set will be re evaluated after incorporating improved parameters such as ME and PE for ILS function and new linelists for secondary  $\text{XCH}_4$  products
- Important to establish a dense data set of ground based measurements in order to
  - to confirm the comparison to TROPOMI in the future
  - provide more common days for comparison to the satellite (especially for methane)
  - explore variability of the measurements



## References

Frey, M., Hase, F., Blumenstock, T., Groß, J., Kiel, M., Mengistu Tsidu, G., Schäfer, K., Sha, M. K., and Orphal, J.: **Calibration and instrumental line shape characterization of a set of portable FTIR spectrometers for detecting greenhouse gas emissions**, Atmos. Meas. Tech., 8, 3047–3057, <https://doi.org/10.5194/amt-8-3047-2015>, 2015

Frey, M., Mahesh K. Sha , Hase, F., Kiel, M., Blumenstock, T., Harig, R., Surawicz G., Deutscher, N.M., Shiomi, K., Franklin, J.E., Bösch, H., Chen, J., Grutter, M., Ohyama, H., Sun, Y., Butz, A., Mengistu Tsidu, G., Ene, D., Wunch, D., Cao, Z., Garcia, O., Ramonet, M., Vogel F., and Orphal, J., **Building the COLlaborative Carbon Column Observing Network (COCCON): long-term stability and ensemble performance of the EM27/SUN Fourier transform spectrometer**, Atmos. Meas. Tech., 12, 1513–1530, 2019, <https://doi.org/10.5194/amt-12-1513-2019>

Gisi, M., Hase, F., Dohe, S., Blumenstock, T., Simon, A., and Keens, A.: XCO<sub>2</sub>-measurements with a tabletop FTS using solar absorption spectroscopy, Atmos. Meas. Tech., 5, 2969–2980, <https://doi.org/10.5194/amt-5-2969-2012>, 2012.

Hasekamp, O., Lorente, A., Hu, H., Butz, A., Aan de Brugh, J., Landgraf, J.: **ATBD for Sentinel-5 Precursor Methane Retrieval**, SRON-S5P-LEV2-RP-001

Hedelius, J. K., Parker, H., Wunch, D., Roehl, C. M., Viatte, C., Newman, S., Toon, G. C., Podolske, J. R., Hillyard, P. W., Iraci, L. T., Dubey, M. K., and Wennberg, P. O.: Intercomparability of XCO<sub>2</sub> and XCH<sub>4</sub> from the United States TCCON sites, Atmos. Meas. Tech., 10, 1481–1493, <https://doi.org/10.5194/amt10-1481-2017>, 2017

Tu, Qiansi, **Observation of atmospheric greenhouse gas abundances on regional scales in boreal areas using portable FTIR Spectrometers**, Dissertation

Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: **The Total Carbon Column Observing Network**, Philos. T. R. Soc. A, 369, 2087–2112, <https://doi.org/10.1098/rsta.2010.0240>, 2011

Lambert, J.-C., S. Compernelle, K.-U. Eichmann, M. de Graaf, D. Hubert, A. Keppens, Q. Kleipool, B. Langerock, M.K. Sha, T. Verhoelst, T. Wagner, C. Ahn, A. Argyrouli, D. Balis, K.L. Chan, I. De Smedt, H. Eskes, A.M. Fjæraa, K. Garane, J.F. Gleason, F. Goutail, J. Granville, P. Hedelt, K.-P. Heue, G. Jaross, M.L. Koukouli, J. Landgraf, R. Lutz, S. Nanda, S. Niemejer, A. Pazmiño, G. Pinardi, J.-P. Pommereau, A. Richter, N. Rozemeijer, M. Sneep, D. Stein Zweers, N. Theys, G. Tilstra, O. Torres, P. Valks, C. Vigouroux, P. Wang, and M. Weber. , **Quarterly Validation Report of the Copernicus Sentinel-5 Precursor Operational Data Products #06: April 2018 – February 2020**, S5P MPC Routine Operations Consolidated Validation Report series, Issue #06, Version 06.0.1, 154 pp., March 2020.



## Acknowledgements

This work was funded by ESA within the Contract No. 4000117151/16/l-LG “Preparation and Operations of the Mission Performance Centre (MPC) for the Copernicus Sentinel-5 Precursor Satellite”. The satellite data were obtained through Sentinel-5P Expert Users Data Hub (<https://s5pexp.copernicus.eu/>).

We acknowledge support of this work by the project “PANhellenic infrastructure for Atmospheric Composition and climatE change” (MIS 5021516) which is implemented under the Action “Reinforcement of the Research and Innovation Infrastructure”, funded by the Operational Programme "Competitiveness, Entrepreneurship and Innovation" (NSRF 2014-2020) and co-financed by Greece and the European Union (European Regional Development Fund).

We wholeheartedly acknowledge the support provided by the IT Center of the Aristotle University of Thessaloniki (AUTh) throughout the progress of this research work.

We wholeheartedly acknowledge the support provided by the Atmospheric Toolbox.