

*EGU 2020, 4-8 May 2020, Session AS3.19: New (Sentinel-5 Precursor) and Evolving (e.g. Sentinel-4)  
Capabilities to Measure Atmospheric Composition from Space*

# Sentinel-5 Precursor methane and carbon monoxide column retrievals and assessments related to localized emission sources

*Using S5P scientific WFM-DOAS retrievals*

**Michael Buchwitz**, Oliver Schneising, Stefan Noël, Maximilian Reuter, Steffen Vanselow,  
Heinrich Bovensmann, John P. Burrows

Institute of Environmental Physics (IUP), Institute of Remote Sensing (IFE)  
University of Bremen, Bremen, Germany

# Outline

- Essentially overview 3 recent publications covering
  - Retrieval algorithm, products, validation
  - Selected results, e.g.,
    - CO: Californian fires Nov 2018
    - CH<sub>4</sub>: Emissions from major natural gas and petroleum production fields

<https://doi.org/10.5194/acp-2020-274>  
Preprint. Discussion started: 14 April 2020  
© Author(s) 2020. CC BY 4.0 License.



3

Atmospheric  
Chemistry  
and Physics  
Discussions  
Open Access  
EGU

Schneising et al., ACP (in review), 2020

## Remote sensing of methane leakage from natural gas and petroleum systems revisited

Oliver Schneising, Michael Buchwitz, Maximilian Reuter, Steffen Vanselow, Heinrich Bovensmann, and John P. Burrows

Atmos. Meas. Tech., 12, 6771–6802, 2019  
<https://doi.org/10.5194/amt-12-6771-2019>  
© Author(s) 2019. This work is distributed under the Creative Commons Attribution 4.0 License.



1

Atmospheric  
Measurement  
Techniques  
Open Access  
EGU

Schneising et al., AMT, 2019

## A scientific algorithm to simultaneously retrieve carbon monoxide and methane from TROPOMI onboard Sentinel-5 Precursor

Oliver Schneising<sup>1</sup>, Michael Buchwitz<sup>1</sup>, Maximilian Reuter<sup>1</sup>, Heinrich Bovensmann<sup>1</sup>, John P. Burrows<sup>1</sup>, Tobias Borsdorff<sup>2</sup>, Nicholas M. Deutscher<sup>3</sup>, Dietrich G. Feist<sup>4,5,6</sup>, David W. T. Griffith<sup>3</sup>, Frank Hase<sup>7</sup>, Christian Hermans<sup>8</sup>, Laura T. Iraci<sup>9</sup>, Rigel Kivi<sup>10</sup>, Jochen Landgraf<sup>2</sup>, Isamu Morino<sup>11</sup>, Justus Notholt<sup>1</sup>, Christof Petri<sup>1</sup>, David F. Pollard<sup>12</sup>, Sébastien Roche<sup>13</sup>, Kei Shiomi<sup>14</sup>, Kimberly Strong<sup>13</sup>, Ralf Sussmann<sup>15</sup>, Voltaire A. Velasco<sup>3</sup>, Thorsten Warneke<sup>1</sup>, and Debra Wunch<sup>13</sup>

Atmos. Chem. Phys., 20, 3317–3332, 2020  
<https://doi.org/10.5194/acp-20-3317-2020>  
© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



2

Atmospheric  
Chemistry  
and Physics  
Open Access  
EGU

Schneising et al., ACP, 2020

## Severe Californian wildfires in November 2018 observed from space: the carbon monoxide perspective

Oliver Schneising, Michael Buchwitz, Maximilian Reuter, Heinrich Bovensmann, and John P. Burrows  
Institute of Environmental Physics (IUP), University of Bremen FB1, Bremen, Germany

# WFM-DOAS (or WFMD) algorithm for S5P


Atmos. Meas. Tech., 12, 6771–6802, 2019  
<https://doi.org/10.5194/amt-12-6771-2019>  
 © Author(s) 2019. This work is distributed under the Creative Commons Attribution 4.0 License.

Atmospheric  
Measurement  
Techniques  
Open Access  
EGU


Schneising et al., AMT, 2019

## A scientific algorithm to simultaneously retrieve carbon monoxide and methane from TROPOMI onboard Sentinel-5 Precursor


Oliver Schneising<sup>1</sup>, Michael Buchwitz<sup>1</sup>, Maximilian Reuter<sup>1</sup>, Heinrich Bovensmann<sup>1</sup>, John P. Burrows<sup>1</sup>, Tobias Borsdorff<sup>2</sup>, Nicholas M. Deutscher<sup>3</sup>, Dietrich G. Feist<sup>4,5,6</sup>, David W. T. Griffith<sup>3</sup>, Frank Hase<sup>7</sup>, Christian Hermans<sup>8</sup>, Laura T. Iraci<sup>9</sup>, Rigel Kivi<sup>10</sup>, Jochen Landgraf<sup>2</sup>, Isamu Morino<sup>11</sup>, Justus Notholt<sup>1</sup>, Christof Petri<sup>1</sup>, David F. Pollard<sup>12</sup>, Sébastien Roche<sup>13</sup>, Kei Shiomi<sup>14</sup>, Kimberly Strong<sup>13</sup>, Ralf Sussmann<sup>15</sup>, Voltaire A. Velasco<sup>3</sup>, Thorsten Warneke<sup>1</sup>, and Debra Wunch<sup>13</sup>



## Data flow



## Averaging kernels



## Method:

- Least-squares fit of simulated radiances to S5P radiances
- Very fast Look-Up-Table (LUT) scheme
- Quality flagging (e.g., clouds) and bias correction (only for methane) via Machine Learning (Random Forest; VIIRS for clouds; climatology for methane)

## Products:

- CO columns [molec./cm<sup>2</sup>] and XCO [ppb]; XCH<sub>4</sub> [ppb]

## Differences w.r.t. operational algorithms / products:

- Many as independent algorithms ..., e.g., resulting products:
  - WFMD XCH<sub>4</sub>: Typically better coverage (incl. some ocean coverage)
  - WFMD CO: Cloud-free only, XCO in addition to CO column, ...

# S5P/WFMD products: Validation


Atmos. Meas. Tech., 12, 6771–6802, 2019  
<https://doi.org/10.5194/amt-12-6771-2019>  
 © Author(s) 2019. This work is distributed under the Creative Commons Attribution 4.0 License.




Schneising et al., AMT, 2019

## A scientific algorithm to simultaneously retrieve carbon monoxide and methane from TROPOMI onboard Sentinel-5 Precursor


Oliver Schneising<sup>1</sup>, Michael Buchwitz<sup>1</sup>, Maximilian Reuter<sup>1</sup>, Heinrich Bovensmann<sup>1</sup>, John P. Burrows<sup>1</sup>, Tobias Borsdorff<sup>2</sup>, Nicholas M. Deutscher<sup>3</sup>, Dietrich G. Feist<sup>4,5,6</sup>, David W. T. Griffith<sup>3</sup>, Frank Hase<sup>7</sup>, Christian Hermans<sup>8</sup>, Laura T. Iraci<sup>9</sup>, Rigel Kivi<sup>10</sup>, Jochen Landgraf<sup>2</sup>, Isamu Morino<sup>11</sup>, Justus Notholt<sup>1</sup>, Christof Petri<sup>1</sup>, David F. Pollard<sup>12</sup>, Sébastien Roche<sup>13</sup>, Kei Shiomi<sup>14</sup>, Kimberly Strong<sup>13</sup>, Ralf Sussmann<sup>15</sup>, Voltaire A. Velasco<sup>3</sup>, Thorsten Warneke<sup>1</sup>, and Debra Wunch<sup>13</sup>



## Comparisons with TCCON



**XCH<sub>4</sub>:**  
 Global offset: -1.4 ppb  
 Random (sgl.obs., 1-sigma): 14 ppb  
 Systematic (site-to-site StdDev): 4.4 ppb



**XCO:**  
 Global offset: 4.5 ppb  
 Random (sgl.obs., 1-sigma): 5.1 ppb  
 Systematic (site-to-site StdDev): 1.9 ppb



# S5P/WFMD products: Comparison with operational products





Atmos. Meas. Tech., 12, 6771–6802, 2019  
<https://doi.org/10.5194/amt-12-6771-2019>  
 © Author(s) 2019. This work is distributed under  
 the Creative Commons Attribution 4.0 License.

Atmospheric  
Measurement  
Techniques  


Schneising et al., AMT, 2019

## A scientific algorithm to simultaneously retrieve carbon monoxide and methane from TROPOMI onboard Sentinel-5 Precursor

Oliver Schneising<sup>1</sup>, Michael Buchwitz<sup>1</sup>, Maximilian Reuter<sup>1</sup>, Heinrich Bovensmann<sup>1</sup>, John P. Burrows<sup>1</sup>, Tobias Borsdorff<sup>2</sup>, Nicholas M. Deutscher<sup>3</sup>, Dietrich G. Feist<sup>4,5,6</sup>, David W. T. Griffith<sup>3</sup>, Frank Hase<sup>7</sup>, Christian Hermans<sup>8</sup>, Laura T. Iraci<sup>9</sup>, Rigel Kivi<sup>10</sup>, Jochen Landgraf<sup>2</sup>, Isamu Morino<sup>11</sup>, Justus Notholt<sup>1</sup>, Christof Petri<sup>1</sup>, David F. Pollard<sup>12</sup>, Sébastien Roche<sup>13</sup>, Kei Shiomi<sup>14</sup>, Kimberly Strong<sup>13</sup>, Ralf Sussmann<sup>15</sup>, Voltaire A. Velasco<sup>3</sup>, Thorsten Warneke<sup>1</sup>, and Debra Wunch<sup>13</sup>



- **Overall** reasonable to good agreement
- **CH<sub>4</sub>**: WFMD typically better coverage (e.g., also some ocean coverage but also over land)
- **CO**: WFMD much sparser (cloud-free only)

# S5P/WFMD CO: Some details ...

Atmos. Meas. Tech., 12, 6771–6802, 2019  
<https://doi.org/10.5194/amt-12-6771-2019>  
 © Author(s) 2019. This work is distributed under  
 the Creative Commons Attribution 4.0 License.


Atmospheric  
 Measurement  
 Techniques  
 Open Access  
 EGU

Schneising et al., AMT, 2019

**A scientific algorithm to simultaneously retrieve carbon monoxide  
 and methane from TROPOMI onboard Sentinel-5 Precursor**


Oliver Schneising<sup>1</sup>, Michael Buchwitz<sup>1</sup>, Maximilian Reuter<sup>1</sup>, Heinrich Bovensmann<sup>1</sup>, John P. Burrows<sup>1</sup>,  
 Tobias Borsdorff<sup>2</sup>, Nicholas M. Deutscher<sup>3</sup>, Dietrich G. Feist<sup>4,5,6</sup>, David W. T. Griffith<sup>3</sup>, Frank Hase<sup>7</sup>,  
 Christian Hermans<sup>8</sup>, Laura T. Iraci<sup>9</sup>, Rigel Kivi<sup>10</sup>, Jochen Landgraf<sup>2</sup>, Isamu Morino<sup>11</sup>, Justus Notholt<sup>1</sup>,  
 Christof Petri<sup>1</sup>, David F. Pollard<sup>12</sup>, Sébastien Roche<sup>13</sup>, Kei Shiomi<sup>14</sup>, Kimberly Strong<sup>13</sup>, Ralf Sussmann<sup>15</sup>,  
 Voltaire A. Velasco<sup>3</sup>, Thorsten Warneke<sup>1</sup>, and Debra Wunch<sup>13</sup>

TROPOMI/WFMD XCO 20180918




## Elevated CO due to various CO emission hotspots (urban areas, steel plants, ...)


(a) TROPOMI/WFMD XCO Nov–Dec 2018



TROPOMI/WFMD XCO 20181104



TROPOMI/WFMD XCO 20180606





# S5P/WFMD CO: Californian fires Nov 2018

Atmos. Chem. Phys., 20, 3317–3332, 2020  
https://doi.org/10.5194/acp-20-3317-2020  
© Author(s) 2020. This work is distributed under  
the Creative Commons Attribution 4.0 License.



Schneising et al., ACP, 2020


Atmospheric  
Chemistry  
and Physics  
Open Access  
EGU

## Severe Californian wildfires in November 2018 observed from space: the carbon monoxide perspective


Oliver Schneising, Michael Buchwitz, Maximilian Reuter, Heinrich Bovensmann, and John P. Burrows

Institute of Environmental Physics (IUP), University of Bremen FB1, Bremen, Germany

### Comparisons with CAMS, use of ERA5, ...



### S5P/WFMD CO: Elevated CO due to fires



### Main air quality / health-related conclusion:

Even the most polluted city scenes likely comply with the national ambient air quality standards (10 mgCO/m<sup>3</sup> with 8 h averaging time). This finding based on dense daily recurrent satellite monitoring is consistent with isolated ground-based air quality measurements.

# S5P/WFMD XCH<sub>4</sub>: A GHG-CCI product

<http://cci.esa.int/ghg>

climate change initiative

esa

Home

Greenhouse Gases CCI project

**→ GET CCI DATA**



A single point of access. Open. Free. Easy.  
[cci.esa.int/data](http://cci.esa.int/data)

**greenhouse gases cci**

Objective | Team | Data and documentation | Publications | Images | Contact us

**Summary**  
Carbon dioxide (CO<sub>2</sub>) and methane (CH<sub>4</sub>) are the two most important anthropogenic ("human-made") greenhouse gases (GHG) driving global climate change. Increasing atmospheric concentrations of these Essential Climate Variables (ECVs) leads to global warming with adverse consequences such as rising sea levels. It is therefore important to monitor the spatial distribution and the time evolution of these gases and to improve our knowledge of their various natural and anthropogenic sources and sink.

- aerosol cci
- biomass cci
- cloud cci
- cmug cci
- fire cci
- ghg cci
- glaciers cci
- antarctic ice sheet cci
- ice sheets greenland cci
- land cover cci
- high resolution land cover cci
- lakes cci



## APPLICATIONS

**Satellites providing clear picture of greenhouse gases**

[https://www.esa.int/Applications/Observing\\_the\\_Earth/Space\\_for\\_our\\_climate/Satellites\\_providing\\_clear\\_picture\\_of\\_greenhouse\\_gases](https://www.esa.int/Applications/Observing_the_Earth/Space_for_our_climate/Satellites_providing_clear_picture_of_greenhouse_gases)



# S5P/WFMD XCH<sub>4</sub>: Methane emissions from gas & oil fields

<https://doi.org/10.5194/acp-2020-274>  
Preprint. Discussion started: 14 April 2020  
© Author(s) 2020. CC BY 4.0 License.




Atmospheric  
Chemistry  
and Physics  
Discussions  
EGU

Schneising et al., ACP (in review), 2020


Remote sensing of methane leakage from natural gas and petroleum systems revisited

Oliver Schneising, Michael Buchwitz, Maximilian Reuter, Steffen Vanselow, Heinrich Bovensmann, and John P. Burrows


## Major gas & oil production regions



Permian



## USA: Methane Leakage



- Use of S5P WFMD XCH<sub>4</sub> to get emission estimates
- Conversion to relative leakage by division with total production
- Comparison with published values

# S5P/WFMD XCH<sub>4</sub>: Methane emissions from gas & oil fields

<https://doi.org/10.5194/acp-2020-274>  
Preprint. Discussion started: 14 April 2020  
© Author(s) 2020. CC BY 4.0 License.






Atmospheric  
Chemistry  
and Physics  
Discussions  
EGU

Schneising et al., ACP (in review), 2020

## Remote sensing of methane leakage from natural gas and petroleum systems revisited

Oliver Schneising, Michael Buchwitz, Maximilian Reuter, Steffen Vanselow, Heinrich Bovensmann, and John P. Burrows

## Major gas & oil production regions



**Table 1.** Summary of the emission and production values used to determine the leakage rates (emissions divided by combined oil and gas production). All values have been converted to  $\text{kBOE d}^{-1}$  as described in Section 2. Also shown are the mean percentage variance contributions to the emission estimates for the relative uncertainty components of Equation 2.

Region	Emissions ( $\text{kBOE d}^{-1}$ )	Oil		Gas		Oil+Gas ( $\text{kBOE d}^{-1}$ )	Leakage (%)	Variance contributions (%)			
		( $\text{kBOE d}^{-1}$ )	(%)	( $\text{kBOE d}^{-1}$ )	(%)			<i>E</i>	<i>v</i> , abs	<i>v</i> , dir	$\rho_{dry}$
Permian	81	3897	64	2197	36	6094	1.3	60.2	38.6	0.6	0.6
Appalachia	60	127	2	5052	98	5179	1.2	73.2	26.4	0.2	0.2
Eagle Ford	39	1344	55	1112	45	2456	1.6	64.3	34.7	0.5	0.5
Bakken	23	1361	75	444	25	1805	1.3	64.9	34.5	0.4	0.2
Anadarko	72	548	31	1237	69	1785	4.0	70.8	28.4	0.4	0.4
Galkynysh/ Dauletabad	83	0	0	1533	100	2017	4.1	74.9	21.8	0.5	2.8

# Summary & conclusions

- Scientific WFM-DOAS (or WFMD) algorithm to retrieve  $\text{XCH}_4$  and XCO from TROPOMI/S5P
- Access to data products (free of charge):
  - S5P WFMD XCO: [https://www.iup.uni-bremen.de/carbon\\_ghg/products/tropomi\\_wfmd/](https://www.iup.uni-bremen.de/carbon_ghg/products/tropomi_wfmd/)
  - S5P WFMD  $\text{XCH}_4$ : <http://cci.esa.int/data>
- Estimated quality of WFMD products (relative to TCCON): Single observation uncertainty (= root-sum-square of sgl.obs. random and site-to-site systematic differences):
  - $\text{XCH}_4$ : ~15 ppb (~1%); excluding 1.4 ppb global low bias
  - XCO: ~5 ppb (~5%); excluding 4.5 ppb global high bias
- Comparison of WFMD & OPERational products:
  - Overall good agreement
  - Coverage may be significantly different
- As shown in this presentation, we started to use these information rich products to obtain information on various (localized) methane and CO emission sources
- See also: [https://www.iup.uni-bremen.de/carbon\\_ghg/](https://www.iup.uni-bremen.de/carbon_ghg/)



# Acknowledgements

- Funding:
  - ESA (projects GHG-CCI/GHG-CCI+, S5L2PP, Methane+)
  - German BMBF (project AIRSPACE)
  - State & University of Bremen
- Data:
  - Copernicus (EU / ESA) TROPOMI/S5P products:
    - L1b
    - L2 CH<sub>4</sub> & CO
    - VIIRS-based L2 cloud product for TROPOMI
  - Ground-based validation data: TCCON
  - Meteorological data: ECMWF
  - Other: see publications