



Ex-hurricane Ophelia and air quality impacts over Europe in CAMS forecast systems

Dimitris Akritidis¹, Eleni Katragkou¹, Aristeidis K. Georgoulas¹, Prodromos Zanis¹,
Sergios Kartsios¹, Johannes Flemming², Antje Inness², and Henk Eskes³

¹ DEPARTMENT OF METEOROLOGY AND CLIMATOLOGY, SCHOOL OF GEOLOGY, ARISTOTLE UNIVERSITY OF THESSALONIKI, THESSALONIKI, GREECE

² EUROPEAN CENTRE FOR MEDIUM-RANGE WEATHER FORECASTS (ECMWF), READING, UK

³ ROYAL NETHERLANDS METEOROLOGICAL INSTITUTE (KNMI), DE BILT, THE NETHERLANDS

e-mail: dakritid@geo.auth.gr

The **Copernicus Atmosphere Monitoring Service (CAMS)**¹ is one of the six services that **Copernicus** provides, using a comprehensive global assimilation and forecasting system to **assess the state and composition of the atmosphere** on a daily basis.



CAMS global system

- Based on the European Centre for Medium-Range Weather Forecasts (ECMWF) Integrated Forecasting System (IFS).
- Horizontal resolution of about 40 km and 60 vertical levels reaching up to 0.1 hPa.
- Assimilation of aerosol, ozone, nitrogen dioxide, carbon monoxide, and sulfur dioxide using the 4D-VAR data assimilation system (Inness et al., 2015, 2019)
- An extended version of the Carbon Bond 2005 (CB05) chemical mechanism is used (Flemming et al., 2015) and the Morcrette et al. (2009) aerosol parameterization.




CAMS regional system

- The CAMS regional forecast system provides the ensemble median (RegEns) and members of the European-scale air quality forecasts up to 4 days in advance.
- The products have a 0.1° horizontal resolution and are based on state-of-the-art numerical air quality models developed in Europe: CHIMERE, EMEP, EURAD-IM, LOTOS-EUROS, MATCH, MOCAGE and SILAM.
- CAMS regional models assimilate PM₁₀ and PM_{2.5} surface observations but not satellite aerosol products.


Data

- We use CAMS-global (IFS) and CAMS-regional (RegEns) day-1 forecast data.
- AOD₅₅₀ data from the two MODIS sensors aboard EOS Terra (equator crossing time 10:30 LT) and Aqua (equator crossing time 13:30 LT) satellites are used (Levy et al., 2013; Sayer et al., 2015).
- Also used are aerosol subtype data from CALIOP/CALIPSO (equator crossing time 13:30 LT) (Kim et al., 2018).
- Ground-based observations of PM₁₀ and PM_{2.5} surface concentrations are used from rural background stations, obtained from the European Monitoring and Evaluation Programme (EMEP) (Tørseth et al., 2012).

¹<https://atmosphere.copernicus.eu/>




MODIS AOD₅₅₀ at 16/10/2017




- On October 16th, high AOD₅₅₀ values are observed over North and Baltic Sea, France, Belgium, the Netherlands and Germany.

IFS AOD₅₅₀ at 12Z 16/10/2017




- The dust AOD contribution over Great Britain (GB) ranges with a percentage between 20-35%, while over the North Sea and the Netherlands it reaches values up to 65%.
- The biomass burning emission trails originating from Portugal are also a significant input for AOD, with black carbon and organic matter aerosols contributing up to 20% and 40%, respectively, over GB.

Percentage contribution to the IFS AOD₅₅₀ at 12Z 16/10/2017




CALIOP/CALIPSO AEROSOL SUBTYPES



- Dust and polluted dust are detected at 45°N-50°N and 30°N-35°N up to 3 km.
- Dust is also observed in the upper troposphere (10 km) over France.
- Elevated smoke is identified over the North Sea at 4 km.


IFS DOMINANT AEROSOL TYPE



Vertical cross-sections of CALIOP/CALIPSO aerosol subtypes and IFS dominant aerosol type on 12Z of October 17th 2017. The embedded map depicts the respective CALIOP/CALIPSO track.


- The latitudinal bands of IFS dust aerosol dominance are consistent with CALIPSO, yet, fire-originated aerosols (carbonaceous aerosols) are found over the North Sea higher (above 5 km) compared to CALIPSO. This might be due to sea salt aerosol overestimation that prevents carbonaceous aerosols to appear as the dominant type below 5 km.

Impacts on air-quality (PM₁₀)



Observed (black), IFS (blue), IFS no DA (no Data Assimilation) (grey) and RegEns (red) 3-hourly timeseries of PM₁₀ surface concentrations (ug/m³) (top), and percentage (%) contribution of aerosol type to IFS PM₁₀ surface concentrations (bottom) at Guipry (FR0024R), France.

- From 12Z of October 14th and up to October 16th an **increase in PM₁₀ surface concentrations is observed**, also seen in IFS and RegEns.
- However, **the early peak in observations is not captured by CAMS forecast systems**.
- The percentage contribution of each IFS aerosol type to IFS PM₁₀ concentrations indicates that the enhancement in PM₁₀ surface levels is initially (00Z on 15 October) **due to organic matter aerosols (75%)**, probably related to transported smoke from fires burning over the Iberian Peninsula, while afterwards (21Z on 15 October) the **transport of dust induces a significant contribution from dust aerosols (53%)** as well.
- On 12Z of October 16th a **sharp peak is depicted in observed PM₁₀ concentrations** reproduced from both IFS and RegEns, yet overestimated from IFS, which seems to result from **sea salt transport from the west** and on the south of Ophelia.
- IFS exhibits an increase of PM₁₀ levels on 12Z of October 18th due to dust and organic matter aerosols which is not seen in RegEns and observations.



Observed (black), IFS (blue), IFS no DA (no Data Assimilation) (grey) and RegEns (red) 3-hourly timeseries of PM_{2.5} surface concentrations (ug/m³) (top), and percentage (%) contribution of aerosol type to IFS PM_{2.5} surface concentrations (bottom) at Chilbolton Observatory, GB.

- After 15Z of October 15th the **observed PM_{2.5} surface concentrations increase up to 28 ug/m³**, with IFS peaking with a 6 hours delay.
- A second wave of dust and biomass burning transport is affecting PM_{2.5} surface concentrations over the site on 15Z of October 18th as revealed from IFS, which is in agreement with observations, but not so profound in RegEns.

- African dust was transported towards northwestern Europe on the east side of Storm Ophelia forming an extended dust plume reaching the Baltic States. On it's passage, Ophelia lifted and drifted smoke aerosols from fires burning over the Iberian Peninsula carrying them over several European regions, such as France, Great Britain and the Netherlands.
- After 00Z of October 16th, dust and smoke aerosols are found on the warm and cold front of Ophelia, being transported within it's warm sector.
- The decomposition of IFS AOD₅₅₀ fields indicates that African dust, organic matter from biomass burning over the Iberian Peninsula and sea salt aerosols sprayed in the vicinity of Ophelia, are the major contributors to the plumes of high AOD₅₅₀.
- The CALIOP/CALIPSO and the spatially and temporally collocated IFS cross sections of dominant aerosol types, reveal a good qualitative performance of IFS, as it generally manages to reproduce the type and location of aerosols during the passage of Ophelia.
- In terms of air quality, comparison with ground-based measurements from EMEP stations reveals that both IFS and RegEns are in general able to reproduce the observed signal of increase in PM₁₀ and PM_{2.5} surface concentrations, yet exhibiting some inconsistencies on the quantitative and temporal representation of the observed peaks.

References

- Flemming, J., Huijnen, V., Arteta, J., Bechtold, P., Beljaars, A., Blechschmidt, A.-M., Diamantakis, M., Engelen, R., Gaudel, A., Inness, A., et al.: Tropospheric chemistry in the Integrated Forecasting System of ECMWF, Geoscientific model development, 8, 975–1003, 2015.
- Inness, A., Blechschmidt, A.-M., Bouarar, I., Chabrilat, S., Crepulja, M., Engelen, R. J., Eskes, H., Flemming, J., Gaudel, A., Hendrick, F., Huijnen, V., Jones, L., Kapsomenakis, J., Katragkou, E., Keppens, A., Langerock, B., de Mazière, M., Melas, D., Parrington, M., Peuch, V. H., Razinger, M., Richter, A., Schultz, M. G., Suttie, M., Thouret, V., Vrekoussis, M., Wagner, A., and Zerefos, C.: Data assimilation of satellite-retrieved ozone, carbon monoxide and nitrogen dioxide with ECMWF's Composition-IFS, Atmospheric chemistry and physics, 15, 5275–5303, 2015.
- Inness, A., Flemming, J., Heue, K.-P., Lerot, C., Loyola, D., Ribas, R., Valks, P., van Roozendaal, M., Xu, J., and Zimmer, W.: Monitoring and assimilation tests with TROPOMI data in the CAMS system: near-real-time total column ozone, Atmospheric Chemistry and Physics, 35 19, 3939–3962, <https://doi.org/10.5194/acp-19-3939-2019>, 2019.
- Kim, M.-H., Omar, A. H., Tackett, J. L., Vaughan, M. A., Winker, D. M., Trepte, C. R., Hu, Y., Liu, Z., Poole, L. R., Pitts, M. C., Kar, J., and Magill, B. E.: The CALIPSO version 4 automated aerosol classification and lidar ratio selection algorithm, Atmospheric Measurement Techniques, 11, 6107–6135, <https://www.atmos-meas-tech.net/11/6107/2018/>, 2018.
- Levy, R. C., Mattoo, S., Munchak, L. A., Remer, L. A., Sayer, A. M., Patadia, F., and Hsu, N. C.: The Collection 6 MODIS aerosol products over land and ocean, Atmospheric Measurement Techniques, 6, 2989–3034, <https://www.atmos-meas-tech.net/6/2989/2013/>, 2013.
- Morcrette, J.-J., Boucher, O., Jones, L., Salmond, D., Bechtold, P., Beljaars, A., Benedetti, A., Bonet, A., Kaiser, J., Razinger, M., et al.: Aerosol analysis and forecast in the European Centre for medium-range weather forecasts integrated forecast system: Forward modeling, Journal of Geophysical Research: Atmospheres, 114, <https://doi.org/10.1029/2008JD011235>, 2009.
- Sayer, A. M., Hsu, N. C., Bettenhausen, C., Jeong, M.-J., and Meister, G.: Effect of MODIS Terra radiometric calibration improvements on Collection 6 Deep Blue aerosol products: Validation and Terra/Aqua consistency, Journal of Geophysical Research: Atmospheres, 120, 12,157–12,174, <https://doi.org/10.1002/2015JD023878>, 2015.
- Tørseth, K., Aas, W., Breivik, K., Fjæraa, A. M., Fiebig, M., Hjellbrekke, A. G., Lund Myhre, C., Solberg, S., and Yttri, K. E.: Introduction to the European Monitoring and Evaluation Programme (EMEP) and observed atmospheric composition change during 1972–2009, Atmospheric Chemistry and Physics, 12, 5447–5481, <https://doi.org/10.5194/acp-12-5447-2012>, <https://www.atmos-chem-phys.net/12/5447/2012/>, 2012.

This work is performed within the framework of the service element "CAMS84: Global and regional a posteriori evaluation and quality assurance" of the Copernicus Atmospheric Monitoring Services (CAMS).