

Non-radiative heat dissipation across scales in a water stressed pine forest: From the leaf to the planetary boundary layer

Dan Yakir, Jonathan Muller, Fyodor Tatatrinov, Mathias Mauder, and Eyal Rotenberg

Weizmann Institute of Science, Earth & Planetary Sciences, Rehovot, Israel
Institute of Meteorology and Climate Research, Garmisch-Partenkirchen, Germany



D546 | EGU2020-11279 BG3.25

Boundary layer scale: The 'convector effect'

With suppressed latent heat flux (LE) because of lack of water, the forest is transformed into an effective "convector" that exploits the low tree density and open canopy and, consequently, high canopy-atmosphere aerodynamic coupling.

Contribution of Semi-Arid Forests to the Climate System

Eyal Rotenberg, Dan Yakir*

Science 22 Jan 2010:

Vol. 327, Issue 5964, pp. 451-454

DOI: 10.1126/science.1179998



Water-cooled canopy...




Air-cooled canopy...



Forest cooling -5C annual mean

High roughness \rightarrow low r_h

$$H = \rho_a C_a \frac{(T - T_a)}{r_h}$$





- Latent heat flux (LE) is minor
- Sensible heat (H) balance net radiation R_n



Jonathan Muller


Tree scale 1: Sensible heat from the hottest part of the canopy




Sensible heat (H) production peak where leaf to air temperature difference is maximal

Tree scale 2: Buoyancy below canopy, mech. shear at the top


Measuring flux profiles




The main components of TKE: Mech. shear is dominant at the top of canopy (yellow); decreasing up (blue), and down (black) where Buoyancy becomes relatively more significant



Leaf scale 1: High precision leaf temperature measurements



Measurement methods for leaf temperature



$$L_{tot} = \tau \cdot \epsilon_{obj} \cdot L_{obj} + \tau \cdot (1 - \epsilon_{obj}) \cdot L_{bg} + (1 - \tau) \cdot L_{atm}$$

Quantity of interest

Reference plate with high reflectivity


Ignored due to small distance





Leaf scale 2: Leaf temperature independent of evaporation

- Leaf temperature similar with x10 rate of evaporation
- Efficient sensible heat production at the leaf scale
- Needle leaf: low heat capacity, large surface area, low density

Leaf temp management to protect the biochemistry



Leaf to air temperature difference



$$H = -c_p \cdot \rho \cdot \frac{T_a - T_s}{R_H}, \text{ also at the leaf scale}$$

Conclusions

- The "convective effect" evolves from the leaf, through tree, to the canopy scale
- Main source of sensible heat below top of canopy, with Buoyancy (B) production of turbulence below and Mech. Shear (MS) production above
- Leaf temperature can be controlled independently of evaporation rate
- At all scales, heat is dissipated via non evaporative and non-radiative processes
- Such processes infer resilience to forests under dry conditions
- Indicating potential adjustments in forest in regions undergoing drying and warming climatic trends

