

Eddy covariance and CRDS based techniques of GHGs measurements provide additional constraint in partitioning the net ecosystem exchange

Supriyo Chakraborty, Abirlal Metya, Amey Datye
Pramit K. Deb Burman
Indian Institute of Tropical Meteorology
Pune, India


Dipankar Sarma, Nirmali Gogoi, Abhijit Bora
Tezpur University, Tezpur, India

Abstract: EGU2020-901
<https://doi.org/10.5194/egusphere-egu2020-901>

EGU-2020 General Assembly:
BG Division Meeting
Chat session: May 7, 2020

Eddy-covariance based CO₂ flux estimation in north-east India





IRGA Sensor; Licor 7200 measures CO₂ and water vapor fluxes

Diurnal variation of the Net Ecosystem Exchange on a seasonal timescale at the KNP, India

Sarma et al. 2018, Deb Burman et al. 2019

Partitioning of the CO₂ fluxes into photosynthetic and respiration components

$$NEE = R_{eco} + R_A$$

R_{eco} = eco-system respiration

R_A = photo-synthetic uptake of carbon

Problems in flux partitioning

NEE is small in comparison to its component fluxes.

Partitioning of two components from one equation -> ill-posed problem.

Objective

Hence, it is required to parameterize the one term and then estimate the other one.

Real-time GHG analyzer


Picarro 2201-i CO₂-CH₄ Analyzer was used to get the following parameters:

[CO₂], [CH₄], $\delta^{13}\text{C}_{\text{CO}_2}$, $\delta^{13}\text{C}_{\text{CH}_4}$
(1 Hz frequency)


In association with the IRGA CO₂
data (10 Hz)

$\delta^{13}\text{C}_{\text{CO}_2}$ was calculated
(at 10 Hz frequency).





Photosynthetic uptakes reduces CO_2 and increase $\delta^{13}\text{C}_{\text{CO}_2}$



CH_4 and $\delta^{13}\text{C}_{\text{CH}_4}$
(10 min averaged)

Calculation of “ isoflux”

$$\text{isoflux} = \underbrace{\overline{\rho w'(\delta^{13}\text{C}_a C)'}}_{\text{(I)}} + \underbrace{\rho \frac{d(\delta^{13}\text{C}_a C)}{dt}}_{\text{(II)}} = \underbrace{\delta^{13}\text{C}_R F_R}_{\text{(III)}} + \underbrace{F_A(\delta^{13}\text{C}_a - \Delta)}_{\text{(IV)}}$$


Isoflux = eddy isoflux (term I) + storage isoflux (term II).

(term III) denotes total ^{13}C added to the atmosphere by respiration and

(term IV) presents total ^{13}C removed from the atmosphere by photosynthesis.


Δ denotes the mean carbon isotope discrimination by the canopy.

Identifying the source signature




Moving Miller and Tans plot.

Green and red solid circles denote source signature for CO_2 .
Blue circles present the source signature for CH_4 .




Hourly average NEE (green squares) and isoflux (red dots).



Photosynthesis and respiration flux calculated from NEE (blue dotted line), isoflux, $\delta^{13}\text{C}_A$ and $\delta^{13}\text{C}_R$.

Control on the isotopic values



solid dots = time series of VPD. Blue-dot represents = low VPD. Red-cross = $\delta^{13}C_R$ and magenta-triangles = $\delta^{13}CH_4$. Yellow squared box = daily rainfall.

Conclusions

A large change ($\sim 6 \text{ ‰}$) in mean $\delta_s^{CO_2}$ in subsequent nights was attributed to the change in VPD as a result of sudden rain during the dry season.

A large depletion ($\sim 16 \text{ ‰}$) in $\delta_s^{CH_4}$ on the first observation night was linked to aerobic methane production from the plant methoxyl groups.

The NEE components, F_A and F_R , were estimated using the Isotope-EC combination method proposed by Bowling et al. (2001).

Thank you