

## Above ground response of rainforest functional groups to experimental drought and subsequent rewetting

**Angelika Kübert**<sup>1</sup>, Kathrin Kühnhammer<sup>2</sup>, Ines Bamberger<sup>1</sup>, Erik Daber<sup>1</sup>, Jason De Leeuw<sup>3</sup>, Kinzey Bailey<sup>3</sup>, Jia Hu<sup>3</sup>, S. Nemiah Ladd<sup>1</sup>, Laura Meredith<sup>3</sup>, Joost van Haren<sup>3</sup>, Matthias Beyer<sup>2</sup>, Maren Dubbert<sup>1</sup>, and Christiane Werner<sup>1</sup>

•<sup>1</sup>University of Freiburg, Ecosystem Physiology, Freiburg, Germany (angelika.kuebert@cep.uni-freiburg.de)

•<sup>2</sup>Institute for Geoecology, Technische Universität Braunschweig, Langer Kamp 19c, 38106 Braunschweig, Germany

•<sup>3</sup>School of Natural Resources and the Environment, University of Arizona, Tucson, United States of America

# Experimental design

## Project Water, Atmosphere, and Life Dynamics in Biosphere 2 (WALD)

- At Biosphere 2: controlled tropical rainforest ecosystem
- Extended drought period of 10 weeks
- Subsequent rewetting in 2 steps:
  - 1) deep water pulse (from Dec 3), enriched in  $^2\text{H}$  ( $\sim 1000 \text{ ‰}$ )
  - 2) rain events (Dec 8, from Dec 18 on every 2-3 days), isotopic range: natural abundance



# Experimental setup

## Measurements

- Gas leaf chambers coupled to water & carbon laser spectroscope, measured alternately for 5 minutes
- Chambers continuously flushed, supplied air continuously monitored



## Studied species: canopy trees and understory species


- Canopy: *Clitoria fairchildiana* CF, *Ceiba pentandra* CP, *Hibiscus tiliaceus* HE, *Hura crepitans* HC, PA: *Pachira aquatic* PA
- Understory: *Hibiscus rosa-sinensis* HR, *Pterocarpus indicus* PI

# Above ground response

## Analysis


- 1) Transpiration  $E$  and net assimilation rate  $A$  (von Caemmerer & Farquhar 1981), leaf water use efficiency  $A/E$
- 2) Isotopic signature of transpiration by mass balance (Simonin et al. 2013, Dubbert et al. 2017)






# First results: above ground response

# Leaf water use during drought



Mean A / mean E. n=3-4.




Mean  $\pm$  1 se. n=3-4.

Species

- PA
- CF
- HR
- PI



# Recovery after drought: deep water uptake



# Outlook

## Species-specific water use strategies in response to drought and rewetting:


Study of above and below-ground water use before, during and after drought, i.e.:

plant physiological responses such as leaf water use efficiency, leaf water potential, root water uptake dynamics






# Acknowledgement



- Many thanks to the numerous helpers at Biosphere 2:  
Sydney Kerman, Juliana Gil Loaiza, Adrian Dahlmann, Kathiravan Meeran, Jianbei Huang, Gemma Purse, Johannes Ingrisich and many more.
- Funding:  
ERC-Consolidator Grant (EU) #647008  
Biosphere 2, University of Arizona





# References

---

- Dubbert, M., Kübert, A., & Werner, C. (2017). Impact of Leaf Traits on Temporal Dynamics of Transpired Oxygen Isotope Signatures and Its Impact on Atmospheric Vapor. *Frontiers in Plant Science*, 8. <https://doi.org/10.3389/fpls.2017.00005>
- Simonin, K. A., Roddy, A. B., Link, P., Apodaca, R., Tu, K. P., Hu, J., Dawson, T. E., & Barbour, M. M. (2013). Isotopic composition of transpiration and rates of change in leaf water isotopologue storage in response to environmental variables: Isotopic non-steady-state transpiration. *Plant, Cell & Environment*, 36(12), 2190–2206. <https://doi.org/10.1111/pce.12129>
- von Caemmerer, S., and Farquhar, G.D.(1981).Some relationships between the biochemistry of photosynthesis and the gas-exchange of leaves. *Planta* 153, 376–387.[doi:10.1007/BF00384257](https://doi.org/10.1007/BF00384257)