

# Reduced tree growth across the semiarid United States due to asymmetric responses to intensifying precipitation extremes

Matthew P. Dannenberg<sup>1</sup>, Erika K. Wise<sup>2</sup>, and William K. Smith<sup>3</sup>

<sup>1</sup>Dept. of Geographical and Sustainability Sciences, University of Iowa


<sup>2</sup>Dept. of Geography, University of North Carolina, Chapel Hill

<sup>3</sup>School of Natural Resources and the Environment, University of Arizona

Sunrise, Yosemite Valley  
Albert Bierstadt, ca. 1870



Precipitation variability has increased across most of the United States, especially the Southwest, over the past century



Change in precipitation variability (CV)

Dannenberg, Wise & Smith (2019), *Science Advances*


# question

---

How does forest growth respond to increasing  
**precipitation variability?**




# Conceptual models



## Symmetric

A linear response of growth to precipitation in which reduced growth during dry extremes is compensated by increased growth during wet extremes. No long-term change in mean growth as precipitation variability increases.


# Conceptual models



## Positive Asymmetric

A nonlinear response of growth to precipitation in which reduced growth during dry extremes is **more than offset** by increased growth during wet extremes. Long-term **increase** in mean growth as precipitation variability increases.


# Conceptual models




## Negative Asymmetric

A nonlinear response of growth to precipitation in which reduced growth during dry extremes is **not offset** by increased growth during wet extremes. Long-term **decrease** in mean growth as precipitation variability increases.


Conceptual models  
were tested across the  
U.S. using growth  
records from 1,314  
tree-ring sites and  
**PRISM** precipitation  
estimates.









78 species



78 species




# Methods




Growth responses from each tree-ring site were classified into one of four conceptual models based on the best fit from ordinary least squares regression

# Methods




Responses to extreme events were detected using an extreme value capture method corrected for chance agreement using Cohen's kappa ( $\kappa$ )


Cool season  
(Oct-Mar)



Not precipitation-limited   Symmetric   Positive asymmetry   Negative asymmetry




Widespread negative asymmetries in western U.S.  
(increasing precipitation yields diminishing returns for growth)






Low precipitation extreme




High precipitation extreme




Dry years are very likely  
to have very low growth



Low precipitation extreme



High precipitation extreme





But wet years are less likely  
to have very high growth




Mostly symmetric throughout U.S.

Except for bur oak


Not precipitation-limited Symmetric Positive asymmetry Negative asymmetry




Not precipitation-limited




Symmetric



Positive Asymmetry




Negative Asymmetry



Warm season  
(Apr-Sep)


what **long-term effect** does increasing precipitation variability have on tree growth?

Growth Response




Widespread negative asymmetries between growth and precipitation

Change in precipitation variability



Large increase in precipitation variability




Simulations with two precipitation scenarios  
(same mean, different variance)



# High-variability scenario:

Double the likelihood of **low** growth

No change in likelihood of **high** growth



# summary

---

1

Growth responds **asymmetrically** to precipitation variability (especially western conifers and bur oak)

2

Increased precipitation variability can drive **decreases** in forest growth

## ENVIRONMENTAL STUDIES

# Reduced tree growth in the semiarid United States due to asymmetric responses to intensifying precipitation extremes

Matthew P. Dannenberg<sup>1,2\*</sup>, Erika K. Wise<sup>3</sup>, William K. Smith<sup>2</sup>

Earth's hydroclimatic variability is increasing, with changes in the frequency of extreme events that may negatively affect forest ecosystems. We examined possible consequences of changing precipitation variability using tree rings in the conterminous United States. While many growth records showed either little evidence of precipitation limitation or linear relationships to precipitation, growth of some species (particularly those in semiarid regions) responded asymmetrically to precipitation such that tree growth reductions during dry years were greater than, and not compensated by, increases during wet years. The U.S. Southwest, in particular, showed a large increase in precipitation variability, coupled with asymmetric responses of growth to precipitation. Simulations suggested roughly a twofold increase in the probability of large negative growth anomalies across the Southwest resulting solely from 20th century increases in variability of cool-season precipitation. Models project continued increases in precipitation variability, portending future growth reductions across semiarid forests of the western United States.

<sup>1</sup>Department of Geographical and Sustainability Sciences, University of Iowa, Iowa City, IA, USA. <sup>2</sup>School of Natural Resources and the Environment, University of Arizona, Tucson, AZ, USA. <sup>3</sup>Department of Geography, University of North Carolina, Chapel Hill, NC, USA.

\*Corresponding author. Email: matthew-dannenberg@uiowa.edu

Dannenberg *et al.*, *Sci. Adv.* 2019;5:eaaw0667 2 October 2019

Copyright © 2019  
The Authors, some  
rights reserved;  
exclusive licensee  
American Association  
for the Advancement  
of Science. No claim to  
original U.S. Government  
Works. Distributed  
under a Creative  
Commons Attribution  
NonCommercial  
License 4.0 (CC BY-NC).

[link](#)