

Paleoceanography and Paleoclimatology



RESEARCH ARTICLE

10.1029/2019PA003820

Key Points:

- ^{230}Th normalization is a robust tool for calculating sedimentary mass fluxes
- ^{230}Th may be affected by hydrothermal and boundary scavenging in certain discrete regions
- Generally, ^{230}Th mass fluxes are preferable over age model-based mass accumulation rates

Supporting Information:

- Supporting Information S1
- Table S1

Correspondence to:

K. M. Costa,
kassandracosta@whoi.edu

Citation:

Costa, K. M., Hayes, C. T., Anderson, R. F., Pavia, F. J., Bausch, A., Deng, F., et al. (2020). ^{230}Th normalization: New insights on an essential tool for quantifying sedimentary fluxes in the ocean.

^{230}Th Normalization: New Insights on an Essential Tool for Quantifying Sedimentary Fluxes in the Modern and Quaternary Ocean


Kassandra M. Costa¹ , Christopher T. Hayes² , Robert F. Anderson^{3,4} , Frank J. Pavia^{3,4,5} , Alexandra Bausch^{3,4,6} , Feifei Deng⁷ , Jean-Claude Dutay⁸ , Walter Geibert⁹ , Christoph Heinze¹⁰ , Gideon Henderson⁷ , Claude Hillaire-Marcel¹¹ , Sharon Hoffmann¹² , Samuel L. Jaccard¹³ , Allison W. Jacobel^{14,15} , Stephanie S. Kienast¹⁶ , Lauren Kipp^{3,16} , Paul Lerner¹⁷ , Jörg Lippold¹⁸ , David Lund¹⁹ , Franco Marcantonio²⁰ , David McGee²¹ , Jerry F. McManus^{3,4} , Figen Mekik²² , Jennifer L. Middleton³ , Lise Missiaen²³ , Christelle Not²⁴ , Sylvain Pichat^{25,26} , Laura F. Robinson²⁷ , George H. Rowland²⁷ , Matthieu Roy-Barman⁸ , Alessandro Tagliabue²⁸ , Adi Torfstein^{29,30} , Gisela Winckler^{3,4} , and Yuxin Zhou^{3,4}

¹Department of Geology and Geophysics, Woods Hole Oceanographic Institution, Woods Hole, MA, USA, ²School of Ocean Science and Engineering, University of Southern Mississippi, Stennis Space Center, MS, USA, ³Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY, USA, ⁴Department of Earth and Environmental Sciences, Columbia University, New York, NY, USA, ⁵Now at Division of Geological and Planetary Science, California Institute of Technology, Pasadena, CA, USA, ⁶Now at Department of Earth System Science, Stanford University, Stanford, CA, USA, ⁷Department of Earth Sciences, University of Oxford, Oxford, UK, ⁸Université Paris-Saclay, CNRS, CEA, UVSQ, Laboratoire des sciences du climat et de l'environnement, IPSL, Laboratoire CEA, UVSQ, CNRS, Gif sur Yvette, France, ⁹Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany, ¹⁰Geophysical Institute and Bjerknes Centre for Climate Research, University of Bergen, Bergen, Norway, ¹¹GEOTOP, Université du Québec à Montréal, Québec, Canada, ¹²Department of Earth and Ocean Sciences, University of North Carolina Wilmington, Wilmington, NC, USA, ¹³Institute of Geological Sciences and Oeschger Center for Climate Change Research, University of Bern, Bern, Switzerland, ¹⁴Department of Earth, Environmental, and Planetary Sciences, Brown University, Providence, RI, USA,




**How does ^{230}Th normalization
work?**

Constant production in the water column:



Constant production in the water column:

Very soluble
 $\tau = 400,000\text{yr}$



$$\beta_{^{230}\text{Th}} = 2.562 \times 10^{-5} \text{ dpm/cm}^3\text{kyr}$$

High Sediment Flux




10 dpm

5 g sediment

$^{230}\text{Th}_{\text{xs}}$ concentration
2 dpm/g

Low Sediment Flux




10 dpm

1 g sediment

$^{230}\text{Th}_{\text{xs}}$ concentration
10 dpm/g

^{230}Th Global Database



- Working group in the GeoTRACES program
- Compilation of over 1000 sites!
- Can be combined with measurements of, e.g., CaCO_3 , Opal, Fe, etc
- Update, synthesis, and outlook for ^{230}Th normalization

^{230}Th assumption

- All the ^{230}Th produced in the water column is scavenged by particles and buried in the underlying sediment
 - Flux to Sediment = Production in Water Column
 - $F / P = 1$


1. Boundary Scavenging

2. Nepheloid Layers

3. Hydrothermal Activity


Boundary Scavenging

- High particle fluxes near margins rapidly/efficiently strip ^{230}Th out of the water column
- This can create a *concentration gradient* in ^{230}Th in the water column
- ^{230}Th could then follow the concentration gradient towards the boundary




Modelled F/P captures boundary scavenging

73% of the ocean has ^{230}Th flux within 30% of water column production



*Background : Composite of iLoveClim, HAMOCC, CESM, and NEMO-PISCES models
Dots : measurements in sediment traps*

Boundary scavenging really only an issue in the polar seas and along continental margins




Most of the ocean is doing just fine!!


Other considerations for ^{230}Th application

- Discussed next:
 - *Nepheloid layers*
 - *Hydrothermal scavenging*
- Also in the Paper:
 - *Grain sizes and sediment focusing*
 - *Calcium carbonate dissolution*

Nepheloid layers: “Boundary Scavenging” along the seafloor



Lerner et al., submitted




Gardner et al., 2018


- Sediment that gets resuspended due to turbulence along the seafloor

Nepheloid layers do scavenge extra ^{230}Th ...


- But the particle concentration is so high...
- That the net ^{230}Th concentration decreases !!
- Could possibly lead to *overestimating* the mass flux
- But need to test this with coretop samples under nepheloid layers



Hydrothermal Activity: Fe-Mn rich particles have high affinity for scavenging ^{230}Th




Pavia et al., 2018




Lund et al., 2019

- Burial rates on mid-ocean ridges may be higher than production ($F/P > 1$)

High ^{230}Th fluxes have to be supplied from somewhere...



Lund et al., 2019



Lund et al., 2019; Middleton et al., in prep

- Excess scavenging at the ridge creates a concentration gradient
- Lateral diffusion supplies more ^{230}Th
- Net effect of hydrothermal scavenging is *difficult to predict*

Summary

- Where it works: MOST of the ocean!!
- Where to use caution:
 - Continental margins and polar oceans, where boundary scavenging may be high
 - Fluxes biased too LOW
 - Regions with extensive nepheloid layers
 - Fluxes biased too HIGH
 - Mid-ocean ridges with active hydrothermal systems
 - Fluxes could be biased either HIGH or LOW
- Future directions:
 - Validate hypotheses with joint ^3He - ^{230}Th measurements

Questions?

- Please contact me at
 - kassandracosta@whoi.edu
- I look forward to hearing from you!