

A synchronous change of mid- to late-Holocene hydroclimate and prehistoric population in coastal East Asia indicated by pollen, XRF and grain size data

Jinheum Park¹, Jungjae Park^{1,2}, Sangheon Yi^{3,4}, Jin Cheul Kim³, Eunmi Lee³, Quihong Jin¹ and Jieun Choi¹



¹ Department of Geography, Seoul National University

² Institute for Korean Regional Studies, Seoul National University

³ Geology Division, Korea Institute of Geoscience and Mineral Resources

⁴ Department of Petroleum Resources Technology, University of Science and Technology, Korea


- Growing interest in a relationship between ancient cultures and climate change
 - Researches in many different regions around the world
 - Xie et al. (2013), Collins et al. (2017), Manning et al. (2017), Xu et al. (2019) etc.
- Nonetheless, a direct comparison between continuous and high-resolution climate proxies and archaeological data is in a starting stage in the Korean Peninsula
 - Constantine et al. (2019), Park et al. (2019) etc.
- Here, we analyzed pollen, XRF, and grain size from a core in the southern Korean Peninsula, and compared the data with summed probability distribution (SPD) of archaeological radiocarbon dates, which is an indicator for ancient population.

Acknowledgements

: This research was supported by the Basic Research Project (GP2017-013) titled “Investigation of subtropicalization in the Korean Peninsula using the geological Proxy: Climate-geoecosystem assessment of the mid-Holocene period” of the Korea Institute of Geoscience and Mineral Resources (KIGAM), funded by the Ministry of Science, ICT, and Future Planning. It was also supported by the Ministry of Education of the Republic of Korea and the National Research Foundation of Korea (NRF-2018S1A2A01025813).

1

Introduction





Adapted from Ryan et al. (2009) and Google Earth software

- Core STP18-03 (yellow star)
→ 35°26'18"N, 128°46'41"E
→ floodplain of Miryang River, Korea
- XRF (0.5cm interval)
Pollen (137 samples)
Grain size (5cm interval)
- Chronology
→ 8 radiocarbon & 8 OSL dates
→ 14 of 16 dates used (bacon)

2


Materials and methods



- (c) Brown: sand percentage (%)
- (d) Black: titanium content (area)
- (e) Green: tree percentage (%)
- (f) Magenta: *Artemisia* + *Poaceae* + ferns (%)

- The proxies are generally in phase
- Precipitation \uparrow , river discharge (sand) \uparrow , arboreal plant growth \uparrow , titanium erosion \downarrow
- Precipitation \downarrow , river discharge (sand) \downarrow , arboreal plant growth \downarrow , titanium erosion \uparrow

3 Results



- (a) Crimson: June insolation at 30°N (W/m²) (Berger et al., 1991)
- (b) Green: tree percentage (%)
- (c) Magenta: *Artemisia* (mugwort) + *Poaceae* (grass) + ferns (%)
- (d) Black: XRF titanium content (area)
- (e) Brown: sand percentage (%)
- (f) Red: SPD of archaeological records in Korea (Oh et al., 2017)
- (g) Blue: Western Tropical Pacific SST (°C) (Stott et al., 2004)

- Drier climate (b-e) are synchronous with lower ancient population (f)
 - ca. 4.8 - 3.6 ka & ca. 2.8 - 2.3 ka BP
 - Ancient societies in Korea were vulnerable to climate deterioration
- Active ENSO-like phase was a possible trigger for drier climate in the Korean Peninsula by dampening strength of East Asian Summer Monsoon (EASM) → vertical orange boxes

4

Discussion

- We conducted a multi-proxy analysis (XRF, pollen and grain size) on the hydroclimate change in the Korean Peninsula during ca. 8.3 - 2.3 ka BP.
- Periods of adverse climate conditions were synchronous with times when the density of archaeological records diminished.
 - Prehistoric cultures would have been vulnerable to climate deterioration during ca. 4.8 - 3.6 ka and ca. 2.8 - 2.3 ka BP.
 - Resource availability for hunter-gatherers and agricultural chance would have been confined due to precipitation deficits.
- Hydroclimate in the Korean Peninsula was likely modulated by the Pacific Ocean.
 - ENSO-like forcing dampened strength of the EASM

5 Conclusion

- Berger, A., & Loutre, M. F. 1991. Insolation values for the climate of the last 10 million years. *Quaternary Science Reviews*, 10(4), 297–317.
- Blaauw, M., & Christen, J. A. 2011. Flexible paleoclimate age–depth models using an autoregressive gamma process. *Bayesian analysis*, 6(3), 457–474.
- Bond, G., Showers, W., Cheseby, M., Lotti, R., Almasi, P., DeMenocal, P., ... & Bonani, G. 1997. A pervasive millennial-scale cycle in North Atlantic Holocene and glacial climates. *science*, 278(5341), 1257–1266.
- Chen, J., Chen, F., Feng, S., Huang, W., Liu, J., & Zhou, A. 2015. Hydroclimatic changes in China and surroundings during the Medieval Climate Anomaly and Little Ice Age: spatial patterns and possible mechanisms. *Quaternary Science Reviews*, 107, 98–111.
- Cheng, H., Fleitmann, D., Edwards, R.L., Wang, X., Cruz, F.W., Auler, A.S., Mangini, A., Wang, Y., Kong, X., Burns, S.J., 2009. Timing and structure of the 8.2 kyr BP event inferred from $\delta^{18}O$ records of stalagmites from China, Oman, and Brazil. *Geology* 37, 1007–1010.
- Collins, J. A., Prange, M., Caley, T., Gimeno, L., Beckmann, B., Mulitza, S., ... & Schefuß, E. 2017. Rapid termination of the African Humid Period triggered by northern high-latitude cooling. *Nature communications*, 8(1), 1372.
- Constantine, M., Kim, M., & Park, J. 2019. Mid-to late Holocene cooling events in the Korean Peninsula and their possible impact on ancient societies. *Quaternary Research*, 1–11.
- DeMenocal, P. B. 2001. Cultural responses to climate change during the late Holocene. *Science*, 292(5517), 667–673.
- Faegri, K., Iversen, J., 1989. *Textbook of pollen analysis* (4th edn by Faegri, K., Kaland, P.E. & Krzywinski, K.). Wiley, Chichester.
- Grimm, E.C., 1987. CONISS: a FORTRAN 77 program for stratigraphically constrained cluster analysis by the method of incremental sum of squares. *Computers & Geosciences* 13, 13–35.
- Grimm, E.C., 1991. *Tilia and Tiliagraph*. Illinois State Museum, Springfield, 101.
- Katsuki, K., Yang, D. Y., Lim, J., Lee, J. Y., Asahi, H., & Han, M. 2017. Multi-centennial-scale changes in East Asian typhoon frequency during the mid-Holocene. *Palaeogeography, palaeoclimatology, palaeoecology*, 476, 140–146.
- Lim, J., Lee, J. Y., Hong, S. S., Park, S., Lee, E., & Yi, S. 2019. Holocene coastal environmental change and ENSO-driven hydroclimatic variability in East Asia. *Quaternary Science Reviews*, 220, 75–86.
- Manning, J. G., Ludlow, F., Stine, A. R., Boos, W. R., Sigl, M., & Marlon, J. R. 2017. Volcanic suppression of Nile summer flooding triggers revolt and constrains interstate conflict in ancient Egypt. *Nature communications*, 8(1), 900.
- Moy, C. M., Seltzer, G. O., Rodbell, D. T., & Anderson, D. M. 2002. Variability of El Niño/Southern Oscillation activity at millennial timescales during the Holocene epoch. *Nature*, 420(6912), 162.
- Nahm, W.-H., Kim, J.K., Kim, J.-Y., Yi, S., Lim, J., Kim, J.C., 2013. The Holocene climatic optimum in Korea: Evidence from wetland records. *Palaeogeography, Palaeoclimatology, Palaeoecology* 376, 163–171.
- Oh, Y., Conte, M., Kang, S., Kim, J., & Hwang, J. 2017. Population fluctuation and the adoption of food production in prehistoric Korea: using radiocarbon dates as a proxy for population change. *Radiocarbon*, 59(6), 1761–1770.
- Park, J., Park, J., Yi, S., Kim, J.C., Lee, E., Jin, Q., 2018. The 8.2 ka cooling event in coastal East Asia: High-resolution pollen evidence from southwestern Korea. *Scientific reports* 8, 12423.
- Park, J., Park, J., Yi, S., Kim, J. C., Lee, E., & Choi, J. 2019. Abrupt Holocene climate shifts in coastal East Asia, including the 8.2 ka, 4.2 ka, and 2.8 ka BP events, and societal responses on the Korean peninsula. *Scientific reports*, 9(1), 10806.
- Park, J., Shin, Y.H., Byrne, R., 2016. Late-Holocene vegetation and climate change in Jeju Island, Korea and its implications for ENSO influences. *Quaternary Science Reviews* 153, 40–50.
- Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk Ramsey, C., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., 2013. IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* 55, 1869–1887.
- Ryan, W. B. F., S.M. Carbotte, J. Coplan, S. O'Hara, A. Melkonian, R. Arko, R.A. Weissel, V. Ferrini, A. Goodwillie, F. Nitsche, J. Bonczkowski, and R. Zemsky 2009, Global Multi-Resolution Topography (GMRT) synthesis data set, *Geochem. Geophys. Geosyst.*, 10, Q03014,
- Song, B., Li, Z., Lu, H., Mao, L., Saito, Y., Yi, S., ... & Zuo, X. 2017. Pollen record of the centennial climate changes during 9–7 cal ka BP in the Changjiang (Yangtze) River Delta plain, China. *Quaternary Research*, 87(2), 275–287.
- Stott, L., Cannariato, K., Thunell, R., Haug, G. H., Koutavas, A., & Lund, S. 2004. Decline of surface temperature and salinity in the western tropical Pacific Ocean in the Holocene epoch. *Nature*, 431(7004), 56.
- Xie, S., Evershed, R. P., Huang, X., Zhu, Z., Pancost, R. D., Meyers, P. A., ... & Gu, Y. 2013. Concordant monsoon-driven postglacial hydrological changes in peat and stalagmite records and their impacts on prehistoric cultures in central China. *Geology*, 41(8), 827–830.
- Wang, C., Lu, H., Zhang, J., Gu, Z., & He, K. (2014). Prehistoric demographic fluctuations in China inferred from radiocarbon data and their linkage with climate change over the past 50,000 years. *Quaternary Science Reviews*, 98, 45–59.
- Xu, D., Lu, H., Chu, G., Liu, L., Shen, C., Li, F., ... & Wu, N. 2019. Synchronous 500-year oscillations of monsoon climate and human activity in Northeast Asia. *Nature communications*, 10(1), 1–10.
- Zhou, X., Sun, L., Zhan, T., Huang, W., Zhou, X., Hao, Q., Wang, Y., He, X., Zhao, C., Zhang, J., 2016. Time-transgressive onset of the Holocene Optimum in the East Asian monsoon region. *Earth and Planetary Science Letters* 456, 39–46.