

# Assessment of decadal prediction skills and sensitivity to SIT initialization

Tian Tian<sup>1</sup>, S. Yang<sup>1</sup>, P. Karami<sup>2</sup>, F. Massonnet<sup>3</sup>, T. Kruschke<sup>2</sup> and T. Koenigk<sup>2,4</sup>  
<sup>1</sup>DMI, <sup>2</sup>SMHI, <sup>3</sup>UCLouvain, <sup>4</sup>SU

## Summary (wrt 1997-2016)


**Bias:** >2m ice thickness (SIT) along the Sept ice edge and the Arctic coastlines in **FREE (un-init)**, but multi-year ice (MYI) <50% Arctic sea-ice cover in OBS.

**SS<sub>A12</sub>:** thick MYI SIT skill improved greatly by **A12 (all-init.)** in winter and, in turn constraining Sept min.

**SS<sub>A10</sub>:** thinner first-year ice (FYI) SIC skill mostly benefit from Atlantic heat transport by **A10 (no-ice-init)**.


**AS:** skill increased in predicting Sept. min in FYI/MYI regions, and re-emerged in yr 5-7 due to advection of sea-ice anomalies

### Sept bias ice edge & SIT




### Skill score (SS)

$$SS_{A1x} = 1 - \frac{RMSE_{A1x}}{RMSE_{FREE}}$$



### Added skill (AS)

$$AS = SS_{A12} - SS_{A10}$$



Novelty@DMI	Introducing SIV anomaly to constrain SIT initialization with 5 categories
<b>Model system</b>	EC-Earth3 Climate Prediction System with Anomaly Initialization ( <b>EC-Earth3-CPSAI</b> )
<b>Configuration</b>	Atm and Ocn coupled <b>GCM</b> (atm:T255L91 + ocn: ORCA1L75)
<b>Forcing</b>	<b>CMIP6</b> historical + future scenario SSP2-4.5
<b>Anomaly init. to ocn + ice (NEMO3.6+LIM3)</b>	5-member <b>ORAS5</b> (1979-2019) & single backward extension (1960-1978); anomaly wrt obs climate (1979-2014); initialized variables: 3D ocean (temperature & salinity) and 2D sea-ice (SIC, SIT and snow thickness)
<b>Full-field init. to atm (IFS cycle 36r4)</b>	ERA40 (1960-1978), <b>ERA1</b> (1979-2018), ERA5 (2019) + 3D-T perturbation with random differences (to the order of $10^{-5}$ K)
<b>Init. hindcasts</b>	yearly started on <b>1 November</b> over <b>1960-2019</b> , 10 yrs + 2 months long
<b>Model climate</b>	EC-Earth3 CMIP6 historical run <b>r5i1p1f1</b> (1979-2014)
<b>Joint ensembles</b>	<b>15</b> members labelled <b>r1-15i2p1f1</b> : DMI performing 10 members + SMHI performing 5 members
<b>Sensitivity tests</b>	5-member ens-mean comparison: <b>A12</b> (all-init.), <b>A11</b> (no-SIT-init.), <b>A10</b> (no-ice-init.), <b>FREE</b> (no-init);

## TT and SY supported by four projects:

- ARCPATH – the NordForsk research programme Arctic Climate Predictions: Pathways to Resilient, Sustainable Societies
- NCKF – the Danish National Center for Climate Research
- EUCP – European Climate Prediction System



The Blue-Action project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 727852