



UNIVERSITÄT **BONN**

ETH zürich



Refining chronologies by dating pollen concentrates – new approach of separating pollen using flow cytometry


Christoph Steinhoff¹, Nadine Pickarski¹, Irka Hajdas², Thomas Litt¹

¹Institute of Geosciences, University of Bonn


²Ion Beam Physics, ETH Zürich

06.05.2020

PALEOVAN - Project at Lake Van




- ICDP funded deep drilling project at Lake Van in Eastern Anatolia (2004 and 2010)
- Analysis and reconstruction of the Quaternary climate in the Near East
- Drilling at Ahlat Ridge carried out sedimentary succession of ca. **220 m**, spanning over **600 000 years**



- North-south trending **seismic profile** crossing main **Tatvan Basin** and **Ahlat Ridge** (PALEOVAN location - **375 mbsf**)

Lake Van - Varvechronology

- Interglacial layers are **annually laminated**
- Varve counting provides possibility to create high resolution chronology
- Maximum age for the Holocene-Pleistocene-Transition is **8719 vy BP (yellow)**
- **Discrepancy** of ca. 3000 years to GSSP
- Seven volcanic ash layers (**gray**) allow the correlation to other sedimentary profiles in Lake Van
- Comparison to existing varve chronology by *Landmann et al. (1996)* (**orange**) shows missing varves/erroneous counting in lower part
- ^{14}C measurements above and below reference layers (tephra) to validate varve chronology



From Sediment to Pollen Concentrate

Chemical Pretreatment

Initial volume of **15-45 cm³**, depending on pollen conc.

Use of **HCl**; removal of carbonates

Coarse sieving (**53 µm** mesh width)

Use of **HF**; dissolution of silicates


Ultrasonic sieving (**10 µm**)

Flow Cytometry

Cytometric analysis using **flow cytometry** (*GFZ Potsdam*)

Laser based **multiparameter analysis**

Separation based on fluorescence activated cell sorting (**FACS**)




AMS Dating

AMS dating using gas ion source (*ETH Zürich*)

Flow Cytometry



Application to **analyze, identify** and **isolate** populations of heterogeneous samples according to optical properties



1. Measuring intensity of refracted light on surface structure (**forward scatter** FSC and **side scatter** SSC)
 - Indicator for size, shape and internal complexity of grains
2. Additionally measuring spectra and intensity of **auto fluorescence signals** of particles generated by various lasers


Scatter plots (each dot represents 1 particle) for identification of subpopulations (**cytometric analysis**)

3. Morphological properties: No differentiation between **sporomorphs** (**pink, blue**) and **detritus** (**gray**) possible
4. Fluorescence signal: Separation of distinctive **pollen** (**pink**) and **spores** (**blue**) is now possible



Results – Flow Cytometry


- Cytometric analysis allows **unequivocal identification** of pollen and spores
- Total pollen count varies widely (**40 000 – 900 000 grains**) and is independent of initial volume, sample depth, and sorting strategy
- Ratio of sporomorphs in pollen concentrate is largely **>70%**
 - Some detrital particles with inherent color remain in concentrate after separation
 - But recently developed strategy allows isolation of pollen concentrates with purity **>90%**
- Sample weight (µg C) varies between **13 – 160 µg carbon** and is **suitable for AMS dating**



© Authors. All rights reserved

Results – AMS Dating

- ^{14}C ages (**green**) **older** than varvechronology
- Good agreement in lower part of section
 - **Gap** between varve counting and GSSP for onset of the Holocene can be **closed**
- **Hiatus** in sedimentation above tephra 1
 - **Landslides** due to **ridge position**
- **Statistical errors** increase from ± 100 years in upper part to ± 170 years in lower part
- **Scattering** of all **measurements** varies between 2500 years around tephra 1-3 and 700 years for tephra 5 and 7
- Large uncertainty below **tephra 3**, wide range in pollen ages



© Authors. All rights reserved

Outlook

It is possible to **generate ^{14}C ages** derived from **pollen concentrates**!

Discrepancies concerning the Holocene chronology of Lake Van **can be solved**.

Still, we need **improvements** in sample preparation to **increase** pollen **yield**, while **minimizing** risk of **contamination**.

Reliable measurements on **old** and **recent carbon** are necessary to exclude the contamination of the flow cytometry itself.

Further **samples between** the **tephra** layers are already taken, which will help do **reduce** the **scattering** of our AMS ages and further **refine** the **Holocene chronology of Lake Van**

Thank you for your interest!

Special Thanks to:

Team of the Flow Cytometry Core Facility, *University of Bonn*

Dr. Jens Kallmeyer and Axel Kitte (both *GFZ Potsdam*) for assistance during cytometric analysis

We thank Dr. Caroline Welte for her help with the GIS 251 AMS analysis as well as the team at the Laboratory of Ion Beam Physics, *ETH Zürich*

ukb universitäts
klinikum bonn

GFZ
Helmholtz-Zentrum
POTSDAM

ETH zürich

DFG