

# Rapid, continuous radiocarbon analysis of carbonate archives using laser ablation

M. Wertnik<sup>1,2</sup>, C. Welte<sup>1,2</sup>, L. Wacker<sup>1</sup>, C. Yeman<sup>1</sup>, B. Hattendorf<sup>3</sup>, J. Koch<sup>3</sup>, M. Christl<sup>1</sup>,  
J. Fohlmeister<sup>4</sup>, D. Riechelmann<sup>5</sup>, H.-A. Synal<sup>1</sup>, T. Eglinton<sup>2</sup>

1 Laboratory of Ion Beam Physics, ETHZ, Otto-Stern Weg 5, HPK, 8093 Zurich, Switzerland

2 Geological Institute, ETHZ, Sonneggstrasse 5, 8092 Zurich, Switzerland

3 Laboratory of Inorganic Chemistry, D-CHAB, ETHZ, Vladimir-Prelog Weg 1, 8093 Zurich, Switzerland


4 Potsdam Institute for Climate Impact Research (PIK) e.V., 14473 Potsdam, Germany

5 Institut für Geowissenschaften, Johannes Gutenberg-Universität Mainz, Johann-Joachim-Becher-Weg 21, 55099 Mainz

wertnikm@phys.ethz.ch, cwelte@phys.ethz.ch, wacker@phys.ethz.ch, yemanc@phys.ethz.ch,  
bodo@inorg.chem.ethz.ch, koch@inorg.chem.ethz.ch, mchristl@phys.ethz.ch,  
jens.fohlmeister@pik-potsdam.de, d.riechelmann@geo.uni-mainz.de, synal@phys.ethz.ch,  
timothy.eglinton@erdw.ethz.ch



# Laser-Ablation-AMS



	Original Setup	Modified
fluence on sample	1-2.5 J·cm <sup>-2</sup>	8-23 J·cm <sup>-2</sup>
cell volume	600 µL	900 µL
spot size	680 x 110 µm <sup>2</sup>	140 x 75 µm <sup>2</sup>
ablation rate	100 µg/min	50-100 µg/min
carbon flow	6 µg/min	3.5-7 µg/min
max. ion current	7 µA	9 µA
blank level	0.011±0.002 up to 36'000 y	0.009 ± 0.002 up to 38'000 y

pulsed ArF excimer laser (GAM LASER, USA)  
 laser repetition rate: up to 250 Hz  
 (Yeman, C. (2019))

Carbonate conversion efficiency: 70 ± 4%  
 (Welte et. al. (2017))




# Stalagmite STAM-4


STAM-4 has been chosen as a sample because it is a fast grown stalagmite. Could a continuous  $^{14}\text{C}$  record be used to detect a subannual signal?

## Previous measurements:

- Uranium/Thorium: inconclusive (too young, did not contain enough Thorium)
- $^{14}\text{C}$  graphite-based measurements have been taken and the results agree well with those from LA-AMS



## Growth Rate



## Cloșani Cave

- 45.1°N, 22.8°E southern slope of Carpathians in SW Romania
- 433 m above sea level (msl)
- In Upper Jurassic limestone mainly calcite (93%), dolomite (7%)
- 1458 m long over a vertical range of 15 m (Warken et. al. (2018))



# Radiocarbon Record of Stalagmite STAM-4

## velocities:

young  $\rightarrow$  old:

$v = 5 \mu\text{m/s}$

(equals  $\sim 19$   
pts/mm)


old  $\rightarrow$  young:

$v = 2 \mu\text{m/s}$

(equals  $\sim 48$   
pts/mm)

data points are  
correlated to sample  
location with a  
python script.

Sample measuring  
time: about 6 h  
(+ 3 h standards)




# Exploratory Analysis of STAM-4

**Savitzky-Golay** filter with polynomials of degree 3 and window width of 33 for scan with 5  $\mu\text{m/s}$  and 67 for scan with 2  $\mu\text{m/s}$ .

Preliminary layer width measurement using high resolution scan picture.


Further analysis required to see what kind of signal we can detect in STAM-4.

While a subannual signal seems unlikely in this case, interesting features are found e.g. around 1980.





# Zoom-in of Savitzky-Golay filter



This work was supported by ETH Research Grant ETH-03 18-2.