


Fast determination of surface water mass changes using regional orthogonal functions

Guillaume Ramillien (1, 3, 4), Lucia Seoane (2, 3, 4) and José Darrozes (2, 3, 4)

Guillaume.Ramillien@get.omp.eu; Lucia.Seoane@get.omp.eu

1. Centre National de la Recherche Scientifique (CNRS)
2. Université Paul Sabatier (UPS)
3. Géosciences Environnement Toulouse (GET)
4. Observatoire Midi-Pyrénées (OMP)

Spatial scales of GRACE signal representation (*)




We propose a representation method that maximize the information inside a domain

by construction of a finite set of regional orthogonal functions known as *Slepian* functions (Simons 2009)




(*) modified from Freedden & Schreiner 2008

Examples of regional *Slepian* functions : cases of Africa & Congo basin



Recovery of WaterGap Global Hydrology Model (WGHM) maps by Kalman Filtering estimation


$$s(\theta, \lambda) = \sum_{K=1}^{K_{max}} slepian(\theta, \lambda)^{(k)} a^{(k)}$$



Inversion of real GRACE data – First results (1/2)


Recovery of the first modes from GRACE-based geopotential differences

First modes versus time by KF estimation



Inversion of real GRACE data – First results (2/2)

Recovery of the first modes from GRACE-based geopotential differences



Highlights

A promising approach based on orthogonal *Slepian* functions has been successfully developed to invert GRACE satellite geopotential difference data and map continental water storage changes.

It is possible to follow the evolution of the GRACE *Slepian* coefficients adjusted by Kalman Filtering (KF).

Perspectives

Numerical optimization of the computation of *Slepian* coefficients of higher orders ($K \sim 7000$) would permit to reach better spatial resolutions in continental hydrology.

Multi-year series of coefficients estimated from real GRACE (-FO) data would enable spatio-temporal analysis and characterization of water mass processes.