

A diachronous opening of the Iapetus Ocean in the Neoproterozoic

Boris Robert¹, Mathew Domeier¹, Johannes Jakob¹


¹Centre for Earth Evolution and Dynamics, University of Oslo, Norway



UiO : **University of Oslo**


The Iapetus Ocean

Opening
~700-550 Ma




Merdith et al., 2017

500 Ma



Domeier, 2016

Closure
~430 Ma



When does the Iapetus Ocean open?

Three distinct times proposed for the opening between Laurentia and Amazonia :

- (1) 550-530 Ma (e.g. Pisarevsky et al. 2008)
- (2) 615-570 Ma (e.g. Cawood et al. 2001)
- (3) 750-680 Ma (e.g. Li et al. 2013)

In this study, **review of the geological observations** from 750 to 520 Ma to test these scenarios.

Distribution of magmatism linked to rifting




Two main phases of rifting at 750-680 Ma and 615-550 Ma.

Phase (II) usually interpreted as leading to breakup.

Phase (I) only found between Laurentia and Amazonia and Baltica.


Which one lead to breakup?




Robert et al. 2020
Geology, in press.

Rift-related magmatic pulses at 750–680 Ma (purple) and 615–550 Ma (red) in Laurentia, Amazonia and Baltica. Esp: Eastern Sierra Pampeanas, Wsp: Western Sierra Pampeanas, AA: Antofalla-Arequipa, RP: Rio de la Plata, Dr: Dashwoods ribbon, Y: Yucatan, S: Sierra Madre, M: Mixteca Oaxaca.

Detrital zircon record



Robert et al. 2020, *Geology*, in press.



Almost no Neoproterozoic rock exposures along west Amazonia.

An enigmatic 650-500 Ma detrital zircon age peak in younger sediments along west Amazonia.

Three possible origins:


- Magmatism related to Iapetus rifting
- 0.9-0.48 Ma magmatism from the Brasiliano orogenic system
- A hidden magmatic arc along western Amazonia (Chew et al. 2008)

A. Histograms and kernel density estimates of detrital zircon U-Pb ages of post-500 Ma metasedimentary rocks from western Amazonia, early Cambrian and Ediacaran synrift sediments along eastern Laurentia. B & C: Two alternative scenarios that may explain the provenance of the 650–500 Ma zircons along West Amazonia. Sampling sites: 1: Marañon Complex, 2: Cordillera Real of Ecuador, 3: Humber zone in Newfoundland, 4: Blue Ridge inlier, 5: Birmingham basement graben system.

There is no direct observation of this arc but this scenario better explains amplitude and temporal restriction of the peak, and detrital zircon record in Laurentia.

Our favorite scenario: opening of 2 oceans

- Opening at ~700 Ma of Paleo-lapetus, opening ~600 Ma of Neo-lapetus, explain:
 - the 2 pulses of rift-related magmatism
 - the presence of a subduction zone along western Amazonia by 650 Ma.
- Paleomagnetic data at 615 Ma are in agreement with an early opening between Amazonia and Laurentia.
- Between 750 and 615 Ma, paleomagnetic data are sparse, which preclude any paleomagnetic test.



Robert et al. 2020
Geology, in press.

Paleomagnetic poles:


LR: Long Range dyke (Murthy et al. 1992; Hodych et al. 2004)

ED: Egersund dykes (Walderhaug et al. 2007)

AD: Adma Diorite (Morel, 1981)

PS: Planalto da Serra complex (Garcia et al. 2013).

lapetus, an analog to Tethys?



- A possible mechanism:
 - Subduction of the Paleo-lapetus oceanic ridge puts under tension the Laurentia plate, which could have detached continental slivers from Laurentia.
 - A similar mechanism is invoked for the opening of Neo-Tethys (Stampfli and Borel, 2002).
 - The emplacement of the Central Igneous Magmatic Province (C) may have facilitated the detachment of a thin continental ribbon, similar to the Panjal traps (P) in Tethys.

References



Merdith, A.S. et al., 2017, A full-plate global reconstruction of the Neoproterozoic: Gondwana Research, v. 50, p. 84–134, doi:10.1016/j.gr.2017.04.001.

Domeier, M. (2016). A plate tectonic scenario for the Iapetus and Rheic oceans. *Gondwana Research*, 36, 275–295.

Pisarevsky, S. A., Murphy, J. B., Cawood, P. A., & Collins, A. S. (2008). Late Neoproterozoic and Early Cambrian palaeogeography: models and problems. *Geological Society, London, Special Publications*, 294(1), 9–31.

Cawood, P.A., McCausland, P.J.A., and Dunning, G.R., 2001, Opening Iapetus: Constraints from the Laurentian margin in Newfoundland: *Geological Society of America Bulletin*, v. 113, p. 443–453, doi:10.1130/0016-7606(2001)113<0443:OICFTL>2.0.CO;2.

Li, Z. X., Evans, D. A., & Halverson, G. P. (2013). Neoproterozoic glaciations in a revised global palaeogeography from the breakup of Rodinia to the assembly of Gondwanaland. *Sedimentary Geology*, 294, 219–232.

Chew, D., Magna, T., Kirkland, C., Miskovic, A., Cardona, A., Spikings, R., and Schaltegger, U., 2008, Detrital zircon fingerprint of the Proto-Andes: Evidence for a Neoproterozoic active margin? *Precambrian Research*, v. 167, p. 186–200, doi:10.1016/j.precamres.2008.08.002.

Eriksson, K.A., Campbell, I.H., Michael Palin, J., Allen, C.M., and Bock, B., 2004, Evidence for Multiple Recycling in Neoproterozoic through Pennsylvanian Sedimentary Rocks of the Central Appalachian Basin: *The Journal of Geology*, v. 112, p. 261–276, doi:10.1086/382758.

Thomas, W.A., Astini, R.A., Mueller, P.A., Gehrels, G.E., and Wooden, J.L., 2004, Transfer of the Argentine Precordillera terrane from Laurentia: Constraints from detrital-zircon geochronology: *Geology*, v. 32, p. 965, doi:10.1130/g20727.1.

Allen, J. S., (2009). Paleogeographic reconstruction of the St. Lawrence promontory, Western Newfoundland [Ph.D. thesis]: Lexington, University of Kentucky, 310 p.

Chakraborty, S., Moecher, D.P., and Samson, S.D., 2012, Provenance of the Lower Ocoee Supergroup, eastern Great Smoky Mountains: *Geological Society of America Bulletin*, v. 124, p. 1278–1292, doi:10.1130/b30578.1.

Murthy, G., Gower, C., Tubrett, M., and Pätzold, R., 1992, Paleomagnetism of Eocambrian Long Range dykes and Double Mer Formation from Labrador, Canada: *Canadian Journal of Earth Sciences*, v. 29, p. 1224–1234, doi:10.1139/e92-098.

Hodych, J.P., Cox, R.A., and Košler, J., 2004, An equatorial Laurentia at 550 Ma confirmed by Grenvillian inherited zircons dated by LAM ICP-MS in the Skinner Cove volcanics of western Newfoundland: implications for inertial interchange true polar wander: *Precambrian Research*, v. 129, p. 93–113, doi:10.1016/j.precamres.2003.10.012.

Walderhaug, H.J., Torsvik, T.H., and Halvorsen, E., 2007, The Egersund dykes (SW Norway): a robust Early Ediacaran (Vendian) palaeomagnetic pole from Baltica: *Geophysical Journal International*, v. 168, p. 935–948, doi:10.1111/j.1365-246X.2006.03265.X.

Morel, P., 1981, Palaeomagnetism of a Pan-African diorite: a Late Precambrian pole for western Africa: *Geophysical Journal International*, v. 65, p. 493–503, doi:10.1111/j.1365-246X.1981.tb02724.X.

Garcia, M.S.R., Trindade, R.I.F., D’Agrella-Filho, M.S., Pinho, F.E.C., 2013, Paleomagnetismo do complexo alcalino Planalto da Serra (Mato Grosso): implicações para a formação do Gondwana: *Latinmag*, v. 3, Special Issue OB19, p. 1–8.

Stampfli, G.M., and Borel, G.D., 2002, A plate tectonic model for the Paleozoic and Mesozoic constrained by dynamic plate boundaries and restored synthetic oceanic isochrons: *Earth and Planetary Science Letters*, v. 196, p. 17–33, doi:10.1016/S0012-821X(01)00588-X.