

# Spatial assessment of inland excess water hazard using combined machine learning and geostatistical methods

Annamária LABORCZI <sup>1</sup>, Csaba BOZÁN <sup>2</sup>, Gábor SZATMÁRI <sup>1</sup>,  
János KÖRÖSPARTI <sup>2</sup>, László PÁSZTOR <sup>1</sup>



<sup>1</sup> *Institute for Soil Sciences and Agricultural Chemistry,  
Centre for Agricultural Research,  
Herman Ottó út 15, H-1022 Budapest, Hungary*



<sup>2</sup> *Research Institute of Irrigation and Water Management,  
National Agricultural Research and Innovation Centre,  
Anna-liget str. 35, H-5540 Szarvas, Hungary*



## What is Inland Excess Water (IEW)?


- Temporary water inundation, a form of surplus surface water.
- In flatlands: due to both precipitation and groundwater emerging on the surface.
- Most frequently in local depressions of large flat areas, irrespective of river floods.
- Literature frequently uses the phrase ‘waterlogging’.


Complex interaction of contributing factors:

- natural (e.g. meteorological, hydrogeological, pedological, topographical),
- anthropogenic (e.g. land use, agricultural engineering).

IEW causes several social, economic, and environmental problems in the flatland regions of Hungary, covering nearly half of the country.







## Study area


'10.07. Kisújszállás Excess  
Water Protection Section'

Overview map  
with relief of  
Hungary

## Reference Data

Map of temporally aggregated legacy information  
on inland excess water inundation  
frequency.

The values do not show real frequencies, they can  
be considered as an indicator (Number of  
inundation events occurred within the period  
from where observations are available).



Source: Laborczy et al. 2020.

<https://doi.org/10.3390/ijgi9040268>

# Environmental Co-Variables

## Soil

Digital Kreybig Soil Information System

- soil physical property layer,
- ‘landscape management soil type’

3D Soil Hydraulic Database of Europe

EU-SoilHydroGrids ver1.0

## Climate

- Average annual
- precipitation,
  - temperature,
  - evaporation,
  - evapotranspiration

Humidity index

## Land use

National CORINE Land Cover database

## Topography

Hungarian HydroDEM  
Digital Elevation Model  
and its derivatives


## Hydrogeology

Depth and the thickness of the  
uppermost aquitard

Standard depth of groundwater

Groundwater recharge and discharge  
areas

**Locations of the  
validation dataset**










# IEW inundation probability result maps

two methods:  
RK – Regression  
Kriging,  
RFK – Random  
Forest combined  
with Ordinary  
Kriging

two co-variable  
packages:  
BS – basic set,  
ES – extended set




# Validation



Category-difference between the independent validation dataset,  
and the categorized predicted values

# Validation



Difference between the legacy map's inundation frequency and the categorized inundation result maps.



# Conclusions

Comparing the results of the two approaches (RK and RFK), we did not find significant differences in their accuracy.

Both methods are appropriate for predicting inland excess water hazard, we suggest the usage of RFK, since

- it is more suitable for revealing non-linear and more complex relations than RK,
- it requires less presupposition on and preprocessing of the applied data,
- and keeps the range of the reference data, while RK tends more heavily to smooth the estimations, while
- it provides a variable rank, providing explicit information on the importance of the used predictors.

# Thank you for viewing our presentation!

If you are interested in more details, you can download  
our Open Access article:



International Journal of  
*Geo-Information*

Laborczi, A., Bozán, Cs., Körösparti, J., Szatmári, G., Kajári, B., Túri, N., Kerezsi, Gy., Pásztor, L. 2020. Application of hybrid prediction methods in spatial assessment of inland excess water hazard. *ISPRS International Journal of Geo-Information* 9: 268  
<https://doi.org/10.3390/ijgi9040268>

*Results & figures of the presentation comes from the above mentioned publication.*

## *Acknowledgements:*

*Our work has been supported by the National Research, Development and Innovation Office (NKFIH; Grant Nos. KH-126725 and K-131820), the INTERREG CE968: "RAINMAN" project, and the Higher Education Institutional Excellence Programme (NKFIH-1150-6/2019) of the Ministry of Innovation and Technology in Hungary, within the framework of the 4th thematic programme of the University of Debrecen.*

*Gábor Szatmári is supported by the Premium Postdoctoral Scholarship of the Hungarian Academy of Sciences (PREMIUM-2019-390).*

*Authors thank J. Matus for her indispensable contribution.*

