

# Combined use of Raman lidar and DIAL measurements and MESO-NH model simulations for the characterization of complex water vapour field structures and their genesis:

Water vapour mixing ratio:  $\Delta T=5$  min;  $\Delta z=30$  m

Paolo Di Girolamo<sup>1</sup>, Marie-Noelle Bouin<sup>2</sup>, Cyrille Flamant<sup>3</sup>, Donato Summa<sup>4</sup>, and Benedetto De Rosa<sup>1</sup>


<sup>1</sup> Scuola di Ingegneria, Università degli Studi della Basilicata, 85100 Potenza, Italy, digirolamo@unibas.it

<sup>2</sup> Météo-France /CNRM, 29280 Plouzané, France

<sup>3</sup> LATMOS/IPSL, UPMC Univ. Paris 06 Sorbonne Universités, UVSQ, CNRS, Paris, France

<sup>4</sup> IMAA-CNR, 85050 Tito Scalco (Pz), Italy


**IOP 8 - BASIL, Candillargues, 43°36'40.10"N ; 4° 4'15.80"E, 28-29 September 2012**



Measurements carried out by BASIL on 28 September 2012 reveal a **water vapour field** characterized by a **quite complex vertical structure**.

**Goal:** Assess the **origin** and **transport path** of the **different humidity filaments** observed by BASIL on this day **based on the comparison** with data from **MESO-NH model**.






Air masses within the lower of the two upper layers are found to overpass Southern Spain and Morocco, descending from an elevation of 2-3.5 km, while air masses within the uppermost layer (up to 4 km) are not tracked by MESO-NH.



MESO-NH back-trajectory analysis reveals that:

- air masses within the surface humidity layer originated over the Atlantic Ocean,
- air masses within the elevated filamentary humidity layer also originated over the Atlantic Ocean, overpassing the sea stretch North of Spain and Southern France at an altitude of ~1 km.



## Back-trajectory analysis from **HYSPLIT** extended further back in time (10 days)

**Back-traject. analysis** from **HYSPLIT** ending in Candillargues at the altitudes of the observed humidity layers : 18:00 UTC on 28 Sept. 2012 and starting **240 hours (10 days) earlier (18 Sept. 2012)**.



The back-trajectory extended back in time by **240 hours (10 days)** reveals that **air masses** in the:

- the **surface layer** and the **filamentary elevated structure** originated over the **North Atlantic**;
- the lower of the two upper layers originated in **tropical Atlantic area**;
- **uppermost humid layer**, overpassed Algeria, originated over **Central Africa (Mali)**.