



Research context

This work was conducted in the context of the **DFG-funded Collaborative Research Centre 1211 (CRC1211)** „Earth - Evolution at the dry limit“, particularly the subproject C03. The CRC pioneers the research on the **mutual evolutionary relationships between earth surface processes and biota in arid and hyper-arid areas** of the **Atacama Desert**, Northern Chile. Subproject C03 aims on understanding driving factors of **hillslope dynamics** in this driest part of the earth on recent and geological timescales.



Introduction

The Atacama-specific zebra stripes generally occur on hillslopes of varying inclination within areas of lowest rainfall and low fog frequency. As perhaps the most enigmatic landform in the Atacama, zebra stripes are made up by sequences of **contour-parallel**, lateral bands of **angular clasts** with **stripe-internal grain sorting** followed by rather **clast-free interspaces**. The lateral succession of 1-3m wide frontal lobes resembles **granular fingering** identified in experimental tests on granular flows of different particle mixtures (i.e. Pouliquen et al., 1997; Valderrama et al., 2018). In a region, where water, vegetation or soil formation processes are virtually absent, and their influence on surface geomorphology is limited, we emphasize the role of seismicity in the formation of zebra stripes and the Atacama landscape in general. Inter-site comparison points to considerable differences between individual zebra stripe appearances, potentially reflecting different timescales of seismic activity.




Acknowledgements

Funding by Deutsche Forschungsgemeinschaft (CRC 1211) is gratefully acknowledged. Logistical support during field campaigns in 2018 and 2019 was kindly provided by Eduardo Campos Sepulveda, Departamento de Ciencias Geológicas, Universidad Católica del Norte (UCN), Chile. Field work was assisted by Marie Gröbner, Lennart Meine and Florian Steininger.


References

Gastineau, R., Conway, S. J., Johnsson, A., Eichel, J., Mangold, N., Grindrod, P. M., & Izquierdo, T. (2019). Small-scale lobate hillslope features on Mars: A comparative 3D morphological study with terrestrial solifluction lobes and zebra stripe lobes. *Icarus*, 113606.
May, S. M., Hoffmeister, D., Wolf, D., & Bubenzer, O. (2019). Zebra stripes in the Atacama Desert revisited—Granular fingering as a mechanism for zebra stripe formation ?. *Geomorphology*, 344, 46–59.
Owen, J. J., Dietrich, W. E., Nishizumi, K., Chong, G., & Amundson, R. (2013). Zebra stripes in the Atacama Desert: Fossil evidence of overland flow. *Geomorphology*, 182, 157–172.
Pouliquen, O., Delour, J., Savage, S. B., 1997. Fingering in granular flows. *Nature* 386, 816–817.
Valderrama, P., Roche, O., Samaniego, P., van Wyk des Vries, B., Araujo, G., 2018. Granular fingering as a mechanism for ridge formation in debris avalanche deposits: Laboratory experiment and implications for Tutupaca volcano, Peru. *J. Volcanol. Geoth. Res.* 34, 409–418.


Geomorphological features indicating „activity“



Active forms of zebra stripes are indicated by **well-developed frontal lobes**, **reduced dust accumulation** in clast interstices, clearly developed **downslope sorting within individual stripes** and concurrent **slightly decreasing mean sorting values** of entire stripes towards lower slope sections. High proportions of surface cover are assumed to be related to free floating of particles. In contrast, presumed palaeo-activity of stripes is related to poorly visible to degraded frontal lobes, higher dust contents, indistinct sorting trends and high portions of clasts embedded into the underlying vesicular horizon.




Cosmogenic nuclide analysis (¹⁰Be; ²¹Ne)



Preliminary chronological data of cosmogenic nuclide analyses (¹⁰Be; ²¹Ne) and OSL rock surface dating suggests the potential of these techniques as a tool for **deciphering relative intra- and inter-site chronologies** and constraining timescales of activity/stability patterns. A parallel **downslope increase of terrestrial cosmogenic nuclide concentrations** on PAP4 and PAP8 supports the assumption of successive downslope transport of clasts, likely related to zebra stripe formation. Comparative OSL rock surface dating experiments suggest that **active stripes contain clasts with bleaching fronts on both surface and bottom sides**, pointing to late Pleistocene or Holocene clast overturning. Inactivity is indicated by similar IR50 and pIR225 bleaching curves (equilibrium of bleaching and dosing) in upper clast surfaces and the lack of a bleaching front at the bottom side, pointing to long exposures without clast overturning.

OSL rock surface dating



Future perspectives

Future investigations will concentrate on:

- The role of (sub-) surface gypsum dynamics and the significance of (rare) fog influence (shrink-swell dynamics?)
- Long term soil-moisture and temperature trends (freeze-thaw processes?)
- Measurements of local seismic activity
- Particle-based simulations of granular flows and stripe formation
- Constraining timescales of zebra stripe formation
- Focusing on intersite comparisons