

1-D photochemical model to predict oxygen isotope anomalies in early Earth atmospheres

Bethan Gregory^{1,2*}, Mark Claire^{1,2,3}, and Sarah Rugheimer^{1,4}

*Final-year PhD candidate; correspondence email: bsg@st-andrews.ac.uk

¹School of Earth and Environmental Sciences, University of St Andrews, UK; ²Centre for Exoplanet Sciences, University of St Andrews, UK; ³Blue Marble Space Institute of Science, 1001 4th Ave, Suite 3201, Seattle, USA; ⁴Dept. of Atmospheric Physics, University of Oxford, UK

The goal of this project is to reduce uncertainties in the evolution of O₂ levels over Earth history, using a 1-D photochemical model and triple oxygen isotope data.

1. The photochemical model, Atmos




Fig. 1: Schematic of model set-up. Adapted from Claire, personal communication.

We use and develop the 1-D coupled photochemical-climate model Atmos (e.g. [1, 2, 3, 4]).

The model calculates production and loss of atmospheric species at each altitude layer, due to chemical reactions and vertical transport (Fig. 1).

Photolysis, lightning, particulate formation & deposition, rain, and influx of gases are included.

Outputs include species mixing ratio profiles.

2. Modelling of oxygen levels suggests two stable states





Fig. 2 (above): O₃ column density with O₂ mixing ratio for 2024 model atmospheres (from Gregory et al., in review). For Case 1, we varied O₂ flux and kept CH₄:O₂ flux ratio constant; for Case 2, we varied CH₄:O₂ flux ratio at constant O₂ flux; for Case 3, we included an oxidative weathering flux term for O₂.

Fig. 3 (right): Potential constraints on oxygen levels over Earth history, from the clustering of model solutions from this study with high-O₂ or trace-O₂ mixing ratios (purple regions)^[8]. Adapted from [9].

- The model Atmos is run by varying lower boundary conditions. For O₂, mixing ratio or flux boundary conditions can be chosen.
- Previous fixed mixing ratio-driven models can simulate atmospheres with O₂ ground mixing ratios between 2.1x10⁻⁶ and 0.21^[5, 6, 7], but many of these model atmospheres may be unstable equilibrium solutions.
- Models with flux boundary conditions produce few stable solutions for atmospheres with ground-level O₂ mixing ratios between 6x10⁻⁷ and 2x10⁻³ (1% of present atmospheric levels) (Figure 2, Gregory et al., in review ([8])).
- This provides a **potential lower limit** on Proterozoic O₂ levels (Figure 3).



3. Incorporating triple oxygen isotopes into Atmos in order to predict Δ¹⁷O

i) Background to atmospheric Δ¹⁷O

In the formation of stratospheric ozone:

O₃ gains a **large, positive** Δ¹⁷O value...

... which is propagated to O(¹D), CO₂, NO₃⁻, SO₄²⁻, and H₂O₂ via chemical reactions (e.g. [10]).

Due to mass balance, O₂ has a **small, negative** Δ¹⁷O value, dependent on pCO₂, pO₂, and primary productivity...

...which can be incorporated into stable sulphate and preserved in the geological record.

(where Δ¹⁷O = δ¹⁷O - 0.528 δ¹⁸O)


ii) Developing Atmos

We have incorporated the three isotopes of oxygen in order to predict Δ¹⁷O profiles of atmospheric species and Δ¹⁷O values of species reaching the Earth's surface through wet and dry deposition.

iii) Model validation and calibration...

... involves comparison with existing data (e.g. [11, 12, 13]), existing model results (e.g. [14, 15]), and recently-collected salt concentration and oxygen isotope data from Atacama nitrates and sulphates.

Fig. 4: Collecting samples from the Atacama Desert, Chile (Dec 2017). Credit: A. Zerkle.



iv) Research in progress

- Q What is the minimum pO₂ required for a non-zero Δ¹⁷O to be measurable in geological record sulphates?
- Q What kind of Δ¹⁷O_{O₂} values might be expected for the high-O₂ and trace-O₂ atmospheres from the flux-driven modelling results presented above (Figure 3)?

SUMMARY: We are developing an oxygen isotope model to predict Δ¹⁷O in different species under different conditions, in order to explore our hypothesis of two stable states for oxygen concentrations.

Fig. 5: Maria Elena, Atacama Desert, Chile

References

[1] Zahnle, K., Claire, M. and Catling, D., 2006. The loss of mass-independent fractionation in sulfur due to a Palaeoproterozoic collapse of atmospheric methane. *Geobiology*, 4(4), pp.271-283. [2] Catling, D.C., Claire, M.W., Zahnle, K.J., Quinn, R.C., Clark, B.C., Hecht, M.H. and Kounaves, S., 2010. Atmospheric origins of perchlorate on Mars and in the Atacama. *Journal of Geophysical Research: Planets*, 115(E1). [3] Domagal-Goldman, S.D., Segura, A., Claire, M.W., Robinson, T.D. and Meadows, V.S., 2014. Abiotic ozone and oxygen in atmospheres similar to prebiotic Earth. *The Astrophysical Journal*, 792(2), p.90. [4] Claire, M.W., Kasting, J.F., Domagal-Goldman, S.D., Stüeken, E.E., Buick, R. and Meadows, V.S., 2014. Modeling the signature of sulfur mass-independent fractionation produced in the Archean atmosphere. *Geochimica et Cosmochimica Acta*, 141, pp.365-380. [5] Kasting, J.F. and Donahue, T.M., 1980. The evolution of atmospheric ozone. *Journal of Geophysical Research: Oceans*, 85(C6), pp.3255-3263. [6] Kasting, J.F., Holland, H.D. and Pinto, J.P., 1985. Oxidant abundances in rainwater and the evolution of atmospheric oxygen. *Journal of Geophysical Research: Atmospheres*, 90(D6), pp.10497-10510. [7] Segura, A., Krelow, K., Kasting, J.F., Sommerlatt, D., Meadows, V., Crisp, D., Cohen, M. and Milaver, E., 2003. Ozone concentrations and ultraviolet fluxes on Earth-like planets around other stars. *Astrobiology*, 3(4), pp.689-708. [8] Gregory, B. S., Claire, M. W. and Rugheimer, S., in review. Photochemical modelling of atmospheric oxygen levels suggests two stable states. [9] Kump, L.R., 2008. The rise of atmospheric oxygen. *Nature*, 451(7176), pp.277-278. [10] Thiemens, M.H., 2006. History and applications of mass-independent isotope effects. *Annu. Rev. Earth Planet. Sci.*, 34, pp.217-262. [11] Michalski, G. et al., 2003. First measurements and modeling of 17O in atmospheric nitrate. *Geophys. Res. Lett.*, 30(16). [12] Michalski, G., Böhlke, J.K. & Thiemens, M., 2004. Long term atmospheric deposition as the source of nitrate and other salts in the Atacama Desert, Chile: New evidence from mass-independent oxygen isotopic compositions. *Geochimica et Cosmochimica Acta*, 68(20), pp.4073-4088. [13] Bao, H. & Gu, B., 2004. Natural perchlorate has a unique oxygen isotope signature. *Environmental Science and Technology*, 38(19), pp.5073-5077. [14] Young, E.D., Yeung, L.Y. & Kohl, I.E., 2014. On the Δ¹⁷O budget of atmospheric O₂. *Geochimica et Cosmochimica Acta*, 135, pp.102-125. [15] Lyons, J.R., 2001. Transfer of mass-independent fractionation in ozone to other oxygen-containing radicals in the atmosphere. *Geophysical Research Letters*, 28(17), pp.3231-3234.



University of
St Andrews

Sharing
Geoscience
Online, EGU,
5th May 2020

