


Using the Net Carbon Profit for optimizing vegetation properties along a precipitation gradient

R.C. Nijzink¹, J. Beringer², L.Hutley³, S. Schymanski¹

¹ Luxembourg Institute of Science and Technology, Belvaux, Luxembourg,


²University of Western Australia, Crawley, Australia

³Charles Darwin University, Darwin, NT, Australia




TERRESTRIAL BIOSPHERE MODELS TODAY


Some data of the latent heat flux, Howard Springs, Australia




Some models seem pretty good!



Others underestimate seasonality



And some overestimate seasonality



Many different outcomes for different reasons!

- BESS
- BIOS2
- LPJ-GUESS
- SPA
- CABLE
- MAESPA
- Schymanski et al. (2015)
- Flux tower
- VOM

Introduction

Methods

Results

Conclusions

Previous


Home

Next


TERRESTRIAL BIOSPHERE MODELS TODAY

What's going on... ?

Models use Plant Functional Types and prescribe constant rooting depths ...




Vegetation cover and leaf area index are often prescribed on mean monthly remotely sensed data...




Relying on data and past observations !!!

Models often assume freely draining conditions...



Unknown parameters often tuned for specific conditions or calibrated on observations...



Introduction

Methods

Results

Conclusions

Previous

Home

Next

VEGETATION OPTIMALITY


Net Carbon Profit :

Total difference of carbon uptake by photosynthesis and carbon costs of the system



Vegetation Optimality Model

Optimizes vegetation properties to maximize NCP



No vegetation data needed

No calibration on historical data

Introduction

Methods

Results

Conclusions

Previous

Home

Next

More info

HYPOTHESES



- Conventional models capture the temporal and spatial variation of carbon and water fluxes better compared to the optimality-driven model.
[Go to results →](#)
- Optimality-based dynamics of vegetation cover will lead to worse reproduction of fluxes compared to using mean monthly vegetation cover values for each site obtained from remote sensing time series.
[Go to results →](#)
- Optimality-based rooting depths will not result in better reproduction of carbon and water fluxes compared to a prescribed, homogeneous rooting depth.
[Go to results →](#)
- Re-calibration of the costs for the water transport system, i.e. costs for the vascular system in roots, stems and branches, at each site will not result in a large variation of the cost parameter for these costs.
[Go to results →](#)

[Introduction](#)[Methods](#)[Results](#)[Conclusions](#)[Previous](#)[Home](#)[Next](#)


NORTH AUSTRALIAN TROPICAL TRANSECT

- Mean annual rainfall: 500-1800 mm
- Pronounced wet season: Nov-Feb
- Evergreen trees + seasonal grass
- Flux towers for evaporation and CO₂-fluxes

- Introduction
- Methods
- Results
- Conclusions



VEGETATION OPTIMALITY MODEL



Optimized constants

- Tree cover fraction
- Tree rooting depth
- Grass rooting depth
- Water use strategies

Dynamically optimized variables:

- Grass cover fraction
- Photosynthetic capacity
- Stomatal conductances
- Fine root surface area

Introduction

Methods

Results

Conclusions

Previous

Home

Next

NET CARBON PROFIT

Foliage benefits:

- Bigger foliage
→ more CO₂-uptake

Assimilation

Water transport benefits

- Bigger roots
→ more root water uptake
- Bigger vegetated area
→ more CO₂-uptake

Foliage costs

Water transport costs

Root benefits

- More roots
→ more water uptake

Root respiration

+ photosynthesis

- foliage costs
- water transport costs
- root respiration

----- +
Net carbon profit

Introduction

Methods

Results


Conclusions

Previous

Home

Next

MODEL COMPARISON



The VOM has a correct seasonal amplitude in most cases

Introduction

Methods

Results

Conclusions

- BESS
- BIOS2
- LPJ-GUESS
- SPA
- CABLE
- MAESPA
- Schymanski et al. (2015)
- Flux tower
- VOM

Model data from: Whitley et al. (2015): Biogeosciences 13

Background

Previous Home Next













MODEL COMPARISON

Introduction

Methods

Results

Conclusions




Off-set assimilation during transition from wet to dry period

Previous Home Next

Background

MODEL COMPARISON




VOM underestimates latent heat for the wetter sites

Introduction

Methods

Results

Conclusions



MODEL COMPARISON



- VOM shows spatial pattern similar or better than other models
- Absence of spatial pattern for several other models
- VOM over-estimates assimilation
- Prescribing cover reduces over-estimation assimilation. [More...](#)
- Evaporation at wetter sites under-estimated after prescribing roots. [More...](#)

Introduction

Methods

Results


Conclusions




FOLIAGE COSTS

- Temporal signal largely reproduced
- Model always reaches full cover

100% vegetation cover?



Constructed time series of vegetation cover to prescribe in the VOM



Introduction

Methods

Results

Conclusions

Previous

Home


Next

Background

FOLIAGE COSTS

Prescribing vegetation cover:

- Assimilation generally lower. See also the [model comparison](#).




- Introduction
- Methods
- Results
- Conclusions

Lower assimilation rates when projective cover is prescribed.

FOLIAGE COSTS

Prescribing vegetation cover:

- Assimilation lower, but not always closer to observations



Introduction

Methods

Results

Conclusions

Previous



Home

Next

Background

ROOTING DEPTHS

- Large differences between other models and VOM-results
- Pattern over the transect




LPJ-GUESS:
 Dynamically adapts vegetation and most similar to the VOM →
 Prescribe 2 meter rooting depths to the VOM for comparison

- Introduction
- Methods
- Results**
- Conclusions

ROOTING DEPTHS

- Prescribing roots worsens the under-estimation of E and A in the dry season. See also the [model comparison](#).



Introduction

Methods

Results

Conclusions

Previous


Home

Next

Background

WATER TRANSPORT COSTS

- Fluxes are sensitive to variations in the water transport costs
- Differences occur especially during the dry season



Introduction

Methods

Results

Conclusions

Previous



Home

Next

Background

WATER TRANSPORT COSTS

- Vegetative cover during the dry season sensitive to cost factor



Introduction

Methods

Results

Conclusions

Previous





Home

Next

Background

WATER TRANSPORT COSTS

- Remotely sensed vegetation cover during the dry season only reproduced with different cost factors per site.



Introduction

Methods

Results

Conclusions

Previous

Home

Next

Background

CONCLUSIONS

- The VOM captured the temporal and spatial variation of carbon and water fluxes similar or better compared to the conventional models.
- Optimality-based vegetation cover has a consistent bias during the wet-season reaching full cover. Traditional prescribed vegetation covers lead to lower CO₂-assimilation.
- Optimality-based rooting depths result in a better reproduction of carbon and water fluxes during the dry season.
- Re-calibration of water transport costs for each site resulted in a large variation of the cost parameter for these costs.

Introduction

Methods

Results

Conclusions


Previous

Home


Next

[Back to hypotheses →](#)

APPENDIX



MODEL COMPARISON



This figure shows the time series of an average year, based on the full model runs and smoothed by a 7-day running mean.

The models BESS, BIOS2, LPJ-GUESS, SPA, CABLE and MAESPA were used in a previous model intercomparison study by Whitley et al. (2016). From these models, LPJ-GUESS is the most similar to the VOM, as it uses a carbon allocation scheme to dynamically adapt vegetation. The other models mainly use prescribed vegetation parameters.

The VOM-results of Schymanski et al. (2015) differ from the new VOM-results due to differences in hydrological parameterizations.

Back

Introduction

Methods

Results

Conclusions

Previous


Home

Next

MODEL COMPARISON

- VOM shows spatial pattern similar or better than other models
- Absence of spatial pattern for several other models
- VOM over-estimates assimilation
- Prescribing cover
- Evaporation wet roots. [More...](#)

This figure shows the mean annual fluxes of evaporation and assimilation for the models and observations from the flux towers. The sites are ordered from the wettest site on the left (Howard Springs) to the driest on the right (Sturt Plains).



The VOM shows a clear pattern of decreasing values of evaporation and assimilation. Some models show this too, but MAESPA and LPJ-GUESS do not show a pattern over the transect. At the same time, LPJ-GUESS is actually the most similar to the VOM as it dynamically models vegetation.

Back

Introduction

Methods

Results

Conclusions


Previous

Home

Next

WATER TRANSPORT COSTS

- Fluxes are sensitive to variations in the water transport costs
- Differences occur especially during the dry season



This figure shows the fluxes of evaporation and assimilation for different values of the cost factor for water transport c_{rv} in comparison with flux tower observations (blue).

Water transport costs are a function of rooting depth (y_r), vegetated fraction (M_A , -) and a cost factor (c_{rv} , $mol s^{-1} m^{-3}$):

$$R_v = c_{rv} * M_A y_r$$

Different values of this cost factor lead to different model fluxes especially during the dry season, as can be seen from this figure.

Back

Introduction

Methods

Results

Conclusions


Previous

Home

Next

WATER TRANSPORT COSTS

- Vegetative cover during the dry season sensitive to cost factor



This figure shows the projective vegetation cover for different values of the cost factor for water transport c_{rv} in comparison with remote sensed vegetation cover (blue).

Water transport costs are a function of rooting depth (y_r), vegetated fraction (M_A , -) and a cost factor (c_{rv} , $mol\ s^{-1}\ m^{-3}$):

$$R_v = c_{rv} * M_A y_r$$

Different values of this cost factor lead especially during the dry season to differences in the vegetative cover, as can be seen from this figure.

Back

Introduction

Methods

Results

Conclusions




Previous

Home

Next

WATER TRANSPORT COSTS

- Remotely sensed vegetation cover during the dry season only reproduced with different cost factors per site.



This figure shows the mean vegetation cover during the dry season (May-Sept.) for vegetation cover derived from fPar (Donohue et al. 2016) and the results of the VOM for different values of the cost factor for water transport.

It can be noted here that each site needs a different value of this cost factor in order to reproduce the remotely sensed vegetation cover. It could be argued that this cost factor is not constant over the transect and is a function of other climatic characteristics.

Introduction

Methods

Results

Conclusions

Previous

Home


Next

Back

FOLIAGE

- Temporal
- Model alw

This figure shows the mean vegetation cover predicted by the VOM (red) and derived from remotely sensed fPar-data (blue). It can be noted that the VOM always reaches 100% full cover during the wet season, which is not realistic. This happened consistently for all six study sites along the NATT.



Time series of vegetation cover were constructed based on the remotely sensed vegetation cover in order to prescribe to the VOM. This was done in order to assess whether improvements in projective cover would also reduce over-estimations in the especially CO2-assimilation.

The time series were constructed in two ways:

1. The mean monthly values of vegetation cover were repeated for all years, which is a common approach in land surface modelling.
2. The actual values of remotely sensed vegetation cover were used, but extended with the mean monthly values to cover the full model period.


Construct
cover to

- Introduction
- Methods
- Results
- Conclusions

FOLIAGE COSTS

Prescribing vegetation cover:

- Assimilation generally lower. See also the [model comparison](#).



This figure shows the CO₂-assimilation for the VOM predicting vegetation cover and the VOM that uses prescribed vegetation cover.

The assimilation rates are generally lower for the VOM with prescribed vegetation cover. This relates to the prescribed vegetation cover being generally lower compared to the predicted vegetation cover.

Introduction

Methods

Results

Conclusions

Previous

Home


Next

Back

FOLIAGE COSTS

Prescribing vegetation cover:

- Assimilation lower, but not always closer to observations



The mean annual values of CO₂-assimilation show also that prescribing vegetation cover leads to lower assimilation rates. However, it is not directly clear whether prescribed or predicted vegetation cover leads to mean annual fluxes closer to the observations.

Back

Introduction

Methods

Results

Conclusions


Previous

Home

Next

ROOTING DEPTHS

- Large differences between other models and VOM-results
- Pattern over the transect



This optimized rooting depths for trees and grasses show a pattern over the transect. In comparison with other models, the rooting depths also remain rather shallow. The VOM results of 2015 (Schymanski et al. 2015) predict different rooting depths due to a different schematization and parameterization of the hydrology.

LPJ-GUESS is the most similar to the VOM, as it uses a carbon allocation scheme to dynamically model vegetation. Therefore, the rooting depths of LPJ-GUESS were used to prescribe to the VOM for comparison.

Introduction

Methods

Results

Conclusions

← Previous


Home

Next →

Back →

ROOTING DEPTHS

- Prescribing roots worsens the under-estimation of E and A in the dry season. See also the [model comparison](#).



This VOM-fluxes with optimized rooting depths show strong differences with the results with prescribed rooting depths. Especially during the dry season, differences occur.

Introduction

Methods

Results

Conclusions

← Previous

Home

Next →

Back →