



PLAN
LOIRE
GRANDEUR NATURE

l'Europe
s'engage
le bassin de la Loire
Fonds Européen
de Développement Régional



université
de TOURS

INRAE

➤ Identifying the influence of dams and ponds on the thermal regime at regional scale

The case of Loire catchment

Hanieh Seyedhashemi; Florentina Moatar; Jean-Philippe Vidal;
Aurelian Beaufort; André Chandesris; Laurent Valette

INRAE, RiverLy, Villeurbanne, France



Stream (water) Temperature (ST)



- A key factor in the distribution of aquatic communities (Poole et al., 2001);
- Modified by natural processes and human activities differently (Webb, 1996);
- Exacerbated modifications due to climate change (Webb, 1996; Moatar and Gailhard, 2006; Michel et al., 2020);
- Lack of data on both upstream and downstream of anthropogenic structures (Hill et al., 2013) ;
- New tools required for identifying and predicting human impacts.

Objectives

- Defining thermal signatures to identify human impacts on ST
- Testing them on a large database of ST over a large French basin
- Distinguishing between natural regimes and altered ones
- Detecting the impacts of dams and cumulative effects of upstream ponds

Study area and data

Loire basin and standing surface waters



Loire at the entry of estuary



Coise river

0.8% of the catchment areas with standing surface waters

73 large dams



0.3% of the catchment areas with ponds

One of the largest European catchments (10^5 km^2)

Contrasting natural conditions

Study area and data


Observed stream and air temperature



526 stations of observed ST
2008-2018 with some gap years


Air temperature from Safran reanalysis data
8~km spatial resolution and daily temporal
resolution
(Quintana-Segui et al., 2008; Vidal et al., 2010)

Examples of human impacts: Large dams



✓ Decrease ST and delay the annual cycle


Examples of human impacts: Ponds and lakes



✓ Increase ST and impose a vertical shift in regime

Examples of stream-air temperature relationship: dams and ponds

Using air temperature as a proxy of the heat budget (Mohseni et al., 1999, Caissie et al., 2008)



✓ Lower slope (TS) and R^2 for dams compared to ponds

Methodology

Step 1

Selecting stations that are sensitive to alterations



Step 2

Defining thermal signatures by comparing stream and air temperature based on seasonal regime and linear regression



Step 3


Identifying different thermal regimes through clustering based on thermal signatures




Step 4

Comparing the regimes of different clusters

Step 1: Selecting stations that are sensitive to alterations






(Caissie et al., 2006)




- ✓ The larger a river, the larger its volume (thermal capacity) and the less responsive it is to the alterations

Step 2: Defining thermal signatures by comparing stream and air temperature based on seasonal regime and linear regression



Dam Impact




Pond Impact


Step 2: Defining thermal signatures by comparing stream and air temperature based on seasonal regime and linear regression

Notation	Definition
Dam signatures	
TS	slope
R^2	coefficient of determination
Lag time	$day(Tw_{peak}) - day(Ta_{peak})$
Pond signatures	
SumAbs(Tw-Ta)	$\sum_{Mar.-Oct.}^{Tw>Ta} (Tw - Ta) / nbdays$
Mean(Tw-Ta)	$\sum_{Mar.-Oct.} (Tw - Ta) / nbdays$

Step 3: Identifying different thermal regimes through clustering based on thermal signatures



Step 4: Comparing the regimes of different clusters



- ✓ Dams:
 - delay the annual cycle by 18 days;
 - decrease ST by 2.1°C in average over the summer

- ✓ Cumulative effects of upstream ponds:
 - increase ST by 2.7°C in average over the summer
 - exacerbate the effect in a hot year with 2°C increase in ST in average over the summer

Conclusion and Perspectives

- Dealing with spatial and temporal gaps in ST data
- Defining five thermal signatures by comparing stream and air temperature
- Distinguishing between natural regimes and altered ones
 - Dams:
 - Delay the annual cycle by 18 days;
 - Decrease ST by 2.1°C in average over the summer
 - Cumulative effects of ponds:
 - increase ST by 2.7°C in average over the summer
 - exacerbate the effect in a hot year with 2°C increase in ST in average over the summer
- Identifying highly influenced streams, and taking mitigation actions
- Designing strategic network surveys
- Using natural thermal regime for developing a reference-condition numerical model



Thank you for your attention!

Hanieh.seyedhashemi@inrae.fr



INRAE

Identifying the influence of dams and ponds on the thermal regime

May 7th 2020 / Hanieh Seyedhashemi

Selective References:

Caissie, D. (2006). The thermal regime of rivers: a review. *Freshwater biology*, 51 , 1389-1406.

Chandesris, A., Van Looy, K., Diamond, J. S., & Souchon, Y. (2019). Small dams alter thermal regimes of downstream water. *Hydrology & Earth System Sciences*, 23 .

Dripps, W., & Granger, S. R. (2013b). The impact of artificially impounded, residential headwater lakes on downstream water temperature. *Environmental earth sciences*, 68 , 2399-2407.

Hill, R. A., Hawkins, C. P., & Carlisle, D. M. (2013). Predicting thermal reference conditions for usa streams and rivers. *Freshwater Science*, 32 , 39-55.

Olden, J. D., & Naiman, R. J. (2010). Incorporating thermal regimes into environmental Ows assessments: modifying dam operations to restore freshwater ecosystem integrity. *Freshwater Biology*, 55 , 86-107.

Maheu, A., St-Hilaire, A., Caissie, D., El-Jabi, N., Bourque, G., & Boisclair, D. (2016c). A regional analysis of the impact of dams on water temperature in medium-size rivers in eastern canada. *Canadian Journal of Fisheries and Aquatic Sciences*, 73 , 1885-1897.

Moatar, F., & Gailhard, J. (2006). Water temperature behaviour in the river loire since 1976 and 1881. *Comptes Rendus Geoscience*, 338 , 319-328.

Mohseni, O., Erickson, T. R., & Stefan, H. G. (1999). Sensitivity of stream temperatures in the united states to air temperatures projected under a global warming scenario. *Water Resources Research*, 35 , 3723-3733.

Michel, A., Brauchli, T., Lehning, M., Schaei, B., & Huwald, H. (2020). Stream temperature and discharge evolution in switzerland over the last 50 years: annual and seasonal behaviour. *Hydrology & Earth System Sciences*, 24 .

Poole, G. C., & Berman, C. H. (2001). An ecological perspective on in-stream temperature: natural heat dynamics and mechanisms of human-caused thermal degradation. *Environmental management*, 27 , 787-802.

Webb, B. (1996). Trends in stream and river temperature. *Hydrological processes*, 10 , 205-226.

