

Microbial mechanisms in biomineralization



Erika Kothe, Eileen Meier, Julia Kirtzel, Flavio Costa.


Institute of Microbiology, Friedrich Schiller University Jena, Germany

The concept of biomineralization distinguishes **microbially controlled** and **microbially induced** formation;
both may be intracellular or extracellular



Our research on heavy metal resistance mechanisms in bacteria and fungi led us to consider a more microbiological distinction based rather on **metabolic active** or **passive** processes

Crystal formation is dependent on living biomass

Streptomyces acidiscabies forms nickel struvite crystals surrounded by the bacterial mycelium. Autoclaved biomass does not lead to crystal formation



Different strains form different crystals




Habit of single crystals of $\text{Ni}(\text{NH}_4)(\text{PO}_4) \cdot 6\text{H}_2\text{O}$

A) possible habit formed by $\{00-1\}$, $\{100\}$, $\{010\}$, $\{113\}$ and $\{001\}$ faces

B) view direction parallel to $[010]$ of an idealized struvite crystal formed by $\{010\}$, $\{100\}$, $\{103\}$ and $\{10-3\}$ faces

Active pH control leads to different minerals formed






Streptomyces lividans TK 23 acidifies the medium



Streptomyces naganishii P9A-1 leads to alkalization (without urea in the medium!)



Minerals at different distance to the colony



Genetic determinants

hoxN: Ni/Co permease



nreB: Ni efflux transporter

copY & *copZ*: Cu sensing regulation system



A megaplasmid of *Streptomyces mirabilis* P16-B1 with several metal resistance genes leads to enhanced biomineral formation if transferred to a metal sensitive *S. lividans*.


The plasmid is naturally transferable in soil microcosms




Bioweathering and biomineralization are connected

The basidiomycete fungus *Schizophyllum commune* can degrade shale (by laccase involvement)

This releases metals which induce metal resistance gene (as shown with transcriptome and proteome studies)



Oxalates are formed, likely connected to extracellular matrix (the glucan schizophyllan could be shown)






Thanks to all current and former members of the Microbial Communication lab!!

Georg Büchel
Dirk Merten
Applied Geology,
FSU

Sandor Nietzsche
ElMi Center

Falko Langenhorst
Juraj Majzlan
Mineralogy




Natalie Wielsch
Ales Svatos
Willi Boland
MPI Chemical
Ecology

Geoffrey Gadd
Dundee, UK

Volker Deckert
Tanja Gaudig
IPHT and FSU Jena



„Alteration and element mobility
at the microbe-mineral interface“

Research Training  Deutsche
Forschungsgemeinschaft