

Reactive transport of dichloromethane in laboratory aquifers: insights from dual-element isotope analysis and biomolecular approaches

M. Prieto-Espinoza^{1*}, S. Weill¹, B. Belfort¹, F. Lehmann¹, J. Masbou¹, E. Muller², S. Vuilleumier², G. Imfeld^{1*}

¹ University of Strasbourg, CNRS/EOST, LHyGeS UMR 7517, Laboratory of Hydrology and Geochemistry of Strasbourg, Strasbourg, France

² University of Strasbourg, CNRS, GMGM UMR 7156, Génétique Moléculaire, Génomique, Microbiologie, Strasbourg, France

Dichloromethane in aquifers

Dichloromethane (DMC) is one of the most common chlorinated methanes often detected in groundwater as a result of extensive use, inappropriate disposal and accidental spills^{1,4}.

Why studying water table fluctuations?



- > Spreading of pollutants across the saturated and unsaturated zone (e.g., volatile compounds).
- > Changes in redox conditions and microbial composition that may further impact pollutant degradation.

Research questions



- > What is the impact of water table fluctuations on the extent of DCM *in-situ* degradation in aquifers?
- > What are the main mechanisms and pathways of DCM dissipation in aquifers?

Dual-element CSIA and high-throughput analysis may help?

Laboratory aquifers under near-natural settings




Dimensions: 160 x 80 x 7 cm³
 Flow rate: 0.33 mL/min
 DCM injection: 0.45 mM/L
 Oxygen content: < 0.3 mg/L
 Residence time: 31 days




Material and methods

High-resolution sampling:

- Quantification of DCM
- Isotope analysis of ^{13}C (GC-IRMS) and ^{37}Cl (GC-MS)
- DNA analysis
- Hydrochemical parameters



Iron-reducing conditions in both aquifers





Core sampling for DNA analysis





Mass dissipation and carbon isotope composition of DCM


Fluctuating condition



Steady-state condition



Dual-element CSIA: mechanisms of DCM degradation




- Dual C-Cl isotope plot of DCM degradation under transient (red dots) and steady-state (gray triangles) conditions.
- Reported mechanisms by *Hyphomicrobium MC8b*², *D. formicoaceticum*³, *Ca. Dichloromethamonas elyunquensis*^{3,6} and *Dehalobacterium sp.*⁴ have been added for comparison.

- Similar mechanisms as *Ca. Dichloromethamonas elyunquensis*.
- Combination of different mechanisms (apparent).


Unknown mechanism under strictly anoxic conditions.

Bacterial community analysis



- NMDS ordination of bacterial communities in pore water and sand compartment. Depth in cm from surface to bottom.

Evidence of anaerobic DCM degraders:
Dehalobacterium⁴, Geobacter⁵,
Desulfosporosinus^{6,7}.



- Representation of relative abundance of taxa from phylum to genus level in pore water.

Highlights

- Pronounced carbon isotope fractionation of DCM associated with large DCM mass removal under fluctuating conditions (>90%) compared to steady-state conditions (mass removal of 35%).
- Distinct DCM degradation pathways under steady and fluctuating conditions:
=> mechanistically distinct C-Cl bond cleavage reactions subjected to microbial adaptations during dynamic hydrogeological conditions?
- Occurrence of anaerobic DCM degraders under both steady and fluctuating conditions:
=> supports DCM degradation under iron-reducing prevailing conditions.

Water table fluctuations enhance DCM biodegradation and influence DCM degradation pathways compared to steady-state conditions.

References

1. Hermon, L.; Denonfoux, J.; Hellal, J.; Joulain, C.; Ferreira, S.; et al. 2018. Dichloromethane biodegradation in multi-contaminated groundwater: Insights from biomolecular and compound-specific isotope analyses. *Water Res.* 142, 217–226.
2. Heraty, L.J., Fuller, M., Huang, L., Abrajano, T., Sturchio, N., 1999. Isotopic fractionation of carbon and chlorine by microbial degradation of dichloromethane. *Org. Geochem.* 30, 793–799.
3. Chen, G., Shouakar-Stash, O., Phillips, E., Justicia-Leon, S.D., Gilevska, T., Sherwood Lollar, B., Mack, E.E., Seger, E.S., Löffler, F.E., 2018. Dual carbon-chlorine isotope analysis indicates distinct anaerobic dichloromethane degradation pathways in two members of the *Peptococcaceae*. *Environ. Sci. Technol.* 52, 8607–8616.
4. Blázquez-Pallí, N., Shouakar-Stash, O., Palau, J., Trueba-Santiso, A., Varias, J., Bosch, M., Soler, A., Vicent, T., Marco-Urrea, E. and Rosell, M., 2019. Use of dual element isotope analysis and microcosm studies to determine the origin and potential anaerobic biodegradation of dichloromethane in two multi-contaminated aquifers. *Science of The Total Environment*, 696, p.134066.
5. Yoshikawa, M., Zhang, M. and Toyota, K., 2016. Integrated Anaerobic-Aerobic Biodegradation of Multiple Contaminants Including Chlorinated Ethylenes, Benzene, Toluene, and Dichloromethane. *Water, Air, & Soil Pollution*, 228(1).
6. Kleindienst, S., Higgins, S., Tsementzi, D., Chen, G., Konstantinidis, K., Mack, E. and Löffler, F., 2017. ‘*Candidatus* Dichloromethanomonas elyunquensis’ gen. nov., sp. nov., a dichloromethane-degrading anaerobe of the Peptococcaceae family. *Systematic and Applied Microbiology*, 40(3), pp.150-159.
7. Wright, J., Kirchner, V., Bernard, W., Ulrich, N., McLimans, C., Campa, M., Hazen, T., Macbeth, T., Marabello, D., McDermott, J., Mackelprang, R., Roth, K. and Lamendella, R., 2017. Bacterial Community Dynamics in Dichloromethane-Contaminated Groundwater Undergoing Natural Attenuation. *Frontiers in Microbiology*, 8.

THANK YOU!

¹LHyGES:

Gwenaël Imfeld
Sylvain Weill
Benjamin Belfort
François Lehmann
Raphaël Di Chiara
Benoît Guyot
Jérémy Masbou
Colin Fournet
Thierry Perrone

²GMGM:

Stéphane Vuilleumier
Emilie Muller
Carmen Lazaro

*E-mail corresponding authors:

M. Prieto-Espinoza: prietoespinoza@unistra.fr
G. Imfeld: imfeld@unistra.fr

This project has been funded by the EC2CO – CNRS INSU and the Ecole doctorale Sciences de la Terre et Environnement (ED413).

