



Snowmelt contribution to seasonal runoff

Lessons learned from using a bucket-type model on a large set of catchments

Michal Jeníček, Ondřej Ledvinka

michal.jenicek@natur.cuni.cz, <https://www.natur.cuni.cz/geography/physgeo>




FACULTY OF SCIENCE
Charles University



MOTIVATION, RESEARCH QUESTIONS, OBJECTIVES


- 1) to quantify how inter-annual variations in snow storages affect spring and summer runoff, including summer low flows
- 2) to assess the importance of snowmelt in generating runoff compared to rainfall

STUDY AREA: 59 MOUNTAIN CATCHMENTS




HBV-LIGHT MODEL

1. HBV-light model (Seibert and Vis, 2012)
2. Observed data (1980-2014): precipitation, air temperature, discharge and SWE (CHMI)
3. Three objective functions used to calibrate the model; goodness-of-fit assessed against Q and SWE.
4. 100 best parameter sets calibrated resulting in 100 simulations for each catchment




STREAMFLOW SIGNATURES

Snowmelt contribution to runoff (Q_s)




Simulated by the HBV model using „effect tracking“ (complete mixing in a virtual mixing tank)

Groundwater recharge (G_w)




In the HBV, it represents the outflow from the soil box into groundwater boxes

Summer baseflow (Q_b)




In the HBV, it represents an outflow from the lower groundwater box (Q_2) controlled by a recession coeff. (K_2)

Summer (JJA) deficit volumes D_v





Calculated from simulated runoff using variable threshold level method ($Q_{90\%}$)

RELATIVE IMPORTANCE OF SNOW ON RUNOFF




(a) A relation between snowfall fraction (S_f) and snow runoff fraction (Q_{sf}) for all study catchments. (b) A dependence of Q_{sf} and S_f difference on snowfall fraction. Individual points represent mean snowfall fractions and snow runoff fractions for snow-poor years (brown points) and snow-rich years (blue points) for individual catchments.

SNOW RUNOFF FOR SNOW-POOR AND SNOW-RICH YEARS




(a) Difference between mean daily Qsf [-] for snow-poor and snow-rich years. Rows represent individual catchments sorted from top to bottom according to mean catchment elevation from highest to lowest (y-axis not-to-scale), columns represent day of year. (b) Difference between mean monthly runoff (blue bars) and cumulative monthly differences in runoff (red line) for snow-poor and snow-rich years.

RECHARGE AND RUNOFF FOR SNOW-POOR AND SNOW-RICH YEARS




Difference in selected signatures for snow-rich and snow-poor years for study catchments; (left) Seasonal recharge fractions (Dec-May), (right) Annual runoff. Colour scale used for snowfall fraction.

LOW FLOW FOR SNOW-POOR AND SNOW-RICH YEARS




Difference in selected signatures for snow-rich and snow-poor years for study catchments; (left) Summer baseflow (JJA), (right) Deficit volumes (JJA). Colour scale used for snowfall fraction.

HBV SNOWFALL/RAIN SEPARATION: MODELLING EXPERIMENTS




- **Threshold temperature T_T :** a parameter included in the HBV snow routine which separates precipitation to snowfall and rain.
- By changing the T_T , the amount of accumulated snow and snowmelt timing can be controlled, while other variables remain unaffected (e.g. precipitation)
- The T_T was progressively changed from -5 °C to $+5\text{ °C}$. Changes in this parameter influenced the simulated snowfall and thus SWE, snowmelt onset, melt rates and melt-out.
- This way, we simulated what would happen if snowfall turns to rain due to increase in air temperature.
- **Results in next two slides.**

CHANGES IN ANNUAL AND SUMMER RUNOFF WITH DECREASING S_f



Relative change of selected signatures with increasing T_T for study catchments. (a) annual runoff, (b) summer (JJA) runoff. Rows represent individual catchments sorted from top to bottom according to mean catchment elevation from highest to lowest (y-axis not-to-scale), columns represent T_T used in model simulations. Colours show normalized values relative to their means (different scales used for individual panels).





CHANGES IN G_W AND D_V WITH DECREASING S_f



Relative change of selected signatures with increasing T_T for study catchments. (c) groundwater recharge (Dec-May) and (d) summer (JJA) deficit volumes. Rows represent individual catchments sorted from top to bottom according to mean catchment elevation from highest to lowest (y-axis not-to-scale), columns represent T_T used in model simulations. Colours show normalized values relative to their means (different scales used for individual panels).

CONCLUSIONS

- About **17-42% of the total runoff** originates as snowmelt, despite the fact that **12-37% of the precipitation** is falling as snow -> snow is **more effective** to generate catchment runoff compared to liquid precipitation.
- **Snow-poor years** are characterized with **lower snow runoff** and **earlier snowmelt** compared to snow-rich years. **Snow** in snow-poor years contributes for **shorter period to runoff**.
- **Combined effect** of summer precipitation and winter snowpack on summer baseflow -> the **summer low flows might drop in the future**.
- **Lower** snowfall fraction -> lower recharge -> lower baseflow -> **lower** annual runoff -> **higher** summer deficit volumes (the latter is unclear for many catchments)



MORE INFORMATION ...

Hydrology and Earth System Sciences

An interactive open-access journal of the European Geosciences Union

| EGU.eu | EGU Publications | EGU Highlight Articles | Contact | Imprint | Data protection |

<https://doi.org/10.5194/hess-2019-611>

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



Preprints

Abstract

Discussion

Metrics

Submitted as: research article

13 Jan 2020

Importance of snowmelt contribution to seasonal runoff and summer low flows in Czechia

Michal Jeníček¹ and Ondřej Ledvinka²

¹Charles University, Department of Physical Geography and Geoecology, Prague, Czechia

²Czech Hydrometeorological Institute, Prague, Czechia

Review status

A revised version of this preprint is currently under review for the journal HESS.

Received: 13 Nov 2019 – Accepted for review: 10 Jan 2020 – Discussion started: 13 Jan 2020

<https://www.hydrol-earth-syst-sci-discuss.net/hess-2019-611/>

