

# Development of a neural network for groundwater recharge estimation in karst aquifers

## – The future of karstic recharge management? –



Tomy-Minh Truong<sup>1,2</sup>, Márk Rudolf Somogyvári<sup>1</sup>, Martin Sauter<sup>2</sup>, Reinhard Hinkelmann<sup>3</sup>, and Irina Engelhardt<sup>1</sup>

### Motivation:

- Empirical and conceptual models often struggle to adequately describe the highly nonlinear recharge process
- Numerical models require a high amount of system information, which is usually not available, and are computational demanding
- Artificial neural networks (ANNs) are capable of modeling any kind of relationship
- Once trained, the computational cost of ANNs is very low → **very suited for real time modeling**

### Study Area:

- South of France
- Mediterranean climate
- 709 mm/a precipitation
- Hydrogeological basin estimated at 380 km<sup>2</sup>
- Recharge area roughly 130 km<sup>2</sup>
- Supplies Montpellier with drinking water (1100 l/s pumping on average)




### Preprocessing:

- No general rules → “what works best” approach

Used methods include:


- Normalizing
- Smoothing
- Input size reduction
- (Seasonal) differencing
- Series decomposition for modular approach




### Inner workings:

- What is the best architecture?  
→ wrapper feature selection approach (basically **brute forcing** many different network parameter combinations)

### Modular network approach:



### Results:



- So far best results if time series is smoothed
- Normalizing the inputs made the network more stable
- Interesting: well pumping seems to have a minor influence despite high pumping rates  
→ might be because of correlation between temperature and pumping

### Prospect:

- Decoupling of temperature and pumping  
→ removing the trend and seasonality by decomposing the time series  
→ removing the trend and seasonality by differencing/seasonal differencing

<sup>1</sup>Chair of Hydrogeology, Technische Universität Berlin, Berlin, Germany

<sup>2</sup>Department of Applied Geology, Georg-August-University Göttingen, Göttingen, Germany

<sup>3</sup>Chair of Water Resources Management and Modeling of Hydrosystems, Technische Universität Berlin, Berlin, Germany