


Seasonal forecasts of discharge in the Upper Danube upstream of Vienna

Session HS4.4: Ensemble and probabilistic hydro-meteorological forecasts: predictive uncertainty, verification and decision making

I. Martin-Santos, M. Herrnegger and H. Holzmann
(*University of Natural Resources and Life Sciences Vienna*)


K. Fröhlich and J. Ostermüller
(*Deutscher Wetterdienst*)




Overview

- Introduction
- Study area
- Hydrological model – COSERO
- Seasonal discharge forecast system
 - Methodology
 - Bias correction
 - Forecast outputs
- Seasonal forecasts assessment
 - Ensemble spread
 - Forecast skill
- Conclusions

Introduction – Climate services



- Increasing needs for transferring the advances in climate research into hydrological applications within the framework of climate services
- Clim2Power is a research project building a bridge between complex seasonal & long-term climate data and targeted, usable information for decision making in the energy sector



Introduction - Seasonal discharge forecasting

- Seasonal forecasts are becoming one of the most important elements in some policy/decision making systems (e.g. energy sector). Reliable forecasts can result in better anticipation of water related risks in the near months
- Seasonal time scale typically up to a year ahead (chaotic nature of climate system will always limit our ability to predict the weather beyond a few days). Seasonal forecasts offer guidance on large-scale weather patterns and whether a given location will more likely behave above-normal or below normal over a month, referring to a variable of interest
- Dynamical predictions (used since mid 1990s) vs statistical predictions (used since the late 1800s)
- The more sophisticated models typically contain atmospheric, oceanic, land surface and sea ice modules at a relatively high level of complexity. Once forecasts have been produced by models, the process is far from finished (bias correction, downscaling techniques, verification assessments needed)

Seasonal discharge forecasting valuable for water sector



Reservoir management



Hydropower generation



Navigation





Agriculture



Drough risk management

Study Area


UPPER DANUBE BASIN upstream of Vienna (Korneuburg)



- Total Area: 101.204 km²
- Spatial discretization:
 - 65 Sub-basins
 - 3777 HRUs
- Elevation range: ~150m to 4000m
- Mean annual prec.: 1100 mm
- Mean annual discharge: 1923 m³/s


Hydrological model

- COSERO model is a continuous, semi-distributed, conceptual rainfall-runoff model (developed HyWa BOKU)
- It accounts for:
 - Accumulation and melting of snow
 - Actual evapotranspiration from interception
 - Snow and soil layer
 - Storage of water in the soil
 - Separation runoff into different runoff components
- by means of a cascade of linear and non linear reservoirs
- Ability for automatic parameter calibration
- Spatial modelling : river basin divided into several subbasin, simulated discharge computed at outlet.




Seasonal discharge forecasting - Methodology


Calibration/Validation Hydrological model







Seasonal Forecasting System



Seasonal discharge forecasting - Methodology




Seasonal discharge forecasting - Methodology





- Spring forecast 2020 (current season)
- SB-65 (Korneuburg – Danube)

Seasonal discharge forecasting – Bias correction



- Hindcasts (1999-2014): Forecasted discharge generally shows an overall over-estimation over the whole catchment
- To solve this issue: Bias correction based on the long-term mean observed discharges using a „Scale factor“




Seasonal discharge forecasting - display

Calibrated - Hydrological
Model COSERO
(Updated Initial State)

Bias correction
(scale factor)


**Ensemble seasonal
discharge forecasts
(monthly aggregated)**

Bias Corrected SPRING FORECAST DISCHARGE - SB23.
year 2013




- Spring forecast 2013 (exceptionally wet season)
- SB-23 (Wasserburg – Inn)

Seasonal Runoff forecast for SB23.
MAM/2013



DISCHARGE ANOMALY PLUME
Seasonal Forecast from February - SB23.

Monthly mean anomalies relative to 1999-2014 hydroclimatology






CAS

- Shaded quantile and anomaly plume charts give an overview of the spread and uncertainty of the discharge forecasts
- Probability forecasts can be aggregated to terciles, classifying each individual ensemble forecast according to pre-defined thresholds computed for the whole reference period (1999-2014)


Seasonal forecasts assessment – Ensemble spread

- Ensemble spread not able to capture discharge in some basins, particularly in Winter (spread too narrow)
- Evaluation: Averaged-percentage of cases, where the observation fell within the ensemble spread over the hindcast period for each sub-basin
- Aggregated to 7 defined sub-catchment (see plots)




Seasonal forecasts assessment – overall performance

Theoretical skill (based on hindcast forecasts)





Only months where forecasts show >50% probability of being drier, normal or wetter are considered for the verification



Percentage of cases where the observation matched the forecasts (dry, wet or normal) for the hindcast period

Aggregated by season and sub-catchment


Actual skill (based on observations)




Seasonal forecasts assessment - issues

How can we improve the forecast skill?

- **Warm-up with finer meteorological input data (e.g. using reanalysis instead of assimilation datasets)**



- **Considering only high probability forecast situations (at least >60% probability of occurrence)**



Conclusions

- The applied bias correction partially compensate errors introduced by using low resolution assimilation data (in comparison to using COSMO-REA6) for initiating system states)
- Observed discharge values are frequently outside of the ensemble spread, especially in Winter
- In summary, the seasonal forecasts show a very low skill regarding discharge predictions. However, in high probability forecast situations (i.e. many ensemble members agree on dry, normal or wet conditions) acceptable skill is achieved in some sub-catchments
- Improvement potentials and challenges: Finer spatial resolution of meteorological data / forecasts (especially for assimilation data)

THANK YOU FOR YOUR ATTENTION

Contact information:

Ignacio Martin Santos, MSc.

ignacio.martin-santos@boku.ac.at

Institute for Hydrology and Water Management (HyWa)
University of natural Resources and Life Sciences, Vienna