

EVALUATING THE ROLE OF BIOMASS IN THE SORPTION OF FOUR UV-FILTERS

X. SANCHEZ-VILA; . P. RODRÍGUEZ-ESCALES

✉ xavier.sanchez-vila@upc.edu; paula.rodriguez.escales@upc.edu

Grup d'Hidrologia Subterrània (GHS)

Universitat Politècnica de Catalunya; Barcelona, Spain



CSIC

CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS



**UNIVERSITAT POLITÈCNICA
DE CATALUNYA
BARCELONATECH**

INTRODUCTION

Why do we have to study UV filters?

UV filters are used as **personal and care products** like sunscreens and cosmetics (perfumes, creams or shampoos), and in a **number of industrial applications**



Some of them are endocrine disruptors and have estrogenic activity

INTRODUCTION

Why do we have to study UV filters?

UV filters are used as **personal and care products** like sunscreens and cosmetics (perfumes, creams or shampoos), and in **a number of industrial** applications

We're killing the world's coral reefs with SUNCREAM and mascara

SLAPPING on the sunscreen is killing off one of the planet's greatest natural spectacles.

By **STUART WINTER**

PUBLISHED: 12:27, Wed, Oct 21, 2015 | UPDATED: 13:46, Wed, Oct 21, 2015

SHARE      



They are also
destroying coral
reefs...

INTRODUCTION

... and what is conditioning their fate in the subsurface?


The fate of UV filters in subsurface is governed by **degradation** and **sorption** processes

Degradation processes:

UV-filters are **recalcitrant compounds** that are degraded under **co-metabolic** processes in the presence of more labile compounds. Most of current models of **degradation of EOCs** only focus on apparent processes

Shaffer et al. 2014.
Water Research

$$\frac{\partial C}{\partial t} = -kC \Rightarrow T_{1/2} = \frac{\ln K}{2}$$



$T_{1/2}$ 4.58 d

Shaffer et al. 2015

Complex processes (like-cometabolism)
needs more complex rates...


INTRODUCTION

... and what is conditioning their fate in the subsurface?

The fate of UV filters in subsurface is governed by **degradation** and **sorption** processes

Sorption processes:

Sorption is also modelled as a phenomenological process...



$$K_{d,i} = \frac{C_{i,s}}{C_{i,w}} = \frac{C_{i,om} f_{oc}}{C_{i,w}} = K_{oc,i} f_{oc}$$

It is not as simple....


INTRODUCTION

... and what is conditioning their fate in the subsurface?

The fate of UV filters in subsurface is governed by **degradation** and **sorption** processes



Sorption processes:

Sorption is also modelled as a phenomenological process...



Sorbate

- Not all EOCs are neutral.
- Ionics EOCs can form ionic forms at certain pH.




INTRODUCTION

... and what is conditioning their fate in the subsurface?

The fate of UV filters in subsurface is governed by **degradation** and **sorption** processes

Sorption processes:

Sorption is also modelled as a phenomenological process...



- **Organic matter** is the most important surface, but not the only one... mineral surfaces can act as sorbents
- What are the real processes of sorption?


INTRODUCTION

... and what is conditioning their fate in the subsurface?

The fate of UV filters in subsurface is governed by **degradation** and **sorption** processes

Sorption processes:

Sorption is also modelled as a phenomenological process...



$$K_d = \frac{C_s}{C_w}$$



$$K_d = \frac{C_{om} f_{om} + C_{min} A + C_{ie} \sigma_{ie} + C_{rxn} \sigma_{rxn} A}{C_{w,neut} + C_{w,ion}}$$

INTRODUCTION


...what is **solid organic matter** in an aquifer?

Sedimentary organic matter

- It is the traditional SOM (0.01-0.1)
- Normally, it is recalcitrant.

Biomass

- Not important if aquifers are not biologically active
- Biomass can be an important sorbent → biofilm formation (e.g. during Managed Aquifer Recharge, bioremediation...)



The goal of this work is to evaluate the effect of redox conditions in the fate of a cocktail of UV-filters and to evaluate the effect of biomass in sorption processes.



Developing a conceptual and a numerical model of sorption and degradation of UV-filters




Validating the model with experimental data of Liu et al. 2013

EXPERIMENTS


EXPERIMENTS

Liu et al (2013) performed a set of experiments evaluating the **degradation of a cocktail of UV-filters** in different redox conditions. They measured the temporal evolution of water and solid concentrations.



EXPERIMENTS


Liu et al (2013) performed a set of experiments evaluating the **degradation of a cocktail of UV-filters** in different redox conditions. They measured the temporal evolution of water and solid concentrations.



		pKa	logD _{ow}
Benzophenone-3	BP3	7.07	3.13
Octyl 4-methoxycinnamate	OMC	Neutral	5.38
Octocrylene	OC	Neutral	6.78
2-(3-t-butyl-2-hydroxy-5-methylphenyl)5-chloro benzotriazole	UVA-326	10.08	5.33
2-(2'-hydroxy-5'-octylphenyl)-benzotriazole	UVA-329	9.30	5.9

CONCEPTUAL AND NUMERICAL MODELS

CONCEPTUAL MODEL



OC_{sed} Sedimentary organic carbon

OC_{lab} Labile organic carbón

TP Transformation product

TEAPs Electron Acceptor

BM Biomass


1 UV-filter sorption

2 Hydrolisis of OC_{sed}

3 Oxidation of labile OC and growth of biomass

4 Co-metabolic degradation of UV-filters

CONCEPTUAL MODEL



OC_{sed} Sedimentary organic carbon

OC_{lab} Labile organic carbon

TP Transformation product

TEAPs Electron Acceptor

BM Biomass

1 UV-filter sorption

2 Hydrolysis of OC_{sed}

3 Oxidation of labile OC and growth of biomass

4 Co-metabolic degradation of UV-filters

NUMERICAL MODEL

1. Extension of the geochemical database:




1. incorporation of the ionic UV-filters into the geochemical speciation and interaction with pH
2. incorporation of sorption reactions with SOM and with biomass

2. Construction of biogeochemical modeling

3. Construction of co-metabolic degradation model coupled to biogeochemical processes

NUMERICAL MODEL

Extension of the geochemical database: incorporation of the ionic UV-filters into the geochemical speciation and interaction with pH

Reaction	logK
Speciation reactions of ionic UV filters ⁽¹⁾	
Speciation of BP3 (as an organic acid) and formation of BP3 ⁻ 	7.07
Speciation of UVA-326 (as an organic acid) and formation of UVA-326 ⁻ 	10.08
Speciation of UVA-329 (as an organic acid) and formation of UVA-329 ⁻ 	9.3

NUMERICAL MODEL

Extension of the geochemical database: incorporation of sorption reactions with SOM and with biomass

$$K_{d_{UV,TOT}} = \sum_{j=1}^j \sum_{i=1}^i K_{d_{i,j}}$$

where "i" is referred to the different sorbents (1=SOM and 2=biomass), and "j" to the different form of the UVs (1=neutral and 2=ionic form).

Sedimentary Organic Matter:

$$K_{d_{UV,SOM}} = \frac{[K_{oc,UV_0} + K_{oc,UV_{[-]}} K_a / [H^+]] f_{oc}}{1 + \frac{K_a}{[H^+]}} \xrightarrow[\substack{K_{oc,UV_{[-]}} \ll K_{oc,UV_{[0]}}}{K_a / [H^+] \ll 1000} K_{d_{UV,SOM}} \sim \frac{K_{oc,UV_0} f_{oc}}{1 + \frac{K_a}{[H^+]}}$$

Biomass:

$$K_{d_{UV,X}} = \frac{[K_{x,UV_{[-]}} K_a / [H^+]] f_{sites}}{1 + \frac{K_a}{[H^+]}}$$

Only ionic compounds interact with biomass

NUMERICAL MODEL

Extension of the geochemical database: incorporation of sorption reactions with SOM and with biomass

Reaction	logK ⁽¹⁾
Sorption of ionic UV filters to SOM ⁽²⁾	
BP3 + ≡SOM ↔ BP3≡SOM	-97.4736
OMC + ≡SOM ↔ OMC≡SOM	-96.6744
OC+ ≡SOM ↔ OC≡SOM	-96.4540
UVA-326 + ≡SOM ↔ UVA-326≡SOM	-96.6803
UVA-329 + ≡SOM ↔ UVA-329≡SOM	-96.6187
Sorption of ionic UV filters to Biomass (X) ⁽³⁾	
BP3 ⁻ + ≡X ↔ BP3 ⁻ ≡X	Fitted
UVA-326 ⁻ + ≡X ↔ UVA-326 ⁻ ≡X	Fitted
UVA-329 ⁻ + ≡SOM ↔ UVA-329 ⁻ ≡SOM	Fitted

Experimentally determined in abiotic experiments (only SOM)

$$K_{d_{UV,X}} = \frac{[K_{x,UV}] K_a / [H^+]] f_{sites}}{1 + \frac{K_a}{[H^+]}}$$

It was assumed 1 (all biomass surface acts as a sorbent)

Co-metabolic degradation: coupling geochemical processes with UV filters degradation

Degradation of labile organic carbon:

$$r_{ED} = -k_{\max} \frac{[ED]}{[ED] + K_{S,ED}} \frac{[EA]}{[EA] + K_{S,EA}} [X]$$

$$r_{EA} = Qr_{ED} - Sb[X]$$

$$r_X = -Y_h r_{ED} - b[X]$$

Release of labile organic carbon from Sedimentary Organic Matter:

$$r_{DOC} = -k_{\max} [SOM]$$


UV filter degradation coupled to oxidation of organic carbon:

$$r_j = -C_j k_{j,i} F_i \quad j = BP3, OMC, OC, UV326, UV329$$

RESULTS


RESULTS

Modeling results: evolution of UV filters in water and solid surfaces




RESULTS

Modeling results: evolution of UV filters in water and solid surfaces



RESULTS


Modeling results: evolution of UV filters in water and solid surfaces



Rodríguez-Escales and
Sanchez-Vila, Water
Research 2020

RESULTS

Effect of redox conditions in the fate of UV filters



Rodríguez-Escales and
Sanchez-Vila, Water
Research 2020

CONCLUSIONS

CONCLUSIONS

1. We have developed a geochemical model for UV filters, that considers co-metabolic degradation and sorption to two different pools of sorbents: sedimentary organic carbon and biomass.
2. In this work we have demonstrated that **biomass can act as an important sorbent for ionic UV filters** (BP-3, UV-326 and UV-329) in porous media.
3. We further conclude that redox conditions, by themselves, do not condition the fate of UV filters (similar degradation constants). Nevertheless, **redox conditions affect the biomass growth, showing a much higher growth of aerobic biomass than anaerobic one, and also modifying the pH.**

More information is available in [Rodríguez-Escales et al. 2020](#)

EVALUATING THE ROLE OF BIOMASS IN THE SORPTION OF FOUR UV-FILTERS

X. SANCHEZ-VILA; . P. RODRÍGUEZ-ESCALES

✉ xavier.sanchez-vila@upc.edu; paula.rodriguez.escales@upc.edu

Grup d'Hidrologia Subterrània (GHS)

Universitat Politècnica de Catalunya; Barcelona, Spain



CSIC

CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS



**UNIVERSITAT POLITÈCNICA
DE CATALUNYA
BARCELONATECH**