



TECHNISCHE
UNIVERSITÄT
DRESDEN



HELMHOLTZ
ZENTRUM FÜR
UMWELTFORSCHUNG
UFZ

DFG

Deutsche
Forschungsgemeinschaft



DAAD

Deutscher Akademischer Austauschdienst
German Academic Exchange Service



Faculty of Environmental Sciences Institute of Groundwater Management



D261 | EGU2020-6347
(8th May 2020)


The Relevance of Fluid and Porous Media Properties for DNAPL Migration and Entrapment: A Numerical Evaluation of Laboratory Experiments

Christian Engelmann
Kaveh Sookhak Lari
Marc Walther

► christian.engelmann@tu-dresden.de


© Engelmann et al. All rights reserved.

Background

- Ongoing worldwide contamination of groundwater bodies by compounds belonging to the chemical group of Dense Non-Aqueous Phase Liquids (DNAPLs)  [next slide](#)

© Engelmann et al. All rights reserved.

Background (modified after [1])



© Engelmann et al. All rights reserved.

Background


- Ongoing worldwide contamination of groundwater bodies by compounds belonging to the chemical group of **Dense Non-Aqueous Phase Liquids (DNAPLs)**
- DNAPLs are toxic, mutagenic, highly persistent, and soluble in water (may reach wells used for water abstraction), so that **assessment and remediation are essential**
- **Source zone geometry (SZG) has major influence on dissolved contaminant plumes, but often considered as over-simplified SZG**
- **Exploration of SZGs limited by technical/economical constraints**
- **Complex processes vs. poor knowledge on SZGs [2] ► next slide**

© Engelmann et al. All rights reserved.

Concept of DNAPL contamination [2]

complex real world / laboratory

(semi-)analytical / numerical implementation



- DNAPL source zone
- DNAPL plume
- Over-simplified implementation of source zone geometry (W... width, M... thickness)
- Effective source zone geometry
- Phase migration
- Mass transfer from organic to aqueous phase
- Advective-dispersive transport incl. sorption and reactions

© Engelmann et al. All rights reserved.

Goals

- Enhance understanding of DNAPL source zone formation
- Identification of factors relevant for phase migration

© Engelmann et al. All rights reserved.

Goals


- Enhance understanding of DNAPL source zone formation
- Identification of factors relevant for phase migration

Approach


- Perform experimental trials to generate observation data
- Develop and apply image processing and analysis (IPA) framework
- Definition of reference model base-case scenarios
- Calibration of numerical multiphase flow model against IPA data
- Estimation of uncertainties and parameter relevance

© Engelmann et al. All rights reserved.

Experimental setup for DNAPL release [3,4,5,6]





 Experiment domain
 IPA domain
 TMVOC model domain



© Engelmann et al. All rights reserved.

IPA framework [3,4,5,6]




© Engelmann et al. All rights reserved.

Modeling strategy ^[4]

- Applied code: parallelized version of TMVOC ^[7], as implemented in TOUGH3 ^[8]
- Assumptions: 2-D, two-phase flow, single component, isothermal, homogeneous domain, no diffusion, no degradation, no reactions
- Quantifiable parameters fixed, variation of type curve parameters α_{nw} and n_{nw} ^[9], manual calibration through Monte Carlo analysis
- Utilization of HPC cluster Taurus (TU Dresden) for simultaneously performing large number of model realizations ($\Sigma \sim 40,000$)
- Use of Python 3.x for automatized pre- and post-processing


© Engelmann et al. All rights reserved.

First exemplary results (work in progress) [4]




Model fitness for glass bead scenario at $t = 20s$ (end of release)

- ... model realization
- 0 ... worst fitness
- 1... best fitness



© Engelmann et al. All rights reserved.

First exemplary results (work in progress) [4]



DNAPL saturation distribution for
glass bead scenario at $t = 20s$
(end of release)

$$n_{nw} = 14.0$$

$$\alpha_{nw} = 28.0 \text{ cm}^{-1}$$

© Engelmann et al. All rights reserved.

Conclusions

- Model results prove high sensitivity of type curve parameters [4]
- Good agreement between observation and simulation data [4]
- Parts of DNAPL saturation within noise spectrum of IPA data (glass beads: < 4%) [4]

© Engelmann et al. All rights reserved.

Conclusions

- Model results prove high sensitivity of type curve parameters [4]
- Good agreement between observation and simulation data [4]
- Parts of DNAPL saturation within noise spectrum of IPA data (glass beads: < 4%) [4]

Outlook

- Finalize calibration for remaining porous media type scenarios [4]
- Quantify and compare uncertainties (experiment, IPA, model) [4]
- Improve relationship between gray-scale intensity and DNAPL saturation to reduce noise spectrum in IPA data
- Add processes to increase system complexity
- Perform SZG predictions

© Engelmann et al. All rights reserved.

References

- [1] **Kueper**, B.H., Stroo, H.F., Vogel, C.M., Wards, C.H. (Eds.), Chlorinated Solvent Source Zone Remediation. Springer, **2014**, p. 713.
- [2] **Engelmann**, C., Händel, F., Binder, M., Yadav, P. K., Dierich, P., Liedl, R., Walther, M. (**2019a**), The fate of DNAPL contaminants in non-consolidated subsurface systems – Discussion on the relevance of effective source zone geometries for plume propagation, Journal of Hazardous Materials 375, 233-240, doi: 10.1016/j.jhazmat.2019.04.083
- [3] **Engelmann**, C., Ibrahim, S.I., Yadav, P.K., Dietrich, P., Liedl, R., Walther, M. (**2019c**), Delineation of Dense Non-Aqueous Phase Liquid Source Zone Geometries and their Dependency on Fluid and Aquifer Properties Utilizing Numerical Modeling, Groundwater Quality conference 2019 (GQ19), Liège, Belgium, 9-13 September 2019
- [4] **Engelmann**, C., Sookhak Lari, K., Schmidt, L., Werth, C.J., Walther, M., Towards Predicting DNAPL Source Zone Formation: A Numerical Evaluation of Laboratory Experiments to Estimate the Relevance of Type Curve Characteristics, **in prep.**
- [5] **Engelmann**, C., Schmidt, L., Ibrahim, S.I., Yadav, P.K., Dietrich, P., Liedl, R., Walther, M. (**2019b**), Delineation of Dense Non-Aqueous Phase Liquid Source Zone Geometries Utilizing Optical Imaging, Digital Image and Signal Processing (DISP) conference, Oxford, U.K., 30 April 2019
- [6] **Engelmann**, C., Schmidt, L., Werth, C.J., Walther, M. (**2019d**), Quantification of Uncertainties from Image Processing and Analysis in Laboratory-Scale DNAPL Release Studies Evaluated by Reflective Optical Imaging, Water 11, 2274, doi: 10.3390/w11112274
- [7] **Pruess**, K., Battistelli, A. (**2002**), TMVOC, A Numerical Simulator for Three-Phase Non-isothermal Flows of Multicomponent Hydrocarbon Mixtures in Saturated-Unsaturated Heterogeneous Media
- [8] **Jung**, Y., Heng Pau, G.S., Finsterle, S., Doughty, C. (**2018**), TOUGH3 User´s Guide
- [9] **Parker**, J.C., Lenhard, R.J., Kuppusamy, T. (**1987**), A Parametric Model for Constitutive Properties Governing Multiphase Flow in Porous Media, Water Resources Research 23 (4), 618-624

© Engelmann et al. All rights reserved.

Funding and legal information

- The authors are thankful for hosting the first author for a short-term research stay at the Commonwealth Scientific and Industrial Research Organisation (CSIRO), Floreat site, Perth, WA, where funding was provided by the German Academic Exchange Service (DAAD). grant number [57511402]
- The authors gratefully acknowledge the support of GWK (Gemeinsame Wissenschaftskonferenz) for funding this study indirectly by providing computing time through the Center for Information Services and HPC (ZIH) at TU Dresden on the HRSK-II.



DAAD

Deutscher Akademischer Austauschdienst
German Academic Exchange Service

Funded by

DFG Deutsche
Forschungsgemeinschaft
German Research Foundation

grant numbers [WA 3973/2-1], [LI 727/29-1],
and [DI 833/22-1]

© Engelmann et al. All rights reserved.