

Drought tolerant quinoa and irrigation scheduling in the Sahel

EGU-2020 (6-8th May)

Jorge Alvar-Beltrán, Ph.D.
jorge.alvar@unifi.it

Prof. Simone Orlandini
Prof. Anna Dalla Marta
Prof. Jacob Sanou
Dr. Abdalla Dao
Dr. José Luís Camacho



UNIVERSITÀ
DEGLI STUDI
FIRENZE



WORLD
METEOROLOGICAL
ORGANIZATION



WORLD
METEOROLOGICAL
ORGANIZATION



RESEARCH COLLABORATION


World Meteorological Organization (WMO)
Switzerland

Flemish Institute Tech. Research (VITO)
Belgium

Institut de l'Environnement et
Recherches Agricoles (INERA)
Burkina Faso

Research Approach: Quinoa

- *Chenopodium quinoa* Willd. Herbaceous, C3 crop
- Traditionally grown in the Andes (7000 years)
- Thrive in a wide range of ecosystems:
Altiplano, Inter-valleys, Salares, Coastal and Yunga
- Genetic diversity (over 16 thousand accessions)
- Abiotic stress resilience:
 - a) Drought (200-400 mm)
 - b) Halophyte (sea water of 600 mM NaCl)
 - c) Frost (-14°C seedling & -4°C milky grains)
 - d) Heat (+40°C)
 - e) pH versatile & poor soils (sandy & low nutrient)
- High nutritional properties
 - Essential amino-acids & high protein content
 - Rich in Ca, Fe & Mg; vitamins A, B2 & E
 - Gluten free



Source: Vacher, 1998; Jacobsen et al., 2003; Mamedí et al., 2007; Jacobsen et al., 2010; Steduto et al., 2012; Fuentes, 2015

Tackling Problem: Quinoa Field Experiments


AIM

- Evaluate the adaptability of quinoa in the Sahel

EXPERIMENTAL DESIGN

2017-2018

4/11 & 8/12




T: Titicaca

N: Negra Collana



2018-2019

25/10 & 19/11



Tackling Problem: Overcoming Research Barriers

IRRIGATION SCHEDULING

Evapotranspiration (ETo in mm)

$$ETo = 0.0023 (T_{mean} + 17.78) * Ro * (T_{max} - T_{min}) ^{0.5}$$

Ro is the solar radiation ($1 \text{ mm day}^{-1} = 2.45 \text{ MJ m}^{-2} \text{ day}^{-1}$).

Ro monthly adjusted during the growing season



Potential Crop ET ($PETc$ in mm)


$$PETc = ETo * Kc$$

Kc is the crop coefficient.

Kc weekly adjusted as follows:

Kc at E, 2L, 4L, 8L, PF, F, MG, PG, PM

Kc values: 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 0.9, 0.8, 0.7



Total Irrigation (TI in l m^{-2} or mm)

$$TI = \left(\left(\frac{m^3}{1000} \right) \div m^2 \right) - 70$$

70 liters of water to attain drip-irrigation working pressure



Tackling Problem: Quinoa Field Experiments

FIELD & LAB MEASUREMENTS

Irrigation

- Amount, frequency & timing

Agro-meteorology


- Tmax, Tmin, T mean, precipitation, RH & PETc
- Soil temperature
- Solar radiation & photoperiodicity

Plant phenology and physiology

- Time E, 2L, 4L, 8L, PF, F, LS, MG, PG, PM
- Plant height, panicle length & width, root architecture, n° of branches & stem diameter
- Kernel weight, biomass/yield production & canopy cover

Soil characteristics

- pH, soil texture, org. matter, N, C, P, K content & bulk density



Tackling Problem: Quinoa Field Experiments

RESULTS

TITICACA VS. NEGRA COLLANA

Crop variety	Titicaca	Negra Collana
Seed yield (kg ha ⁻¹)	686 a	102 b
Biomass (kg ha ⁻¹)	1686 a	1725 a

⚠ *Average of all treatments (irrigation & N-fertilization) & sowing dates (4-Nov and 8-Dec). Experiment 2017-2018*



TITICACA (Two year experiment 4-sowing dates)

- Main effect N-fertilization ($p > 0.05$).
- Main effect Irrigation ($p < 0.05$). 100 & 80 vs. 60 & 50 PETc
- Sowing dates ($p < 0.05$). 25-Oct vs. 8-Dec

Sowing date	25-Oct	4-Nov	19-Nov	8-Dec
Seed yield (kg ha ⁻¹)	1128a	898ab	659ab	540b

⚠ *Titicaca: best irrigation schedules (FI & PD) & all N-fertilization levels (100, 50, 25 & 0 kg N ha⁻¹).*



Publications

Effect of drought and nitrogen fertilisation on quinoa (*Chenopodium quinoa* Willd.) under field conditions in Burkina Faso

Italian Journal of
Agrometeorology
Rivista Italiana di Agrometeorologia



Effetto della siccità e della fertilizzazione azotata su quinoa (*Chenopodium quinoa* Willd.) in Burkina Faso

JORGE ALVAR-BELTRÁN^{1,*}, COULIBALY SATURNIN², ABDALLA DAO², ANNA DALLA MARTA¹, JACOB SANOU², SIMONE ORLANDINI¹



Article

Effect of Drought, Nitrogen Fertilization, Temperature and Photoperiodicity on Quinoa Plant Growth and Development in the Sahel

Jorge Alvar-Beltrán ^{1,*} , Abdalla Dao ², Anna Dalla Marta ¹ , Coulibaly Saturnin ², Paolo Casini ¹, Jacob Sanou ² and Simone Orlandini ¹ 