

Comparing the discriminating power of contrasted sediment tracing techniques to quantify the impact of nickel mining on river and lagoon siltation in New Caledonia

Virginie Sellier, Olivier Evrard, Oldrich Navratil, J. Patrick Laceby,
Michel Allenbach and Irène Lefèvre



EGU Sharing Geosciences online



Contact: virginie.sellier@lsce.ipsl.fr



MINING SOURCES:

Mining sites, overburden
stocks
roads

- Open-cast mining in operation since 1880s
- Strong increase in **soil erosion** and **sediment transfer** in river systems leading to **the island's ecosystem degradation** (e.g. flooding, water pollution)
- Two sediment sources : **Mining sources & Non-mining sources**
- Need to estimate the contributions of suspended sediment to guide the implementation of efficient management measurements

Objectives

- (1) Quantify the ongoing contributions of sediment sources
- (2) Reconstruct the temporal changes of sediment sources



NON-MINING SOURCES:


fires, shallow landslides, cattle
trampling

Thio River catchment (397 km²), one of the first areas exploited for nickel mining

(1) Lag deposits collected following the 'tributary tracing approach' (Lacey et al., 2017)

Legend

- Mining samples
- Non-mining samples
- Thio River samples
- Thio
- Mining tributaries
- Non-mining tributaries
- ▨ Abandoned mining sites
- Active mining sites
- Mining exploration
- Peridotite massifs



(2) Select the optimal sediment fingerprinting method(s) among

5 conventional sediment fingerprinting methods :

- Fallout radionuclides (^{137}Cs , $^{210}\text{Pb}_{\text{xs}}$)
- Geogenic radionuclides (Th, U, K)
- Elemental Geochemistry
- Colour parameters
- Elemental Geochemistry + colour parameters

Reliability of the models

=

Test on artificial mixture samples

One alternative sediment fingerprinting method

- 2015 : Overbank flood by a tropical depression
- 2017 : Overbank flood by Cyclone Cook
- Sediment core in 2016

- Visible spectrum (PLS regression model)

Results

✗ Fallout radionuclides : **low activities**
in mining and non-mining sources

Geogenic radionuclides


Elemental Geochemistry

✗ Colour parameters : Not provide
satisfactory source discrimination

Elemental geochemistry + colour
parameters

✗ Visible spectrum : Not provide
satisfactory source discrimination

Elemental Geochemistry




**83% of variance
explained**

**15% overestimation of
mining sources**

Elemental geochemistry + colour parameters

**93% of variance
explained**


**7% overestimation of
mining sources**



Results

Recent overbank floods 2015-2017

Elemental geochemistry + colour parameters



68 ± 25 % of the Thio River sediment originating from mining sources for the 2015 flood event

$88 \pm 8\%$ for the 2017 flood event

$74 \pm 13\%$ in the sediment core

Sediment core

Elemental Geochemistry

Colour parameters are not conservative

