


Visualizing anthropogenic methane plumes from the California Methane Survey




Andrew K. Thorpe^{1*}, Riley M. Duren^{2,1}, Robert K. Tapella¹, Brian D. Bue¹, Kelsey T. Foster³, Vineet Yadav¹, Talha Rafiq⁴, Francesca M. Hopkins⁴, Kevin M. Gill¹, Joshua D. Rodriguez¹, Aaron Plave¹, Daniel Cusworth¹, Charles E. Miller¹, AVIRIS team, et al.


*Andrew.K.Thorpe@jpl.nasa.gov, ¹Jet Propulsion Laboratory, California Institute of Technology, ²University of Arizona, ³Stanford University, ⁴University of California Riverside




AVIRIS-NG for CH₄ plume mapping



Thorpe et al., 2014, 2016, 2017
Thompson et al., 2015, 2016




AVIRIS-NG for CH₄ emission flux quantification



Duren et al., 2019

California Methane Survey


- AVIRIS-NG flights funded by the California Air Resources Board (CARB) and California Energy Commission (CEC)
- Targeting CH₄ hotspots from the energy, agriculture, and waste management sectors




Methane Source Finder data portal

- Publicly available: <https://methane.jpl.nasa.gov/>
- Developed under NASA's ACCESS and CMS programs to explore CH₄ data
- Data includes:
 - Remote sensing (AVIRIS-NG imaging spectrometer CH₄ plume results; Thorpe et al., 2017)
 - Surface monitoring (in situ towers for CH₄ flux inversions for the LA basin; Yadav et al., 2019)
 - Bottom up CH₄ emissions inventories (EPA)
 - Infrastructure database (potential CH₄ sources; Carranza et al., 2018)
- Goals:
 - Provide CH₄ dataset to a diverse range of stakeholders to improve understanding of regional CH₄ emissions
 - Provide opportunities for mitigation


Methane Source Finder: LA Basin CH₄ hotspots




LA Basin CH₄ from energy sector




LA Basin CH₄ from energy sector



LA Basin CH₄ from landfill



San Joaquin Valley CH₄ emissions



San Joaquin Valley CH₄ from energy sector



San Joaquin Valley CH₄ from dairy digester



Methane Source Finder facilitates stakeholder engagement

CA Methane Survey



BAY AREA
AIR QUALITY
MANAGEMENT
DISTRICT



South Coast
AQMD



CALIFORNIA
AIR RESOURCES BOARD

Energy sector



Waste management sector





REPUBLIC
SERVICES

Dairy groups



Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not constitute or imply its endorsement by the United States Government or the Jet Propulsion Laboratory, California Institute of Technology.

Methane Source Finder facilitates mitigation



California Methane Survey scientific findings

- 1) Multiple revisits of facilities permitted assessment of persistence:
 - Oil & gas, dairy: 20-35% (mean) persistence
 - Landfills: 100% persistence
- 2) Emissions were calculated for 564 methane point sources
- 3) Estimated emissions from methane point sources in California:
 - $0.618 \text{ TgCH}_4 \text{ yr}^{-1}$ (95% confidence 0.523-0.725)
 - Equivalent to 34-46% of 2016 methane inventory
- 4) Super-emitter activity occurs in every surveyed sector (10% of point sources contributed ~60% of point source emissions)

nature > articles > article



Article | Published: 06 November 2019

California's methane super-emitters

Riley M. Duren , Andrew K. Thorpe, Kelsey T. Foster, Talha Rafiq, Francesca M. Hopkins, Vineet Yadav, Brian D. Bue, David R. Thompson, Stephen Conley, Nadia K. Colombi, Christian Frankenberg, Ian B. McCubbin, Michael L. Eastwood, Matthias Falk, Jorn D. Herner, Bart E. Croes, Robert O. Green & Charles E. Miller

Future work: AVIRIS-NG CO₂ and CH₄ (complete carbon footprint)

