

# Statistical Approaches for Modelling Ice Sheet Interconnectivity

Andrew B. Martinez

with


Luke Jackson, Katarina Juselius and Felix Pretis

The views expressed here should not be attributed to the Department of the Treasury or the U.S. Government


- ▶ Sea level rise uncertainty driven by Greenland and Antarctic (Church et al., 2013)
  - ▶ Typically modeled separately and without interactions
- ▶ Empirical linkages exist over very long time scales; Johnsen et al. (1972)
  - ▶ Broecker (1998); Stocker (1998): Bipolar seesaw model
- ▶ Bamber and Aspinall (2013) use expert surveys to derive correlations
  - ▶ Correlations hard to interpret when series are non-stationary
- ▶ We take an empirical approach:
  - ▶ Monthly data on ice sheet mass (IMBIE: Shepherd et al., 2012)
  - ▶ Econometric tools can test for non-stationarity and interdependence jointly
    - ▶ Cointegration captures the long-run relationships (Johansen, 1991)
    - ▶ Simulate how shocks propagate through the system

# IMBIE (2012) Data








### Ice Sheet Mass Acceleration (Gt per month<sup>2</sup>)



### Ice Sheet Mass Balance (Gt per month)



### Ice Sheet Mass (Gt, Base=1992)



If data are non-stationary then empirical correlation coefficient does not capture the true correlation because average is not the expectation:


$$\mathbb{E}(x_{i,t}) \neq \frac{1}{T} \sum_{t=1}^T x_{i,t}$$

# Bivariate Correlations

If data are non-stationary then empirical correlation coefficient does not capture the true correlation because average is not the expectation:

$$\mathbb{E}(x_{i,t}) \neq \frac{1}{T} \sum_{t=1}^T x_{i,t}$$

The recursively estimated correlation coefficients show instability:



→ Cointegration addresses non-stationarity and correlation simultaneously

## I(2) Cointegrated VAR Model

Test for and estimate I(2) CVAR model from Juselius (2006):

$$\begin{bmatrix} \Delta^2 \text{GR}_t \\ \Delta^2 \text{WA}_t \\ \Delta^2 \text{EA}_t \end{bmatrix} = \begin{bmatrix} -0.007 & 0 \\ -0.003 & -0.005 \\ 0.01 & -0.01 \end{bmatrix} \begin{bmatrix} \widetilde{\text{LR1}}_{t-1} \\ \widetilde{\text{LR2}}_{t-1} \end{bmatrix} + \begin{bmatrix} 0.07 & 0.02 \\ 0.007 & 0.21 \\ 0.18 & 0.30 \end{bmatrix} \begin{bmatrix} \widetilde{\text{MR1}}_{t-1} \\ \widetilde{\text{MR2}}_{t-1} \end{bmatrix} \\ + \begin{bmatrix} 0.36 & -0.14 & -0.02 \\ 0.10 & 0.30 & -0.04 \\ 0.07 & 0.23 & 0.38 \end{bmatrix} \begin{bmatrix} \Delta^2 \text{GR}_{t-1} \\ \Delta^2 \text{WA}_{t-1} \\ \Delta^2 \text{EA}_{t-1} \end{bmatrix} + \text{Deterministic terms} + \begin{bmatrix} \epsilon_{\text{GR},t} \\ \epsilon_{\text{WA},t} \\ \epsilon_{\text{EA},t} \end{bmatrix}$$

# I(2) Cointegrated VAR Model

Test for and estimate I(2) CVAR model from Juselius (2006):

$$\begin{bmatrix} \Delta^2 \text{GR}_t \\ \Delta^2 \text{WA}_t \\ \Delta^2 \text{EA}_t \end{bmatrix} = \begin{bmatrix} -0.007 & 0 \\ -0.003 & -0.005 \\ 0.01 & -0.01 \end{bmatrix} \begin{bmatrix} \widetilde{\text{LR1}}_{t-1} \\ \widetilde{\text{LR2}}_{t-1} \end{bmatrix} + \begin{bmatrix} 0.07 & 0.02 \\ 0.007 & 0.21 \\ 0.18 & 0.30 \end{bmatrix} \begin{bmatrix} \widetilde{\text{MR1}}_{t-1} \\ \widetilde{\text{MR2}}_{t-1} \end{bmatrix} \\ + \begin{bmatrix} 0.36 & -0.14 & -0.02 \\ 0.10 & 0.30 & -0.04 \\ 0.07 & 0.23 & 0.38 \end{bmatrix} \begin{bmatrix} \Delta^2 \text{GR}_{t-1} \\ \Delta^2 \text{WA}_{t-1} \\ \Delta^2 \text{EA}_{t-1} \end{bmatrix} + \text{Deterministic terms} + \begin{bmatrix} \epsilon_{\text{GR},t} \\ \epsilon_{\text{WA},t} \\ \epsilon_{\text{EA},t} \end{bmatrix}$$

Medium-run relationships:

Bi-polar Relation,  $\widetilde{\text{MR1}}_t : \Delta \text{WA}_t = 0.4 \Delta \text{GR}_t - 4$

Antarctic Relation,  $\widetilde{\text{MR2}}_t : \Delta \text{EA}_t = -\Delta \text{WA}_t - 7$

# I(2) Cointegrated VAR Model

Test for and estimate I(2) CVAR model from Juselius (2006):

$$\begin{bmatrix} \Delta^2 \text{GR}_t \\ \Delta^2 \text{WA}_t \\ \Delta^2 \text{EA}_t \end{bmatrix} = \begin{bmatrix} -0.007 & 0 \\ -0.003 & -0.005 \\ 0.01 & -0.01 \end{bmatrix} \begin{bmatrix} \widetilde{\text{LR1}}_{t-1} \\ \widetilde{\text{LR2}}_{t-1} \end{bmatrix} + \begin{bmatrix} 0.07 & 0.02 \\ 0.007 & 0.21 \\ 0.18 & 0.30 \end{bmatrix} \begin{bmatrix} \widetilde{\text{MR1}}_{t-1} \\ \widetilde{\text{MR2}}_{t-1} \end{bmatrix} \\ + \begin{bmatrix} 0.36 & -0.14 & -0.02 \\ 0.10 & 0.30 & -0.04 \\ 0.07 & 0.23 & 0.38 \end{bmatrix} \begin{bmatrix} \Delta^2 \text{GR}_{t-1} \\ \Delta^2 \text{WA}_{t-1} \\ \Delta^2 \text{EA}_{t-1} \end{bmatrix} + \text{Deterministic terms} + \begin{bmatrix} \epsilon_{\text{GR},t} \\ \epsilon_{\text{WA},t} \\ \epsilon_{\text{EA},t} \end{bmatrix}$$

Medium-run relationships:

$$\text{Bi-polar Relation, } \widetilde{\text{MR1}}_t : \Delta \text{WA}_t = 0.4 \Delta \text{GR}_t - 4$$

$$\text{Antarctic Relation, } \widetilde{\text{MR2}}_t : \Delta \text{EA}_t = -\Delta \text{WA}_t - 7$$

Long-run relationships:

$$\text{Bi-polar Relation, } \widetilde{\text{LR1}}_t : \text{WA}_t = 0.4 \text{GR}_t - 4t - 15 \Delta \text{WA}_t - 8 \Delta \text{GR}_t + 3$$

$$\text{Antarctic Relation, } \widetilde{\text{LR2}}_t : \text{EA}_t = -\text{WA}_t - 7t - 52 \Delta \text{WA}_t - 43 \Delta \text{EA}_t + 14$$

→ Trend likely a proxy for physical forcing: temperature / ocean warming

→ East Antarctica only affects Greenland indirectly in the long-run

## I(2) Cointegrated VAR Model

Test for and estimate I(2) CVAR model from Juselius (2006):

$$\begin{bmatrix} \Delta^2 \text{GR}_t \\ \Delta^2 \text{WA}_t \\ \Delta^2 \text{EA}_t \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ -0.004 & 0 \\ 0.01 & -0.01 \end{bmatrix} \begin{bmatrix} \widetilde{\text{LR1}}_{t-1} \\ \widetilde{\text{LR2}}_{t-1} \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0.17 & 0.28 \end{bmatrix} \begin{bmatrix} \widetilde{\text{MR1}}_{t-1} \\ \widetilde{\text{MR2}}_{t-1} \end{bmatrix} \\
 + \begin{bmatrix} 0.30 & 0 & 0 \\ 0.12 & 0.30 & 0 \\ 0 & 0.23 & 0.36 \end{bmatrix} \begin{bmatrix} \Delta^2 \text{GR}_{t-1} \\ \Delta^2 \text{WA}_{t-1} \\ \Delta^2 \text{EA}_{t-1} \end{bmatrix} + \text{Deterministic terms} + \begin{bmatrix} \epsilon_{\text{GR},t} \\ \epsilon_{\text{WA},t} \\ \epsilon_{\text{EA},t} \end{bmatrix}$$

Medium-run relationships:

$$\text{Bi-polar Relation, } \widetilde{\text{MR1}}_t : \Delta \text{WA}_t = 0.4 \Delta \text{GR}_t - 4$$

$$\text{Antarctic Relation, } \widetilde{\text{MR2}}_t : \Delta \text{EA}_t = -\Delta \text{WA}_t - 7$$


Long-run relationships:


$$\text{Bi-polar Relation, } \widetilde{\text{LR1}}_t : \text{WA}_t = 0.4 \text{GR}_t - 4t - 15 \Delta \text{WA}_t - 8 \Delta \text{GR}_t + 3$$

$$\text{Antarctic Relation, } \widetilde{\text{LR2}}_t : \text{EA}_t = -\text{WA}_t - 7t - 52 \Delta \text{WA}_t - 43 \Delta \text{EA}_t + 14$$

- Trend likely a proxy for physical forcing: temperature / ocean warming
- East Antarctica only affects Greenland indirectly in the long-run
- Can simplify using automatic model selection (Doornik, 2009)


# Simulated impact of a sudden loss of ice in Greenland






- ▶ Unconditional forecasts perform well for first 1-2 years
- ▶ Track West Antarctica up through 6-years-ahead
- ▶ Is East Antarctica still reacting to the long-run relationships?
- ▶ I(2) and simplified I(2) models are similar at these horizons

## Greenland's Cumulative Contribution to Sea Level Rise by 2100



## Antarctica's Cumulative Contribution to Sea Level Rise by 2100



- ▶ **Greenland:** I(2) in upper tail and Simple I(2) in lower tail of IPCC range
- ▶ **Antarctica:** Both models in lower tail of IPCC range (East Antarctica?)

**THANK YOU**

- Bamber, J. L. and Aspinall, W. (2013). An expert judgement assessment of future sea level rise from the ice sheets. *Nature Climate Change*, 3(4):424–427.
- Broecker, W. S. (1998). Paleocene circulation during the last deglaciation: a bipolar seesaw? *Paleoceanography*, 13(2):119–121.
- Church, J. A., Clark, P. U., Cazenave, A., Gregory, J. M., Jevrejeva, S., Levermann, A., Merrifield, M. A., Milne, G. A., Nerem, R. S., Nunn, P. D., et al. (2013). Sea level change. Technical report, PM Cambridge University Press.
- Doornik, J. A. (2009). Autometrics. In Shephard, N. and Castle, J. L., editors, *The Methodology and Practice of Econometrics: A Festschrift in Honour of David F. Hendry*, chapter 4, pages 88–121. Oxford: Oxford University Press.
- Johansen, S. (1991). Estimation and hypothesis testing of cointegration vectors in gaussian vector autoregressive models. *Econometrica: Journal of the Econometric Society*, pages 1551–1580.
- Johnsen, S., Dansgaard, W., Clausen, H., and Langway, C. C. (1972). Oxygen isotope profiles through the antarctic and greenland ice sheets. *Nature*, 235(5339):429–434.
- Juselius, K. (2006). *The cointegrated VAR model: methodology and applications*. Oxford university press.
- Shepherd, A., Ivins, E. R., Geruo, A., Barletta, V. R., Bentley, M. J., Bettadpur, S., Briggs, K. H., Bromwich, D. H., Forsberg, R., Galin, N., et al. (2012). A reconciled estimate of ice-sheet mass balance. *Science*, 338(6111):1183–1189.
- Stocker, T. F. (1998). The seesaw effect. *Science*, 282(5386):61–62.