

# The Spacecraft Potential's Influence on the FOV of Rosetta-ICA at Low Ion Energies

---

**Sofia Bergman**<sup>(1)</sup>, Gabriella Stenberg Wieser<sup>(1)</sup>, Martin Wieser<sup>(1)</sup>,  
Fredrik Leffe Johansson<sup>(2)</sup> and Anders Eriksson<sup>(2)</sup>

[sofia.bergman@irf.se](mailto:sofia.bergman@irf.se)

(1) Swedish Institute of Space Physics, Kiruna, Sweden

(2) Swedish Institute of Space Physics, Uppsala, Sweden

EGU Sharing Geoscience Online, 4-8 May 2020




INSTITUTET FÖR RYMDFYSIK  
Swedish Institute of Space Physics


# Background

- Spacecraft charging is problematic when doing low-energy ion measurements, since the particles are either attracted to or repelled from the spacecraft prior to detection
- This distorts the effective FOV of the instrument
- The Ion Composition Analyzer (ICA) on board Rosetta was measuring positive ions down to energies of a few eV
- Rosetta was commonly charged to a negative potential of -10 to -20 V, which distorted the low-energy part of the ICA data
- In this study we use the Spacecraft Plasma Interaction Software (SPIS) to model the distortion of the effective FOV of ICA at low energies

Trajectories of high-energy ions entering ICA



Trajectories of low-energy (a few eV) ions entering ICA




[Bergman et al., 2020a]

# ICA Instrument Description

- Mass resolving ion spectrometer
- Positive ions
- Energy range: a few eV/q – 40 keV/q

## Nominal FOV:

- Total FOV of  $360^\circ \times 90^\circ$
- In azimuthal direction: 16 sectors of  $22.5^\circ$
- In elevation: 16 elevation steps of  $5.6^\circ$



[Bergman et al., 2020b]

# SPIS Simulations

- PIC approach to model the environment
- Test particles traced from the instrument
- Plasma model (valid close to perihelion):

$\text{H}_2\text{O}^+$

$e^-$

$n = 1000 \text{ cm}^{-3}$


$T = 0.5 \text{ eV}$

$v = 4 \text{ km/s}$

$n = 1000 \text{ cm}^{-3}$

$T = 8 \text{ eV}$

- → Spacecraft potential of -21 V




[Bergman et al., 2020a]

# Results - Example

Results for one pixel of the instrument  
 $E = 5\text{-}10$  eV (at infinity)


[Bergman et al., 2020a]

Nominal FOV




Effective FOV

Sector 2, Elevation 9, 5-10 eV




- Dashed square: Nominal FOV of the studied pixel
- Color scale: Flux of particles at the simulation boundary that are reaching the studied pixel
- The colored area hence corresponds to the effective FOV of this pixel

# Results – Effect on Different Elevation Angles




- Sector 3, E=5-10 eV
- Extreme elevation angles are much more severely distorted than angles close to the aperture plane




[Bergman et al., 2020a]

# Results – Different Debye Lengths of the Surrounding Plasma



- $U_{s/c} = -21 \text{ V}$  in all cases
- The FOV distortion is more severe when the Debye length is shorter
- The sensitivity to a changing Debye length varies, however, between different instrument pixels






# Results – Varying Spacecraft Potentials

**Question:** Does the FOV distortion scale linearly with the spacecraft potential?

$$U_{s/c} = -21 \text{ V}$$


$$E = 5\text{-}10 \text{ eV}$$



$\neq$


$$U_{s/c} = -42 \text{ V}$$

$$E = 10\text{-}20 \text{ eV}$$



[Bergman et al., 2020b]

- The Debye length is the same in both cases
- If the ion energy is scaled according to the potential of the spacecraft, the FOV distortion is not necessarily the same
- The small nonlinearity observed is mainly caused by the photoemission and bulk flow of the cometary plasma





# Conclusions

[sofia.bergman@irf.se](mailto:sofia.bergman@irf.se)

- The FOV of ICA is distorted at low ion energies
- The distortion varies between different instrument pixels
- The distortion is sensitive to changes in the Debye length, but the sensitivity varies between different instrument pixels
- A small nonlinearity exists in the relation between FOV distortion and spacecraft potential, mainly caused by the photoemission and bulk flow of the cometary plasma
- **The distortion is considered insignificant at an energy above twice the spacecraft potential**

## Publications:

Bergman et al. (2020a). *JGR: Space Physics*. <https://doi.org/10.1029/2019JA027478>

Bergman et al. (2020b). *JGR: Space Physics*. <https://doi.org/10.1029/2020JA027870>



INSTITUTET FÖR RYMDFYSIK  
Swedish Institute of Space Physics

# Publications

Bergman, S., Stenberg Wieser, G., Wieser, M., Johansson, F. L. & Eriksson, A. (2020a). The influence of spacecraft charging on low-energy ion measurements made by RPC-ICA on Rosetta. *Journal of Geophysical Research: Space Physics*, 125. <https://doi.org/10.1029/2019JA027478>

Bergman, S., Stenberg Wieser, G., Wieser, M., Johansson, F. L. & Eriksson, A. (2020b). The influence of varying spacecraft potentials and Debye lengths on in situ low-energy ion measurements. Submitted to *Journal of Geophysical Research: Space Physics*.  
<https://doi.org/10.1029/2020JA027870>

[sofia.bergman@irf.se](mailto:sofia.bergman@irf.se)



INSTITUTET FÖR RYMDFYSIK  
Swedish Institute of Space Physics