



Observations of low energy ions around the diamagnetic cavity of comet 67P

Gabriella Stenberg Wieser, Martin
Wieser, Sofia Bergman, Elias Odelstad,
Fredrik Johansson and Hans Nilsson
Swedish Institute of Space Physics

1

OBSERVING LOW ENERGY IONS WITH RPC-ICA

At comet 67P there is **lots of dynamics** seen in the low energy ions. For the ion mass spectrometer **RPC-ICA** (Ion Composition Analyzer) we implemented a new observational mode to observe this.

The top panel below shows measurements made with the normal mode and the bottom panel measurements with the **high time resolution mode**.

Normal mode

192 s time resolution

90x360° field of view


5 eV-40 keV

High time resolution mode

4 s time resolution

4x360° field of view


5-95 eV



SPACECRAFT POTENTIAL AND HIGH VOLTAGE OFFSET

4 September 2015

Negative of the spacecraft potential observed with LAP assuming LAP picks up 80% of the true potential




The spacecraft potential is often negative **accelerating ions into the instrument**. The low energy cutoff in the observations therefore often corresponds to the negative of the spacecraft potential. The Langmuir probes (**RPC-LAP**) measures (a fraction of) the spacecraft potential and **we match LAP and ICA observations to find the true potential**. The high voltages used in ICA drift with time and depends on temperature. This **high voltage offset can also be estimated** by comparing ICA and LAP observations.

3


IONS 'FAR AWAY' FROM THE SPACECRAFT

4 September 2015

Near



Far away



We are interested in the **plasma properties 'far away' where the ions are not affected by the presence of the spacecraft**. Therefore we remove the acceleration caused by the spacecraft potential. The geometric factor used to convert from observed counts to physical differential flux is also adjusted.


4

IONS JUST OUTSIDE OF THE DIAMAGNETIC CAVITY

Spacecraft potential

4 September 2015

Magnetic field




Typical observations of ions just outside the diamagnetic cavity. The top panel shows the original measurements together with the **spacecraft potential** and the observed **magnetic field**. The bottom panel shows the corrected ion data. The white line indicates how **the energy-integrated ion flux** varies (scaled to fit in the panel).

SPACECRAFT POTENTIAL, B-FIELD AND ION FLUX

4 September 2015

Far away



Zooming in on a few ion flux increases, we can look at **the timing of the potential, the magnetic field and the ion flux.**

We see that the magnetic field follows the spacecraft potential. **The main ion flux peaks** are, however, **lagging** the magnetic field and spacecraft potential peaks.


It is possible to see a **smaller peak** in the ion flux coinciding with the steepest gradient in the spacecraft potential (and the magnetic field).

6

SPACECRAFT POTENTIAL, B-FIELD AND ION FLUX

4 September 2015


Far away



B-field
follows $-Vs/c$

Peak
in $-Vs/c$


Averaging over several ion flux peaks we note that **the time shift is about 20 seconds** from the peak in the spacecraft potential.



ION TEMPERATURE AND ION FLUX

4 September 2015


Far away normalized



The **upper energy limit** of the ion observations is an indication of the ion **temperature**. We normalize the flux to get rid of the widening of the spectrum due to an increase in flux. Some possible temperature increases still remain (panel above).

There is **weak correlation between ion temperature and ion flux** as shown to the right.

Ion flux versus ion temperature



- Variations in observed ions outside the diamagnetic cavity coincide with variations in spacecraft potential and magnetic field
- The ion flux peaks lag the spacecraft potential and the magnetic field by 20 s: what is the reason for this?
 - Can there be a change in the electron temperature?
 - Are we missing some ions, considering the limited field of view?
- There is weak correlation between apparent ion temperature and ion flux.