

A detailed look on the interaction of solar wind helium with Mercury's surface in the laboratory

H. Biber¹, P.S. Szabo¹, N. Jäggi², M. Wallner¹, R. Stadlmayr¹, A. Niggas¹, M.V. Moro³, D. Primetzhofer³, A. Nenning¹, A. Mutzke⁴, M. Sauer¹, J. Fleig¹, A. Foelske-Schmitz¹, K. Mezger², H. Lammer⁵, A. Galli², P. Wurz² and F. Aumayr¹



TU Wien,
Vienna



University of Bern,
Bern



Uppsala University,
Uppsala



Max Planck Institute
for Plasma Physics,
Greifswald



Austrian Academy
of Sciences,
Graz




Image of Mercury: NASA

- Several effects alter the surface of airless bodies, called space weathering [1, 2]
- Thin collision-less exospheres are formed
- Solar wind ions are sputtering but can be implanted and stay in the surface
- Implantation and release of these particles can affect both surface structure and exosphere formation [3, 4]

[1] Hapke B.: J. Geophys. Res. Planet., 106, 10039, 2001.


[2] Noble S.K., et al.: Sol. Syst. Res. 37, 31, 2003.

[3] Sasaki S., et al.: Adv. Space Res., 33, 2152, 2004.

[4] Hartle R., et al.: J. Geophys. Res., 80, 3689, 1975.

Experimental techniques

a) Quartz Crystal Microbalance (QCM) irradiations with solar wind He



- Sample irradiated with 4keV $^4\text{He}^+$ under normal incidence and at room temperature
- Resonance frequency dependent on mass:

$$\frac{\Delta m}{m_0} = -\frac{\Delta f}{f_0}$$

b) Thermal Desorption Spectroscopy subsequent to the irradiations

- Sample can be heated up to 700K
- Constant heating ramps used
- Released ^4He detected via Quadrupole Mass Spectrometer (QMA)
- Mass before and after heating can be determined with the QCM
- Experiments performed under Ultra-high vacuum

Sample preparation


Magnesium rich
Pyroxene



Pulsed Laser
Deposition (PLD)



Target layer on QCM




- Elemental composition and thickness of target deduced from combined Rutherford Backscattering Spectrometry (RBS) and coincidence Time-of-flight/Energy Elastic Recoil Detection Analysis (ToF-E ERDA)
- Flat surfaces are observed via Atomic Force Microscope

	H	C	O	Mg	Si	Ca	Fe
[at. %]	1.4	1.1	60.3	8.9	20.5	1.8	6.0

Tab.:Elemental composition from ERDA

Mass change during irradiation




- Net implantation in the beginning (I)
- Region with oscillation in mass change and enhanced release (II)
- Equilibrium reached after $2.2 \cdot 10^{21}$ ions/m² (III) - some 100 years of solar wind on Mercury [1]
- Assuming constant sputtering, the implanted mass can be calculated

Figure reduced due to Copyright issues, full figure available during EGU:

<https://owncloud.tuwien.ac.at/index.php/s/xJ2oyQsyNfy8LeY>

[1] Wurz P., et al.: Planet. Space Sci., 58, 1599, 2010.

Subsequent Thermal Desorption Spectroscopy




- Peak in release at about 450K
- Mass loss due to heating agrees with previously implanted He mass
- He completely removed below 500K
- Mercury surface temperature is 100-700K, depending on illumination

Figure reduced due to Copyright issues, full figure available during EGU:

<https://owncloud.tuwien.ac.at/index.php/s/xJ2oyQsyNfy8LeY>

[1] Wurz P., et al.: Planet. Space Sci., 58, 1599, 2010.

Comparison with SDTrimSP simulations



- Comparing with SDTrimSP [1]
- Limiting to 10 at.% He reproduces the general trend
- Enabling ion induced transport does so too
- Oscillations can not be explained with this methods

Figure reduced due to Copyright issues, full figure available during EGU:

<https://owncloud.tuwien.ac.at/index.php/s/xJ2oyQsyNfy8LeY>

[1] Mutzke A., et al.: IPP Report, 2019.

- Saturation with Helium is reached after about $2.2 \cdot 10^{21}$ ions/m² equal to some hundreds of years of solar wind on Mercury
- When heating the sample, all He gets removed below 500K
- Mercury surface temperatures range from about 100K (dark) to 700K (illuminated), so possibly fluctuations in He implantation occur
- Oscillation in implantation with enhanced release of implants are observed
- SDTrimSP can reproduce general trend of implantation, but not the oscillations

This work was funded by the Austrian Science Fund FWF (Project No. I 4101-N36) as well as the Swiss National Science Foundation Fund (200021L_182771/1). Support by VR-RFI (contract #821-2012-5144) and the Swedish Foundation for Strategic Research (SSF, contract RIF14-0053) supporting operation of the accelerator at Uppsala University is gratefully acknowledged.