

# Studying the propagation of ELF waves in the Martian ground-ionosphere waveguide using an FDTD method with application to the ground tomography

P. Rzonca, A. Kulak, J. Mlynarczyk

AGH University of Science and Technology, Department of Electronics, Krakow, Poland




This work has been supported by the National Science Center under grant 2015/19/B/ST9/01710.

# Mars ground-ionosphere waveguide and the idea of ground tomography

- Short pulses are generated by dust-devils / dust storms
- Ground surface on Mars is very (extremely) dry
- ELF waves can penetrate ground much deeper than on Earth
- Reconstruction of wave propagation can be useful for development of inverse methods and therefore allow measurements of subsurface properties
- Martian ELF stations are currently under development in our team


# Numerical model

- Time domain simulation
- Cylindrical coordinates centered around the event and scaled via focusing factor
- Large scale grid ( $d\rho = 10\text{km}$ ,  $dz = 1\text{km}$ )
- Measurements at large distances (3 Mm)
- Current source – Short VED with a cutoff at 450 Hz
- Double-“knee” daytime ionosphere profile [Pechony and Price 2004]
- Multilayer ground [Kulak and Mlynarczyk 2013]




# Results

- First layer - small conductivity ( $\sigma_1 = 10^{-6}$  S/m), second layer - high conductivity ( $\sigma_2 = 10^{-2}$  S/m)
- Pulse source with  $p = 1$  C · km dipole moment
- Measured amplitude at the distance of 3 Mm from the source for various depths of the first layer



# Results

- Same environment + highly conducting plane (diameter =  $2r_{plane}$ ) with source located in the center
- Small conducting plane – almost no impact
- Possible resonance on some plane size relative to antenna location



# Conclusions

- Amplitude of the electromagnetic field at large distances from the source strongly depends on the ground structure
- This approach allows measurement of waveforms and enables identification of ground structure and its components.
- Possibilities of further studies:
  - local distortions of ionospheric profile
  - impact of the solar terminator